



Website: contoureducation.com.au | Phone: 1800 888 300

Email: hello@contoureducation.com.au

VCE Biology $\frac{3}{4}$
AOS 1 Revision [1.8]
Workbook Solutions

Outline:



[1.2] - Nucleic Acids & The Structure of Genes

Pg 2-9

- Recap
- Questions

[1.3] - Gene Expression & The trp Operon

Pg 10-21

- Recap
- Questions

[1.4] - Proteins, Protein Export & Enzymes

Pg 22-34

- Recap
- Questions

[1.5] - Introduction to DNA Manipulation Techniques

Pg 35-45

- Recap
- Questions

[1.6] - CRISPR-Cas9 & Bioethics

Pg 46-57

- Recap
- Questions

[1.7] - Recombinant Plasmids

Pg 58-65

- Recap
- Questions

Section A: [1.2] - Nucleic Acids & The Structure of Genes (20 Marks)

Sub-Section: Recap



Cheat Sheet

[1.2.1] - Identify and compare the characteristic features of the structures of nucleic acids and their monomers, including DNA, mRNA, tRNA and rRNA, including base pairing

- Nucleic acids (DNA and RNA) encode instructions for protein synthesis.
- DNA is a double-stranded helix composed of nucleotides containing a deoxyribose sugar, phosphate group, and nitrogenous bases (adenine, thymine, cytosine, guanine).
- Complementary base pairing occurs in DNA with adenine pairing with thymine (A-T) and cytosine pairing with guanine (C-G).
- RNA exists in three main forms—mRNA (messenger RNA), rRNA (ribosomal RNA), and tRNA (transfer RNA)—each with distinct functions in protein synthesis.
- mRNA carries the genetic code transcribed from DNA to the ribosomes for translation.
- rRNA forms the structural and functional core of ribosomes, where proteins are synthesised.
- tRNA transports specific amino acids to the ribosome during protein synthesis, matching them to the mRNA codons via its anticodon.
- Unlike DNA, RNA is single-stranded, contains the sugar ribose, and uses uracil (U) instead of thymine (T).

[1.2.2] - Identify and describe the structure of a nucleotide in DNA and RNA

- A nucleotide contains a phosphate group, a 5 carbon sugar, and a nitrogenous base - label these on the diagram below!
- DNA nucleotides include deoxyribose, whereas RNA nucleotides include ribose; both share phosphate groups but differ in one nitrogenous base (U in RNA, T in DNA).



[1.2.3] - Define the key components of a gene, including a comparison between the structure of genes in eukaryotes and prokaryotes

<u>Component</u>	<u>Definition</u>
Exons	Regions of DNA that are expressed as a protein.
Introns	Regions of DNA that aren't expressed in the final protein.
Promoter	A sequence of DNA that is responsible for initiating gene expression.
Operator	Binding site for a transcription factor (protein that regulates gene expression).
Terminator	Where transcription of the gene ends.

[1.2.4] - Identify and practically apply the characteristics of the genetic code - universal, unambiguous, degenerate, triplet - to real-life examples

➤ Characteristics of Genetic Code:

Universal	All living organisms use DNA or the language of DNA as the fundamental basis for all their operations.
Triplet	3 bases of DNA (or nucleotides) code for a single amino acid in a protein.
Degenerate	More than 1 triplet/codon codes for a single amino acid.

Sub-Section: Questions

INSTRUCTION: 20 Marks. 20 Minutes Writing.



Question 1 (1 mark)

Some viruses are called retroviruses and contain RNA as their blueprint. Once inside the host cell, an enzyme called reverse transcriptase converts the RNA into DNA. An RNA retrovirus sequence is shown below.

3' GAUCAGUCCA 5'

What would the DNA code be for this sequence?


- A. 3' CTAGTCAGGT 5'
5' GATCAGTCCA 3'
- B. 5' CTAGTCAGGT 3'
3' GATCAGTCCA 5'
- C. 3' CATGACTGGA 5'
5' GTACTGACCT 3'
- D. 3' CUAGUCAGGU 5'
5' GAUCAGUCCA 3'

Reverse transcriptase will convert the RNA strand in the stem of the question into a complimentary DNA strand. The DNA strand will be anti-parallel to the RNA strand, so it will be 5' CTAGTCAGGT 3'. DNA is double-stranded and so the complimentary DNA strand to the one that was reverse transcribed would be 3' GATCAGTCCA 5'.

Space for Personal Notes

Question 2 (1 mark)

The diagram below shows a small segment of the structure of an organic macromolecule.



Which of the following statements correctly identifies what occurs when this segment is joined to other similar structures?


- A. A net output of energy occurs.
- B. The reaction is catabolic.
- C. A condensation reaction would occur.
- D. An input of water is required for this reaction to occur.

A is incorrect – a process that involves the assembly of a large molecule from smaller monomers is anabolic, not catabolic.
 B is incorrect – anabolic reactions are also endergonic where an input of energy is required to form chemical bonds between monomers.
 C is correct – biological macromolecules are produced as a result of condensation reactions.
 D is incorrect – reactions that require an input of water are called hydrolysis reactions. Water is released when the bonds between monomers are broken.

Space for

The following information applies to the two questions that follow.

The diagram below represents part of a DNA molecule.



Question 3 (1 mark)

A single DNA nucleotide is shown by sub-unit(s):

- A. X alone.
- B. X and Y together.
- C. Y and Z together.
- D. X, Y and Z together.**

Question 4 (1 mark)

A feature of DNA that can be seen in the diagram above is

- A. the anti-parallel arrangement of the two strands of nucleotides.**
- B. the process of semi-conservative replication.
- C. its ribose sugar-phosphate backbone.
- D. its double-helix structure.

Space for Personal Notes

Question 5 (1 mark)

In eukaryotic cells, this biomacromolecule would be located in the


- A. nucleus only.
- B. nucleus and mitochondria.
- C. nucleus and chloroplasts.

The endosymbiotic theory suggests that ancestral aerobic bacteria and ancestral cyanobacteria may have been consumed (endocytosis) by ancestral eukaryotic cells. These structures provided a survival advantage to the consumer (efficient production of ATP as well as production of glucose) and the consumed bacteria were provided protection. One piece of evidence to support this theory is that there is DNA in both mitochondria and chloroplasts. This DNA (virtually) works independently to the nuclear DNA.

D. nucleus, chloroplast and mitochondria.

Question 6 (5 marks)

- a. In the space below, draw and label the generic structure of a nucleotide. (1 mark)



- b. Name two types of nucleic acids and the differences between them. (2 marks)

DNA	RNA
double-stranded	single-stranded
deoxyribose sugar	ribose sugar
thymine	uracil

- c. Briefly explain why it is possible to exchange DNA molecules between different species. (2 marks)


DNA is universal; it is the same nucleotides in all species.

The same codons code for the same amino acids irrespective of what species it is.

Question 7 (10 marks)

RNA molecules consist of long strands of joined nucleotides. Each nucleotide consists of three sub-units.

- a. Label the three sub-units on the diagram of the RNA nucleotide below. (1 mark)



- b. Complete the table below by describing the role in a cell of the two types of RNA listed. (2 marks)

Marks	0	1	2	Average
%	41	29	30	0.9

<u>Type of RNA</u>	<u>Role in a cell</u>
tRNA	Brings an amino acid to the ribosome. OR Complementary anticodon binds with codon of mRNA.
mRNA	Carries genetic information from nucleus to the ribosomes. OR Complementary copy of the DNA.

- c. Describe the process in which both of these RNA molecules are involved and state its end product. (4 marks)

Translation:

- mRNA molecule binds to the ribosome at the 5' end.
- tRNA anticodons complementary to the mRNA codons, delivering specific amino acids in their correct order.
- adjacent amino acids are joined together by condensation polymerisation by the ribosome
- translation that ends when the stop codon is reached.
- A polypeptide chain that becomes the basis of a protein is the end product.

- d. What feature of the genetic code allows the same amino acid to be produced despite a change in the DNA? (1 mark)

The degeneracy of the genetic code, as multiple codons can code for the same amino acid.

- e. List two differences between DNA and RNA. (2 marks)


DNA is double stranded whereas RNA is single stranded. Dexoyribose vs ribose sugar.

Section B: [1.3] - Gene Expression & The trp Operon (35 Marks)

Sub-Section: Recap

Cheat Sheet

[1.3.1] Identify and recall the process of gene expression in eukaryotes, comparing how it differs in prokaryotes



- Transcription and translation occur simultaneously in prokaryotes, and there is no mRNA processing.

[1.3.2] Describe the processes of transcription, mRNA processing, and translation, recognising the significance of each step to the final product and [1.3.3] Explain how a single gene can give rise to multiple proteins

- RNA polymerase binds to the promoter region and the DNA unwinds and unzips.
- RNA polymerase then catalyses the production of the mRNA strand by joining together complementary RNA nucleotides. mRNA is complementary to the DNA template strand - adenine pairs with uracil in RNA instead.
- Continues until a termination or stop sequence is reached.
- **Addition of Methyl 5' Cap:**
 - Added to the 5' end of the mRNA strand.
 - Prevents degradation.
 - Enables the ribosome to detect the RNA.

➤ Addition of Poly-A Tail:

- Added to the 3' end of the mRNA strand.
- Composed of adenine bases.
- Prevents degradation and increases mRNA stability.

➤ Splicing: Exons and Introns:

- Splicing removes **introns** (non-coding regions) and retains **exons** (coding regions) in the mRNA strand.
- This process is performed by a complex molecule called a spliceosome.

➤ Alternative Splicing:

- Allows flexibility in mRNA processing:
 - Exons can be removed, introns can be retained, or the order of exons can be shuffled.
- Significance:
 - Creates multiple mRNA variants from a single gene.
 - Leads to the production of diverse proteins from the same gene.

➤ Translation:

- mRNA molecule binds to the ribosome at the 5' end.
- tRNA anticodons complementary to the mRNA codons, delivering specific amino acids in their correct order to the ribosome.
- Adjacent amino acids are joined together by condensation polymerisation by the ribosome.
- Translation ends when the stop codon is reached.

[1.3.4] Identify and recall the general principles and reasons for gene regulation in both prokaryotes and eukaryotes

➤ **Gene Regulation:**

- ⚙ Ensures specific genes are expressed only when required, conserving energy and resources.
- ⚙ **Structural genes:** Code for proteins that perform specific functions (e.g., enzymes).
- ⚙ **Regulatory genes:** Code for transcription factors that control the expression of structural genes by interacting with DNA sequences like promoters or operators.

➤ **Purpose of Gene Regulation:**

- ⚙ Avoids unnecessary production of proteins that may not be needed.
- ⚙ Responds to environmental changes (e.g., nutrient availability, stress).
- ⚙ Maintains cellular efficiency and resource management.

➤ **What is an Operon?**

- ⚙ A cluster of genes with related functions, controlled by a single promoter.
- ⚙ Includes a promoter, an operator (binding site for regulatory proteins), and structural genes.
- ⚙ Example: The *trp operon* in *E. coli* codes for enzymes that synthesise the amino acid tryptophan.

➤ **Benefits of Operons:**

- ⚙ Simplifies regulation of multiple related genes.
- ⚙ Allows a coordinated response to environmental changes (e.g., nutrient levels).
- ⚙ Common in prokaryotes but absent in eukaryotes.

[1.3.5] Describe the regulation of the trp operon through the action of the repressor protein

- A regulatory gene upstream of the operon produces the trp repressor protein.

➤ When tryptophan levels are high:

- ⚙ Tryptophan binds to the repressor protein, causing a conformational change that activates it.
- ⚙ The active repressor binds to the operator, blocking RNA polymerase from transcribing the structural genes.
- ⚙ This halts the production of enzymes needed for tryptophan synthesis, conserving resources.

➤ When tryptophan levels are low:

- ⚙ Tryptophan dissociates from the repressor, rendering it inactive.
- ⚙ RNA polymerase can bind to the promoter and transcribe the structural genes.
- ⚙ Enzymes for tryptophan synthesis are produced, allowing the cell to make more tryptophan.

➤ **Importance of Repressor Dissociation:**

- ⚙ Prevents unnecessary repression when tryptophan levels drop.
- ⚙ Ensures enzymes are available to synthesise tryptophan when needed.

[1.3.6] Describe the regulation of the trp operon through attenuation in high trp environments

- Unique to prokaryotes because transcription and translation occur simultaneously.
- **Mechanism:**
 - 🔗 The leader sequence between the operator and structural genes contains codons for two tryptophan residues and a region that can form hairpin loops in the mRNA.
 - 🔗 Depending on tryptophan availability, the mRNA forms one of two types of loops:
 - **Low tryptophan levels:**
 - 🔗 Ribosome stalls at leader sequence → Anti-terminator loop forms → RNA polymerase continues transcription → Structural genes are expressed.
 - **High tryptophan levels:**
 - 🔗 Ribosome does not stall → Attenuator loop forms → RNA polymerase is stopped → Structural genes are not expressed.


Sub-Section: Questions

INSTRUCTION: 35 Marks. 35 Minutes Writing.

Check off any learning objectives that obtained full marks from the "Contour Check" booklet!

Question 8 (1 mark)

The diagram below illustrates an important cellular process. Labels *M-Q* relate to various steps and/or chemicals involved in the process.



Which one of the following correctly names the steps/chemicals represented by *M-Q*?

- A. *M*: translation; *N*: mRNA; *O*: pre-mRNA; *P*: protein; *Q*: transcription
- B. *M*: transcription; *N*: pre-mRNA; *O*: mRNA; *P*: protein; *Q*: translation**
- C. *M*: translation; *N*: pre-mRNA; *O*: mRNA; *P*: protein; *Q*: transcription
- D. *M*: transcription; *N*: mRNA; *O*: pre-mRNA; *P*: protein; *Q*: translation

Protein synthesis has been heavily assessed over the past few years, particularly in the short-answer section. A clear understanding of the steps and chemicals involved is required. Initially the DNA must be transcribed (step *M*) and the pre-mRNA (chemical *N*) needs to be modified. This modification removes introns, and the smaller mRNA (chemical *O*) is comprised of only exons. The mRNA leaves the nucleus and is fed through a ribosome to be translated (step *Q*) into a protein (chemical *P*).

Question 9 (1 mark)

The graph shows transcription rates of the **trp operon** measured as mRNA synthesis (arbitrary units) under varying tryptophan concentrations (μM).

Tryptophan Concentration (μM)	Transcription Rate (mRNA Units)
0	100
10	80
50	40
100	10
200	0

Based on the data, what pattern is observed in transcription rates of the trp operon as tryptophan levels increase?

- A. Transcription rates increase steadily due to the need for tryptophan synthesis.
- B. Transcription rates remain constant, regardless of tryptophan levels.
- C. Transcription rates decrease sharply as the repressor binds to the operator region.
- D. Transcription rates oscillate unpredictably, depending on ribosome activity.

Explanation: As tryptophan levels rise, transcription decreases sharply, consistent with repressor protein activity at high tryptophan concentrations.

Question 10 (1 mark)

In a prokaryote, what is the primary benefit of regulating gene expression using operons?

- A. It prevents unnecessary proteins from being synthesised, conserving energy and resources.
- B. It allows eukaryotic-like control of mRNA stability.
- C. It eliminates the need for transcription factors in gene expression.
- D. It increases the efficiency of post-translational modifications.

Space for Personal Notes

Question 11 (1 mark)

What would occur if a mutation prevented the trp repressor from binding tryptophan?

- A. The operon would remain permanently off, regardless of tryptophan levels.
- B. Transcription of the trp operon would continue even when tryptophan levels are high.
- C. RNA polymerase would fail to bind the promoter region of the operon.
- D. Tryptophan biosynthesis would halt, even at low tryptophan concentrations.

Question 12 (1 mark)

How does attenuation regulate the trp operon under high tryptophan levels?

- A. The ribosome quickly translates the leader sequence, forming a terminator structure in the mRNA.
- B. The repressor protein binds to the operator, blocking RNA polymerase.
- C. The ribosome stalls, allowing transcription to continue.
- D. Tryptophan molecules bind to RNA polymerase, halting elongation.

Solution Pending

Space for Personal Notes

Question 13 (11 marks)

Collagen is a very important protein that is generally found in the extracellular environment of our bodies (the space between our cells).

- a.** Name the first step of gene expression and describe this process with reference to making collagen. (4 marks)

Transcription:

- RNA polymerase binds to the promoter region of the collagen gene and the DNA unwinds and unzips.
- RNA polymerase then catalyses the production of the mRNA strand by joining together complementary RNA nucleotides. mRNA is complementary to the DNA template strand – adenine pairs with uracil in RNA instead.
- Continues until a termination or stop sequence is reached, with this pre-mRNA coding for collagen moving on to the next step in gene expression.

b. Describe two events that occur during RNA processing and describe their significance. (4 marks)

- Addition of 5' Methyl G cap – allows the mRNA to be recognised by the ribosome to initiate translation, whilst also ensuring stability as it exits the nucleus.
- 3'Poly A-tail – prevents the degradation of the mRNA chain at the 3' end as it leaves the nuclear membrane.
- Splicing – removal of non-coding regions (introns) to generate mature mRNA.
- Alternative splicing - alternative splicing, exons can shuffle around and introns can be retained, thus resulting in many possible mRNA products, producing many different proteins from a single gene.

c. How can a single gene give rise to multiple proteins? (3 marks)

A single gene can code for multiple different proteins due to the RNA processing occurring in gene expression. Specifically, during alternative splicing, exons can shuffle around and introns can be retained, thus resulting in many possible mRNA products, producing many different proteins from a single gene.

Question 14 (14 marks)


- a. Describe the difference between a structural gene and a regulatory gene. (2 marks)

Structural genes are those which have a protein product that contributes to a structure/function within an organism, whereas regulatory genes code for transcription factors and repressors which seek to control the rate of expression of other genes (structural genes).

- b. Why is gene regulation so important for organisms to be able to do? (2 marks)

- Allows them to control what proteins are made and in what quantities – no energy wasted on useless proteins.
- This also enables specialisation of cells, as they can be regulated to produce specific proteins above others.

The trp operon is one such process that is useful for bacteria to control the production of tryptophan, an amino acid.



c. What is an operon? Why might this be advantageous? (2 marks)

Solution Pending

d. Explain, through the function of the trp repressor, how the level of tryptophan in the cell regulates the expression of the trp operon. (4 marks)

- The regulatory gene upstream of the operon will code for the production of the trp repressor protein.
- When there is a sufficient level of trp in the cell, it will bind to the repressor protein. This causes a conformational shape change in the repressor, therefore allowing it to bind to the operator region of the operon.
- This blocks RNA polymerase from transcribing the genes, resulting in the enzymes responsible for trp synthesis not being produced.
- When there is a shortage of trp however, the repressor will not have trp binding to it, and will therefore not undergo the conformational shape change required to bind to the operator. In this case, RNA polymerase can bind to the promoter and travel along the gene unimpeded, allowing transcription to occur. Hence, the enzymes required for trp synthesis will be produced, allowing the cell to produce trp on its own.

Another method by which the trp operon can be regulated is through attenuation.

e. Explain, using the principle of attenuation, how trp operon can be regulated post-transcriptionally. (4 marks)

- In prokaryotes, transcription and translation can occur simultaneously due to a lack of processing and a nucleus.
- In between the operator and the structural genes, there is a leader sequence.
- This leader sequence contains the code for two trp residues. If there is low levels of tryptophan, then the ribosome will be stalled while translating this sequence.
- Following the leader sequence there is an attenuator – if the ribosome is stalled than this attenuator will form an anti terminator loop and allow transcription to occur.
- If there are high levels of trp then the ribosome will not be stalled – thus resulting in a terminator loop to be formed, preventing full transcription of the structural genes.

Space for Personal Notes

Question 15 (5 marks)

- a. Name the enzyme that is responsible for catalysing transcription. (1 mark)

RNA polymerase.

- b. Name the product of transcription, and describe the processing it undergoes before exiting the nucleus. (3 marks)

Pre-mRNA.

RNA Processing:

- 5' Cap, poly A 3' tail.
- Removal of introns.
- Alternative splicing.

- c. What feature of the genetic code allows for this protein to be produced by a bacterial cell? (1 mark)

The universality of the genetic code.

Space for Personal Notes

Section C: [1.4] - Proteins, Protein Export & Enzymes (41 Marks)

Sub-Section: Recap



Cheat Sheet



[1.4.1] - Define and compare primary, secondary, tertiary and quaternary structures of proteins

- Proteins are made of amino acids joined by condensation polymerisation.
- Proteins have diverse functions determined by their structure.
- The proteome is the full set of proteins expressed by a cell, tissue, or organism.
- Amino acids have an amine group, a carboxyl group, and an R group.
- The R group determines the chemical properties of the amino acid (non-polar, polar, positive, or negative).
- **Primary structure:** The sequence of amino acids in a polypeptide chain, determining the order of R groups and the foundation for higher levels of structure.
- **Secondary structure:** The folding of the polypeptide chain into alpha-helices or beta-pleated sheets, stabilised by hydrogen bonds between non-adjacent amino acids. These structures increase stability and provide mechanical strength.
- **Tertiary structure:** The overall 3D shape of a polypeptide, formed by interactions between R groups, including hydrophobic interactions, hydrogen bonds, ionic bonds, and disulfide bridges. This structure often determines the protein's specific function.
- **Quaternary structure:** The combination of multiple polypeptide chains into a single functional protein complex. Not all proteins have a quaternary structure.
- Protein diversity comes from their shape and post-transcriptional modifications.
- More proteins exist than genes due to alternative splicing and modifications.

[1.4.2] - Identify and describe the roles of ribosomes, rough endoplasmic reticulum and golgi apparatus in the transport and export of proteins from a cell

- Ribosome - site of protein synthesis (amino acids joined together to form a polypeptide chain)
- Rough ER - Site of the initial folding of the polypeptide chain, packaged into transport vesicles for delivery to the Golgi apparatus
- Golgi - Its function is to modify and package the protein into a vesicle for transport to various locations within the cell, including to the cell membrane for secretion.
- Transport vesicles take proteins/polypeptides from the rough endoplasmic reticulum to the Golgi Apparatus, and secretory vesicles take the finished product for exocytosis at the membrane.

[1.4.3] - Explain the lock-and-key model, the induced fit model and why enzymes are specific and only catalyse one reaction.

- Enzymes act as organic catalysts, speeding up chemical reactions that would take much longer to occur naturally.
- They are proteins with a 3D active site that is complementary to the shape of their specific substrate.
- Coenzymes assist enzymes by altering the shape of the active site or by providing necessary electrons and protons for the reaction.
- **Process of enzyme function:**
 1. Substrate binds to the enzyme's active site.
 2. The enzyme changes its shape slightly to fit the substrate better (induced fit).
 3. The reaction occurs, transforming the substrate into products.
 4. The products are released, and the enzyme is ready for another reaction.
- **Lock and Key Model:** The enzyme's active site and substrate fit together perfectly, like a key fitting into a lock.
- **Induced Fit Model:** The enzyme's active site moulds itself to fit the substrate upon binding, like a glove fitting a hand.

[1.4.4] - Explain how enzymes change/denature in different pH and at different temperatures

- **Temperature** - As the temperature increases, the rate of enzyme function will INCREASE (as there is an increase in the frequency of successful collisions), until the optimal temperature is reached, after this,
- **pH** - If the pH is within the optimal pH range of an enzyme, the rate of enzyme function will be maximal (as this allows for the greatest frequency of successful collisions). When the pH is outside of this optimal functioning pH range of an enzyme, the rate of enzyme function will decrease (as the active site of the enzyme will be denatured, allowing for less successful collisions).

[1.4.5] - Explain the function of competitive and non-competitive enzyme inhibitors, how they affect the rate of reaction, and how they may/may not be overcome

- **Competitive Inhibitor** - A competitive inhibitor will reduce the activity of an enzyme, as it will compete with the substrate for the active site. This can be overcome by increasing substrate concentration.
- **Non-competitive inhibitor** - binds to the allosteric site of an enzyme, causing a conformational shape change of the active site, preventing the substrate from binding and reducing the activity.

Sub-Section: Questions

INSTRUCTION: 41 Marks. 41 Minutes Writing.

Check off any learning objectives that obtained full marks from the "Contour Check" booklet!

Question 16 (1 mark)

What would be the consequence of a mutation that alters the R-groups of amino acids in a protein's active site?

- A. Loss of secondary structure stability due to disrupted hydrogen bonds.
- B. Decreased specificity of enzyme-substrate binding.**
- C. Misfolding of the protein at the primary structure level.
- D. Disruption of quaternary structure due to lack of subunit interactions.

Question 17 (1 mark)

Which best explains how protein diversity can exceed the number of genes in an organism?

- A. The primary sequence of a protein determines its folding, which increases diversity.
- B. Alternative splicing and post-translational modifications create multiple proteins from one gene.**
- C. The same amino acid sequence can fold into different shapes under environmental conditions.
- D. Quaternary structure allows a single polypeptide chain to function as multiple proteins.

Space for Personal Notes

Question 18 (1 mark)

Why does the tertiary structure of a protein determine its specific function?

- A. It directly influences how subunits interact in a quaternary structure.
- B. It determines the arrangement of disulphide bridges in the protein backbone.
- C. It creates a specific three-dimensional active site that interacts with substrates or other molecules.
- D. It sets the sequence of amino acids for the protein's primary structure.

Question 19 (1 mark)

What happens to an enzyme that is exposed to a highly acidic pH for an extended period?

- A. The enzyme is denatured due to the breaking of peptide bonds in its primary structure.
- B. The enzyme's tertiary structure is disrupted, altering the shape of its active site.
- C. The enzyme's secondary structure is replaced by random coil regions.
- D. The enzyme becomes inactive but regains function when the pH is neutralised.

Question 20 (1 mark)

How do competitive and non-competitive inhibitors differ in their effects on enzyme activity?

- A. Competitive inhibitors reduce the enzyme's V_{max} , while non-competitive inhibitors increase K_m .
- B. Competitive inhibitors block substrate binding, while non-competitive inhibitors alter enzyme conformation.
- C. Non-competitive inhibitors only act on enzymes with quaternary structure, while competitive inhibitors act on all enzymes.
- D. Competitive inhibitors bind allosterically, while non-competitive inhibitors compete with the substrate for the active site.

Space for Personal Notes

Question 21 (1 mark)

Which cellular organelle is primarily responsible for the modification and packaging of proteins for extracellular export?

- A. Rough endoplasmic reticulum.
- B. Golgi apparatus.**
- C. Lysosome.
- D. Ribosome.

Question 22 (1 mark)

Which factor most directly determines whether a protein exhibits quaternary structure?

- A. The presence of disulphide bridges within a single polypeptide chain.
- B. Interaction between multiple polypeptide chains or subunits.**
- C. Hydrogen bonding within alpha-helices and beta-sheets.
- D. The sequence of amino acids in the primary structure.

Question 23 (1 mark)

What role do enzymes play in chemical reactions, and how is this achieved?

- A. Enzymes increase reaction rates by breaking substrate bonds through hydrolysis.
- B. Enzymes lower the activation energy required for the reaction to proceed.**
- C. Enzymes create energy to drive non-spontaneous reactions.
- D. Enzymes permanently bind to substrates to stabilise intermediate complexes.

Space for Personal Notes

Question 24 (1 mark)

How does the proteome differ from the genome of an organism?

- A. The genome determines cellular function, while the proteome determines gene expression.
- B. The genome contains instructions for proteins, while the proteome represents proteins actively expressed at a given time.**
- C. The proteome contains all RNA transcripts, while the genome contains all DNA sequences.
- D. The genome includes only coding regions, while the proteome includes post-translational modifications.

Question 25 (1 mark)

Which of the following best describes why heat permanently denatures an enzyme?

- A. It disrupts hydrogen bonds in the active site, rendering it non-functional.**
- B. It breaks peptide bonds, leading to loss of primary structure.
- C. It alters the enzyme's ionic bonds without affecting the hydrophobic interactions.
- D. It induces changes in the secondary structure, which the enzyme cannot repair.


Space for Personal Notes

Question 26 (21 marks)

The liver plays an important role in metabolising and breaking down drugs that we consume and enter our body. This ranges from medical drugs, which are important in treating diseases, but also involves other socially acceptable drugs such as alcohol.

The pathway to breaking down these drugs can often be very complex and involves key molecules, called enzymes found within the liver, and if these drugs are left unmetabolised, it may cause issues for the body in general.

- a. Using diagrams, describe the two models that describe enzyme function and describe their differences. (4 marks)



The active site doesn't alter to fit the shape of the substrate in the lock and key model, whereas it does slightly in the induced fit model.

When large quantities of these drugs are consumed, this may impact their metabolism in the liver, and after a certain point, there may be excess drug being left unmetabolised, which in the case of alcohol impacts the body's wider function (sensation of being 'drunk').

- b.** Identifying a factor that impacts the rate of enzyme function, explain why after consuming a lot of alcohol, someone might feel drunk. (2 marks)

A factor that may impact the rate of function in this case would be the concentration of the alcohol.

Increasing concentration would increase the rate of metabolism, however, once all active sites have reached saturation, the rate would plateau, with more alcohol being left unmetabolised and therefore contributing to that feeling of being drunk.

- c.** Identify and describe 2 other factors that impact enzyme function. (4 marks)

➤ Temperature – increasing rate, until it gets too high and breaks down the structure (denaturation). Rate slows at lower temperatures but does NOT denature.

➤ pH – rate is highest when in the optimal pH range, outside the pH range the enzyme may denature thus reducing its effectiveness and function.

The enzyme that is involved primarily with the metabolism of many drugs by the liver is the CYP450 enzymes. A recommendation listed on many medications is that you should not drink grapefruit juice prior to or immediately after consuming these drugs.

- d. How might grapefruit juice impact the CYP450 enzymes and their ability to metabolise medical drugs? (2 marks)

Grapefruit juice is considered to be an inhibitor of the CYP450 enzymes, and this could be either competitive or non competitive inhibition. Ultimately, the grapefruit juice would prevent the substrate from binding the active site, resulting in a reduction in the ability to metabolise medical drugs.

Collagen has to be exported outside of the cell to reach the extracellular spaces.

- e. Place the structures from which it goes through in order, after the nucleus, and describe their function in relation to protein export. (8 marks)

Ribosome	Synthesises the polypeptide chain from mRNA via translation.
Rough Endoplasmic Reticulum	Responsible for folding, modifications to the polypeptide, as well as its transport to the golgi apparatus.
Golgi Apparatus	Golgi Apparatus – responsible for adding modification to protein structure (sugar groups for glycoproteins) and packaging them into secretory vesicles for export via exocytosis.
Secretory Vesicle	Travels to the membrane, responsible for exporting it outside of the cell via exocytosis.

f. Define the tertiary structure of a protein and its importance. (1 mark)

The tertiary structure of a protein is the overall 3D structure of the polypeptide, as developed by the interactions between the R groups on amino acids. This is significant as the tertiary structure generally is responsible for the function of a protein.

Space for Personal Notes

Question 27 (10 marks)

A group of students wanted to investigate the activity of an enzyme that catalyses the breakdown of hydrogen peroxide into water and oxygen.

The students measured oxygen concentration using an oxygen sensor. The oxygen sensor fits into the top of a conical flask, as shown in the photograph below.



The students set up three conical flasks with the contents listed in the table below.

Flask	Contents of flask
1	150 mL of 3% hydrogen peroxide solution. 2 mL of enzyme solution. 50 mL of pH-neutral distilled water.
2	50 mL of 3% hydrogen peroxide solution. 2 mL of enzyme solution. 50 mL of high pH buffer solution.
3	150 mL of 3% hydrogen peroxide solution. 2 mL of enzyme solution. 50 mL of low pH buffer solution.


The buffer solutions and the distilled water did not react with the hydrogen peroxide. All three conical flasks were at room temperature. The students recorded the concentration of the oxygen over a five-minute period.

a. State a hypothesis that the students could be testing with this experiment. (1 mark)

The following are examples of possible answers:

- If the pH of the solution is changed, then the rate of the production of oxygen is altered.
- Higher pHs increase the rate of production of oxygen OR a low pH does not result in the production of oxygen.

b. The results of the experiment are shown below.



Explain the difference in the results that the students obtained for the three different flasks. (3 marks)

In Flask 1 the pH is neutral, the enzyme is able to work efficiently, and hydrogen peroxide is being broken down to water and oxygen at the fastest/optimal rate.

Adding 50 mL of alkaline buffer solution will increase the pH of the solution. This change in pH may denature some of the enzyme molecules, or conditions may not be optimal for the enzyme's operation and therefore the rate of the reaction will be slowed down, as seen by the lower percentage of oxygen produced.

Adding 50 mL of acidic buffer solution to the solution has a greater effect on the activity of the enzyme. The pH of the solution may have been altered so that most of the enzyme molecules were denatured and the reaction is very slow or does not proceed.

- c. After completing the experiment, the students decided they could improve their experimental design by adding another conical flask.

Flask	Contents of flask
4	150 mL of 3% hydrogen peroxide solution. 2 mL of distilled water.

- i. Explain how this additional flask could improve their experimental design. (2 marks)

To determine the percentage of oxygen produced without enzyme present and then to act as a comparison for all other results.

- ii. Draw on the graph below the results that would be expected for the production of oxygen in Flask 4. (1 mark)

Students should have drawn a line that was the same as that for Flask 3. The line should not have gone below the initial percentage of oxygen.

- d. Suggest one further improvement that could be made to the students' experimental design. Justify your suggestion. (1 mark)

- Measuring the pH of the solution indicates how much the 50 mL of buffer solution changed the pH.
- Run the experiment for a longer amount of time to check if the percentage of oxygen in the flask plateaus.
- Conduct the experiment over a range of pHs to determine optimal pH for enzyme activity.
- Repeat the experiment.

- e. Consider any experiment. Explain the difference between the accuracy and precision of measurements. (2 marks)

Accuracy of measurement is how close the measurement is to the true value.
Precision of measurement is how close each repeated measurement is to the other measurements.

Section D:[1.5] - Introduction to DNA Manipulation Techniques (28 Marks)

Sub-Section: Recap

Cheat Sheet

[1.5.1] Identify and describe the function of polymerases, endonucleases, and ligases in DNA manipulation

- Polymerases:
 - 🔗 Copy nucleic acid strands by synthesising DNA using an existing template, requiring a primer and helicase to unzip the DNA.
 - 🔗 DNA replication is semi-conservative, producing molecules with one original and one new strand.
- Endonucleases:
 - 🔗 Cut DNA at specific recognition sites, creating either sticky ends (overhangs) or blunt ends (straight cuts).
 - 🔗 Sticky ends are preferred due to complementary base pairing for easier fragment joining (e.g., EcoRI, AluI).
- Ligases:
 - 🔗 Join DNA fragments by reforming phosphodiester bonds, sealing both sticky and blunt ends to create continuous DNA strands.

[1.5.2] Identify the ingredients required, describe the process, and recall key applications of PCR

- The materials required for PCR include:
 - 🔗 **DNA sample:** Provides the template to produce copies.
 - 🔗 **Primers:**
 - **Forward primer** binds to the start of the DNA segment.
 - **Reverse primer** binds to the complementary strand at the opposite end. These primers designate the sequence to be copied and provide a starting point for DNA synthesis.

- 🔗 **Taq polymerase:** A heat-resistant enzyme that makes multiple copies of the DNA strand by adding nucleotides.
- 🔗 **Free nucleotides (dNTPs):** Added by Taq polymerase to produce the new DNA strand.
- 🔗 **Mix buffer:** Maintains the appropriate pH and provides the required salts for Taq polymerase activity.
- 🔗 **PCR tube:** Serves as the vessel for the reaction, containing all the components.

➤ The process of PCR involves the following steps:

1. **Denaturing (90-95°C):** The two DNA strands are broken apart, performing the role that helicase would in a natural setting. Hydrogen bonds between the complementary bases are broken, separating the strands.
2. **Annealing (50-55°C):** Lowering the temperature keeps the strands separate while allowing primers to anneal to the DNA strands. The primers bind at the 3' end, allowing polymerase to begin replication.
3. **Elongation (72°C):** The raised temperature allows a heat-resistant polymerase (Taq polymerase) to function optimally. It extends the primers, replicating the DNA until the end of the sample or a termination sequence. 72°C is the optimal temperature for Taq polymerase.

- These steps are repeated many times (typically 20-40 cycles) to generate a large sample of DNA.

[1.5.3] Describe the process of gel electrophoresis, and describe how it may be used to differentiate DNA samples or to obtain a "DNA profile"

- Gel electrophoresis is used to analyse DNA samples by sorting and separating DNA fragments based on size.
- Typically performed after DNA has been cut with a restriction enzyme.
- Allows for size comparison of DNA fragments, visualisation using fluorescent dyes, and separation of DNA based on fragment size.
- **Why DNA moves through the gel:**
 - DNA is negatively charged due to its phosphate backbone.
 - When an electric current is applied, DNA moves toward the positive electrode.
- **Process:**
 1. DNA samples are placed in wells in an **agarose gel**.
 - One well contains a **standard ladder** with fragments of known sizes.
 2. The gel is placed in a **buffer solution** that conducts electricity.
 - A negative electrode is at the well end, and a positive electrode is at the opposite end.
 3. An electric current is applied, and DNA fragments move through the gel:
 - Smaller fragments move faster.
 - Larger fragments move slower.
 4. A dye (commonly **ethidium bromide**) is added to visualise the results under UV light.
 - DNA fragments appear as fluorescent bands.
- **Importance of standard ladders:**
 - Contain DNA fragments of known sizes.
 - Used to compare and estimate the size of unknown DNA fragments.
 - Specific to experimental conditions for accurate results.

[1.5.4] Explain the factors that affect the movement of fragments in gel electrophoresis

- **Voltage:** Higher voltage increases movement but may distort separation.
- **Gel Composition:** Denser gels slow larger fragments.
- **Buffer Concentration:** Affects electrical conductivity and DNA movement.
- **Time:** Longer runs allow DNA to travel further, but excessive time may cause DNA to move out of the gel.

[1.5.5] Define satellite DNA and STRs, and explain their use in identifying people through DNA profiling for crimes and paternity testing

➤ **Overview:**

- DNA profiling combines technologies like **PCR** and **gel electrophoresis** to identify individuals using DNA.
- Widely used in forensic science to solve crimes and in genetic testing to establish relationships.

➤ **Process:**

- **DNA Collection:** DNA is collected from a crime scene (e.g., blood, hair, or saliva) and compared to samples from suspects or victims.

- **PCR Amplification:**

- The collected DNA is amplified using **PCR** to produce multiple copies of specific DNA regions, particularly **short tandem repeats (STRs)**.
- STRs are short, repeated DNA sequences found in **non-coding regions (satellite DNA)**, which vary significantly between individuals.

- **Digestion with restriction enzyme:**

- DNA is cut with the **same restriction enzyme** to ensure consistent fragment sizes across all samples.

- **Gel electrophoresis:**


- The cut DNA fragments are separated based on size using gel electrophoresis.
- STRs create fragments of varying lengths unique to each individual.
- Fragments are visualised as fluorescent bands under UV light using dyes like ethidium bromide.

➤ **Why STRs are used:**

- STRs are highly variable between individuals, even within non-coding regions.
- Testing multiple STR loci ensures reliable identification and reduces the chance of matching unrelated individuals.

➤ **Interpreting results:**

- Samples with identical banding patterns indicate a match.
- For forensic cases:
 - Matching the crime scene sample to a suspect's STR profile can link them to the scene.
- For genetic testing:
 - Banding patterns confirm familial relationships (e.g., paternity tests).
 - Children inherit STR fragments from both parents, which can be observed in shared patterns.



Sub-Section: Questions

INSTRUCTION: 28 Marks. 28 Minutes Writing.



Question 28 (1 mark)

A DNA fragment with a negative charge is placed in a gel for electrophoresis. What causes its migration?

- A. The attraction to the negatively charged cathode.
- B. The attraction to the positively charged anode.**
- C. The smaller size of the DNA fragment compared to the gel pores.
- D. The application of heat to the gel during the process.

Question 29 (1 mark)

Why are sticky ends often more efficient than blunt ends in creating recombinant DNA?

- A. Sticky ends form hydrogen bonds that make ligation more stable.**
- B. Sticky ends are longer, allowing more opportunities for alignment.
- C. Sticky ends prevent DNA degradation during ligation.
- D. Sticky ends are recognised by ligase without the need for additional enzymes.

Question 30 (1 mark)

A PCR reaction contains the following components: DNA template, Taq polymerase, dNTPs, and primers. What is the purpose of the primers?

- A. To terminate DNA synthesis at specific sequences.
- B. To provide a starting point for Taq polymerase to extend the DNA.**
- C. To unwind the DNA helix during denaturation.
- D. To separate DNA strands during the annealing phase.

Question 31 (1 mark)

A gel electrophoresis experiment shows bands for a crime scene sample matching bands from a suspect's DNA. What does this indicate?

- A. The suspect's DNA has different sequences than the crime scene DNA.
- B. The suspect's DNA matches the crime scene sample, supporting involvement.**
- C. The suspect's DNA contains fewer restriction sites than the crime scene DNA.
- D. The suspect's DNA has undergone more mutations than the crime scene DNA.

Question 32 (1 mark)

What effect would increase the annealing temperature in PCR have on the reaction?

- A. Amplification of non-specific products due to less stringent primer binding.
- B. Increased specificity as primers bind more accurately to the target DNA.
- C. Reduced yield as primers fail to bind effectively to the DNA.**
- D. Decreased denaturation efficiency, leading to incomplete strand separation.

Space for Personal Notes

Question 33 (12 marks)

With the development of DNA manipulation technologies in the 20th century, forensic science has seen significant improvements in its ability to catch the perpetrators of crimes, with even a single hair or cell left at a crime scene proving enough.

- a. What technology can be used to ensure that even trace amounts of DNA can be analysed by technicians? Describe the process. (4 marks)

PCR:

- Denaturation – heating the solution to 90-95 degrees, to separate the two DNA strands.
- Annealing – cool slightly to 50-55 degrees, allowing the primers to bind to each strand of DNA, without them coming together.
- Elongation – reheat this to 70-72 degree, allowing the optimal function of Taq polymerase to extend DNA.
- Repeat these cycles until enough DNA has been amplified.

Once they have enough of a sample, researchers can then apply an endonuclease to a range of DNA samples, and then run the fragments through gel electrophoresis and compare their findings.

- b. How does doing this allow the researchers to tell DNA samples apart and match them to a criminal? (3 marks)

Each individual will have different DNA.
Applying the same restriction enzyme will mean, fragments of different sizes are generated – due to the fact that restriction sites will be found in different places as the sequences are different.
This will result in a different band pattern for each person, allowing them to be told apart.

c. Which property of DNA explains why it will move through the gel? (1 mark)

DNA is negatively charged and hence will move when a current is applied to the gel.

d. Discuss three factors that affect the migration of fragments through the agarose during gel electrophoresis. (3 marks)

Suitable responses for factors affecting the migration of DNA fragments through the agarose gel during gel electrophoresis included three of the following:

- The size of the molecules, as the larger molecule will move more slowly.
- The charge of the molecule, as the negative charge means that DNA moves towards the positive electrode.
- The length of time the voltage is applied, as there may not be enough time for the DNA to migrate through the gel.
- The concentration of the agarose, as denser agarose results in the molecules moving more slowly.
- Students needed to discuss each factor rather than just list them.

e. What is the function of DNA ligase? (1 mark)

To join fragments of DNA together by reconnecting their sugar phosphate backbone via phosphodiester bonds.

Space for Personal Notes

Question 34 (4 marks)

DNA includes sections that are called short tandem repeats (STR). Mutations in STRs occur, on average, every 500 generations. Different numbers of these repeats have no obvious effect on the individual.

- a. What is the likely reason for this? (1 mark)


STRs are found in introns which are non-coding sections of DNA.

A young man, Ben, wants to find out more about his genetic ancestry. He sends a sample of cells, obtained from a swab of his mouth to a laboratory. On receipt of the sample, the laboratory treats the cells to release the DNA to enable the identification of STR markers.

- b. Name the process used to produce many copies of the STR markers. (1 mark)

Polymerase chain reaction.

Each of the STR markers produced is labelled with a dye and subjected to gel electrophoresis. Five of Ben's STR markers were compared with three family groups who have the same surname as him. The following gels resulted.



c. Explain which family is Ben's most recent common ancestor. (2 marks)

Family Y is Ben's most recent common ancestor as there are more (four out of five) STR markers in common between Ben and Family Y than the other families.

Space for Personal Notes

Question 35 (7 marks)

Scientists investigating a transgenic strain of the *Arabidopsis* plant called Kojak carried out a gel electrophoresis to find the root hair gene that had been transferred into the Kojak strain from a species of barley.

The root hair gene was removed along with some other DNA using restriction enzymes and underwent PCR prior to the gel electrophoresis being run.

- a. Why was PCR performed on the DNA sample prior to the gel electrophoresis being carried out? (1 mark)

PCR is performed to amplify the DNA segment so that there is enough to be visible in the gel electrophoresis.

Mark allocation: 1 mark

- 1 mark for recognising that PCR is performed to increase the amount of DNA in the sample

- b. Outline the three major steps in PCR. (3 marks)

Worked solution

1. Heat the DNA to 90 °C
2. Cool to attach primers
3. Taq polymerase copies DNA strands


Mark allocation: 3 marks

- 1 mark for identifying that the DNA must be heated to 90 °C
- 1 mark for stating 'cool to attach primers' or equivalent
- 1 mark for stating 'Taq polymerase copies DNA strands' or equivalent

The table below gives the sizes of the various DNA fragments (genes) used in the genetic engineering of the *Arabidopsis* plant.

Gene	Size (Base Pairs)	Well (Gel Electrophoresis)
Normal <i>Arabidopsis</i> root hair gene	520	2
Mutant Kojak <i>Arabidopsis</i> gene	450	3
Barley root hair gene	600	4
Recombined mutant <i>Arabidopsis</i> + barley genes	1050	5

- c. On the diagram below, indicate where each of the DNA fragments would be positioned after the gel electrophoresis has been run. (2 marks)



- d. What is placed in well 1? (1 mark)

Worked solution

Standard DNA fragments of known lengths

Mark allocation: 1 mark

- 1 mark for identifying a standard DNA fragments of known lengths

Space for Personal Notes

Section E: [1.6] - CRISPR-Cas9 & Bioethics (25 Marks)

Sub-Section: Recap

Cheat Sheet

[1.6.1] Describe the function & process of CRISPR-Cas9 as an adaptive defence against viruses in bacteria

➤ Overview:

- CRISPR-Cas9 is a revolutionary tool derived from bacteria's natural defence mechanism against viruses (bacteriophages).
- It stands for **Clustered Regularly Interspaced Short Palindromic Repeats**.
- Function: Allows for precise and efficient gene editing by targeting specific DNA sequences.

➤ Analogy:

- It works like a surveillance system: bacteria "store" viral DNA to recognise and neutralise future invasions.

➤ Key Features:

- Cas9: A programmable enzyme (endonuclease) that cuts DNA.
- Guide RNA (gRNA): Guides Cas9 to the specific target DNA via complementary base pairing.
- PAM (Protospacer Adjacent Motif): A short DNA sequence that enables Cas9 to bind and cut at the correct site.

➤ Process:

- **Exposure:** A bacteriophage injects its DNA into a bacterial cell.
 - Endonucleases cut a protospacer (a small viral DNA fragment) adjacent to a PAM sequence.
- **Incorporation:** The protospacer is integrated into the bacterial genome.

- **Expression:** The spacer is transcribed into crRNA and combines with tracrRNA to form guide RNA (gRNA).

- **Extermination:** gRNA directs Cas9 to recognise and cut complementary viral DNA, inactivating the virus.

[1.6.2] Explain the process of using CRISPR-Cas9 as a gene editing tool, including silencing, knock-ins & knock-outs

- CRISPR-Cas9 has been adapted as a powerful and versatile tool for editing genomes in eukaryotic cells.
- The system uses the principles of bacterial immunity but has been optimised to make precise cuts in specific DNA sequences within any organism.

➤ How it Works:

➤ Target Sequence Identification:

- A target DNA sequence is selected for editing, such as a gene responsible for a disease.
- Researchers design a synthetic **single guide RNA (sgRNA)**, which combines the targeting properties of crRNA and tracrRNA into one molecule.

➤ CRISPR-Cas9 Complex Formation:

- sgRNA binds to the Cas9 enzyme, forming the CRISPR-Cas9 complex.

➤ DNA Binding:

- The complex locates the target DNA by matching the sgRNA sequence to the complementary DNA sequence.
- A **PAM sequence** (e.g., NGG) adjacent to the target ensures proper binding and initiates the cutting process.

DNA Cutting:

- Cas9 creates a double-strand break at the target site.

Repair Mechanisms:

- The cell's natural repair mechanisms are activated:

-  **Non-Homologous End Joining (NHEJ):** Repairs the break but often introduces errors (e.g., insertions or deletions), which can disable the gene (gene knock-out).

-  **Homology-Directed Repair (HDR):** Uses a provided DNA template to repair the break precisely, allowing the insertion of new genetic material (gene knock-in).

Gene Silencing (Knock-Outs):

- Cas9 introduces a double-strand break at a target DNA sequence.
- The cell repairs the break via non-homologous end joining (NHEJ), which often introduces mutations, disabling the gene.
- Example: Disabling genes linked to diseases like cancer or Huntington's disease.

Gene Knock-Ins:

- Homology-directed repair (HDR) uses an introduced template to insert new DNA sequences.
- Example: Adding a functional copy of a defective gene in conditions like cystic fibrosis.

Targeted Gene Modifications:

- Enables precise alterations to DNA to correct errors or enhance traits.

[1.6.3] Describe & compare the function of the PAM sequence in bacteria & gene editing applications of CRISPR-Cas9 technology

- PAM ensures that Cas9 binds to and cuts only the foreign DNA, distinguishing self from non-self DNA. It also ensures that the Cas9 will only bind to DNA and unwind it for detecting a match where it is possible.
- This system is highly specific and efficient, making it adaptable for gene editing.
- In gene editing applications, it allows the system to be more precise and efficient.

[1.6.4] Describe the function & compare the guide RNA (gRNA) & single guide RNA (sgRNA)

- Guide RNA (gRNA) is a combination of crRNA (CRISPR RNA) and tracrRNA (trans-activating crRNA), which directs Cas9 to the target DNA by complementary base pairing.
- Single guide RNA (sgRNA) is a synthetic RNA that combines the functions of crRNA and tracrRNA into one molecule for simplicity and efficiency.
- Both gRNA and sgRNA guide Cas9 to specific DNA sequences for cutting.
- The sgRNA simplifies the CRISPR-Cas9 system by eliminating the need for two separate RNA molecules.

[1.6.5] Apply bioethical principles to the use of CRISPR-Cas9 technology

➤ **Beneficence:**

- Obligation to use CRISPR to maximise benefits, such as curing genetic diseases and improving crop productivity.

➤ **Non-Maleficence:**

- Avoid causing harm, considering unknown long-term effects or off-target mutations that could negatively impact health or the environment.

➤ **Consent:**

- Ethical concerns arise in germline editing where future generations cannot provide consent. Ensuring parents and patients are fully informed is critical.

➤ **Justice:**

- CRISPR's potential to widen socio-economic inequalities, with only wealthy individuals having access to enhancements, must be addressed.

[1.6.6] Define & describe the bioethical concepts of integrity, respect, beneficence, non-maleficence & justice as elaborated in the VCAA study design

➤ **Integrity:**

- Definition: A commitment to honesty and accurate reporting.
- Ensures transparency and trust in scientific research.
- Example: Publishing both positive and negative results from CRISPR experiments.

➤ **Respect:**

- Definition: Recognising the intrinsic value and autonomy of living things.
- Considers welfare, cultural beliefs, and personal rights.
- Example: Gaining informed consent before experimenting on human subjects.

➤ **Beneficence:**

- Definition: Maximising benefits while minimising risks and harms.
- Encourages actions that contribute positively to individuals or society.
- Example: Using CRISPR to cure genetic disorders responsibly.

➤ **Non-Maleficence:**

- Definition: Avoiding harm, or ensuring harm is proportionate to the benefits.
- Balances risk and reward in scientific practices.
- Example: Carefully evaluating potential off-target effects of CRISPR edits.

➤ **Justice:**

- Definition: Ensuring fair treatment, distribution of benefits, and avoidance of undue burdens.
- Focus on equity and accessibility.
- Example: Avoiding socio-economic inequalities in access to CRISPR-based therapies.

[1.6.7] Define & describe the three ethical approaches as elaborated in the VCAA study design

➤ **Consequences-Based Approach:**

- Central focus: Outcomes or consequences of an action.
- Goal: Maximise positive effects and minimise harm.
- Example: Supporting CRISPR use for disease eradication if the societal benefits outweigh the risks.
- Critique: Can justify morally questionable actions if the outcomes are favourable.

➤ **Duty-/Rule-Based Approach:**

- Central focus: Adherence to moral rules or duties, regardless of outcomes.
- Emphasis on ethical responsibilities and strict regulations.
- Example: Requiring full regulatory approval before using CRISPR on embryos.
- Critique: May ignore potential benefits if rules are overly restrictive.

➤ **Virtues-Based Approach:**

- Central focus: The moral character and intentions of the individual performing the action.
- Emphasis on qualities such as honesty, fairness, and humility.
- Example: Scientists exercising restraint and prioritising ethical considerations in CRISPR applications.
- Critique: Relies on subjective interpretations of virtue.

[1.6.8] Describe briefly how to genetically modify organisms to increase crop productivity & disease resistance, using CRISPR-Cas9

➤ **Improving Productivity:**

- Modifications enhance growth, photosynthesis efficiency, or drought resistance.
- Example: Engineering crops to survive arid climates.

➤ **Enhancing Disease Resistance:**

- Targeted edits make crops resistant to diseases or pests.
- Example: Bananas engineered to resist Panama disease.

➤ **Increasing Nutritional Value:**

- Edits improve the nutritional profile of crops.
- Example: Golden rice enriched with Vitamin A.

[1.6.9] Compare transgenic, cisgenic, & genetically modified organisms

➤ **Genetically Modified Organisms (GMOs):**


- Organisms whose genomes have been altered artificially.
- Includes any form of genetic modification, whether involving foreign or native genes.

➤ **Transgenic Organisms:**

- A subset of GMOs where genes from an unrelated species are introduced.
- Example: Adding genes from fish to tomatoes for cold resistance.

➤ **Cisgenic Organisms:**

- Involves transferring genes from the same or closely related species.
- Example: Enhancing disease resistance in crops using native plant genes.



Sub-Section: Questions

INSTRUCTION: 25 Marks. 25 Minutes Writing.



Question 36 (1 mark)

A genetically modified crop is resistant to pests but leads to a reduction in local insect biodiversity. Which bioethical principle is most relevant?

- A. Beneficence, due to increased crop yields.
- B. Non-maleficence, due to harm caused to insect populations.**
- C. Justice, ensuring equal access to modified crops.
- D. Respect, allowing farmers to choose unmodified crops.

Question 37 (1 mark)

A patient is offered CRISPR treatment for a genetic disease but declines due to personal beliefs. Which ethical concept is being upheld?

- A. Integrity
- B. Non-maleficence
- C. Beneficence
- D. Respect for autonomy**

Space for Personal Notes

Question 38 (1 mark)

CRISPR-Cas9 is used to eliminate genetic diseases in embryos, but this leads to unexpected traits in future generations. Which ethical concern is most relevant?

- A. Beneficence, as this prevents genetic diseases.
- B. Non-maleficence, as an unintended consequence, harms future generations.**
- C. Autonomy, as future individuals cannot consent to the modification.
- D. Justice, as such treatments, may only be available to the wealthy.

Question 39 (1 mark)

A country bans all human germline editing, citing potential misuse. Which ethical principle is the government prioritising?

- A. Justice, to ensure all genetic modifications are regulated equally.
- B. Beneficence, to prevent harm from poorly understood technologies.
- C. Integrity, to align scientific advances with societal values.
- D. Non-maleficence, to avoid long-term unintended effects.**

Question 40 (1 mark)

CRISPR-Cas9 is used to enhance crop yields, but local farmers cannot afford the seeds. Which ethical principle is being violated?

- A. Beneficence, as the crops could alleviate food shortages.
- B. Non-maleficence, as farmers are excluded from economic benefits.
- C. Justice, due to inequity in access to technology.
- D. Autonomy, as farmers cannot choose non-GMO crops.

Solution Pending

Space for Personal Notes

Question 41 (15 marks)

CRISPR-Cas9 is a revolutionary gene editing technique, based upon an immune defence mechanism found in bacteria.

a. Describe how the CRISPR-Cas9 system works as found naturally in bacteria. (3 marks)

- A virus inserts its DNA into the bacteria, and a spacer is cut out and incorporated into the CRISPR locus.
- This is then transcribed to form crRNA which is then combined with tracrRNA to form guideRNA.
- This will form a CRISPR-Cas9 complex with Cas9 enzyme and float around the cell until it encounters complementary viral DNA.
- Viral DNA is cleaved and inactivated, thus protecting the bacteria from infection.

b. What is the significance of the PAM, when looking at the function of CRISPR in bacteria? (2 marks)

The PAM is required for Cas9 to make a cut. So, although the bacterial chromosome has the gRNA match sequence, the lack of PAM ensures that Cas9 won't damage the bacterial chromosome and will instead only cut invading viral DNA which will contain PAM.

CRISPR can be applied to the medical field especially, with scientists figuring out how to apply this technology for gene editing.

c. Describe how CRISPR-Cas9 could be used to edit an embryo and solve this single mutation. (4 marks)

- Identify a target sequence for a cut (in this case the gene with the error for cystic fibrosis), and develop a synthetic guide RNA (sgRNA) complementary to it.
- Combine this with the Cas9 enzyme, altered with a PAM to suit the target.
- Inject this into a target cell (the embryo), and then the sgRNA will bind to the target DNA, and then signal the cut.
- Cell repair mechanisms will try and repair causing errors, which will silence the gene, theoretically preventing cystic fibrosis.

d. What are the advantages of CRISPR-Cas9 that separate it from other techniques? (2 marks)

- CRISPR-Cas9 is much more precise.
- Faster to develop solutions.

- e. Using a consequences-based approach discuss the consequences of germline edits using CRISPR-Cas9. (4 marks)

Consequences based – focusing on the outcomes of an action.

Positive outcomes = Ethical, Negative outcomes = unethical.

- May result in improvements in wellbeing ideally – curing diseases and imperfections.
- There may be unknown impacts and consequences of these germline edits down the line, which may be irreparable.
- Emrbyos, cannot consent to any of the edits made, so them being unhappy with any edits would mean this technology to be unethical.

Space for Personal Notes

Question 42 (5 marks)
Biotech firm says it can resurrect extinct woolly mammoth.

A technology entrepreneur and a geneticist launched a new biotech firm that they say will bring the extinct woolly mammoth back to life. Calling itself Colossal, the biosciences company claims CRISPR genetic technology can be used to bring back the animal, which went extinct over 11,000 years ago.

"Never before has humanity been able to harness the power of this technology to rebuild ecosystems, heal our Earth and preserve its future through the repopulation of extinct animals," technology entrepreneur and Colossal co-founder Ben Lamm said in a statement.

Scientists have managed to find mammoth tusks, bones and other material to try to sequence the animal's DNA. This DNA would then be inserted into the genome of the Asian elephant to form an "elephant-mammoth hybrid," according to the company.

Advocates of "de-extinction" say the process could help humans gain new knowledge regarding biology, evolution and technology. The resurrection of extinct species could also repair damaged ecosystems. In the case of the woolly mammoth, Colossal believes the animal could revitalise the Arctic grasslands, whose properties can mitigate global warming.

The idea of de-extinction could have its drawbacks, however. A March 2017 study published in the Nature Ecology & Evolution journal found that de-extinction programs would be incredibly expensive. Other downsides include resurrected animals carrying new pathogens that could possibly infect humans, along with how they will impact the environment.

The United Nations said in a 2019 report that 1 million animals, plants, and fungi species face extinction in the coming decades.

Source: <<https://learnrgerman.dw.com/en/biotech-firm-says-it-can-resurrect-extinct-woolly-mammoth/a-59171358#>>

- a. With reference to evidence from the article, what ethical approach has been used? (2 marks)

Consequence based approach (1). The consequence of the actions (gain new information, restore ecosystems) outweighs the potential drawback of how it may impact the environment (1).

- b. Suggest a way in which the company proposing the de-extinction of the woolly mammoth could demonstrate the ethical principle of integrity. (1 mark)

The company must accurately report the success and failure of all trials to create a woolly mammoth / elephant hybrid.

- c. Identify **one** potential political issue and **one** potential economic issue associated with the de-extinction of the woolly mammoth. (2 marks)

political: any of the following for 1 mark
Does the habitat cross international borders?
Who is responsible for the monitoring the population?
Do any laws exist for the creation of hybrid species?
Or any other suitable response.
economical: any of the following for 1 mark
Who owns the genome – can they sell it?
Will people lose their homes as habitat is required for the mammoths?
Could the money be better spent?
Or any other suitable response.

Question 43 (2 marks)

The following table provides information on three commonly grown genetically modified (GM) crops in Australia.

Crop	Genetic Modification	Characteristics Given by Modification
GM cotton	Several bacterial genes inserted	Insect resistance and herbicide tolerance
GM canola	Two genes from two different bacterial species inserted	Tolerance to several herbicides
GM safflower	A selection of genes silenced within the safflower genome	Elevated levels of oleic acid in its seeds

- a. Select **one** of the GM crops in the table above and justify whether or not this crop could be described as transgenic. (1 mark)

An acceptable answer was one of the following:

- genetically modified (GM) cotton is transgenic as it contains genes from other species OR bacteria.*
- GM canola is transgenic as it contains genes from other species OR bacteria.*
- GM safflower is not transgenic as it does not contain genes from another species.*

- b. One issue with GM canola is the accidental release, during transport, of seeds along roadsides. Usually, unwanted plants that grow on the side of the road are killed using the herbicide glyphosate. However, GM canola is resistant to glyphosate.

Suggest **one** practical solution for treating GM canola that is found growing along roadsides. (1 mark)

An acceptable answer was one of the following:

- use a different herbicide that the GM canola is not resistant to
- remove by mowing
- pick the GM canola growing by the side of the road by hand
- use controlled burning.

Space for Personal Notes

Section F: [1.7] - Recombinant Plasmids (21 Marks)

Sub-Section: Recap

Cheat Sheet

[1.7.1] Describe the role of plasmids as a vector to transform bacteria & other cells

- Involves combining DNA from multiple sources to produce desired proteins.
- Transformation is the insertion of recombinant DNA into a bacterium, often using a plasmid or vector.
- A vector carries foreign genetic material into a host organism for replication or expression.
- Plasmids, usually carrying the target gene, transform bacteria to produce the desired protein.

➤ **Structure of a plasmid:**

- 🔗 **Plasmid:** Small circular DNA in bacteria, replicates independently, transferable.
- 🔗 **Origin of Replication:** Sequence where replication starts.
- 🔗 **Antibiotic Resistance Gene:** Provides resistance to specific antibiotics.
- 🔗 **Multiple Cloning Site (MCS):** Contains recognition sites for inserting genes.
- 🔗 **Selectable Marker:** Distinguishes transformant from non-transformant cells.
- 🔗 **Screening Marker:** Confirms plasmid is recombinant (e.g., lacZ).
- 🔗 **Promoter Region:** Site where RNA polymerase binds to initiate transcription.

[1.7.2] Explain how a gene of interest is isolated & inserted into a plasmid

➤ **Making a Recombinant Plasmid:**

- 🔗 Obtain a gene of interest and plasmid, cut with the same endonuclease to create sticky ends.
- 🔗 Complementary tick ends will attract via hydrogen bonding.
- 🔗 Use DNA ligase to join the gene and plasmid.
- 🔗 Possible outcomes include recombinant and non-recombinant plasmids.

[1.7.3] Explain electroporation & heat shock as methods to transform bacterial cells

➤ **Transforming Bacteria:**

- 🔗 Electroporation and heat shock methods introduce plasmids into bacteria.
- 🔗 **Electroporation:** Electric current creates membrane pores for DNA entry, increasing permeability.
- 🔗 **Heat Shock:** Uses Ca^{2+} ions, ice, and heat to facilitate DNA uptake, increasing permeability.

[1.7.4] Describe the process by which bacterial cells can be used to produce human proteins, including the application of this to human insulin production

➤ **Isolation of insulin genes:**

- 🧬 mRNA strands coding for the A and B chains of insulin are isolated and used as templates to synthesise complementary DNA (cDNA) via **reverse transcription**. This ensures that introns are not present in the insulin genes.

➤ **Preparation of plasmids and DNA:**

- 🧬 The plasmid (vector) and insulin DNA sequences are cut with the same **restriction endonuclease**, creating complementary **sticky ends**. This ensures the insulin DNA can be inserted into the plasmid.

➤ **Formation of recombinant plasmids:**

- 🧬 The insulin DNA sequences (A and B) are separately inserted into plasmids containing a **beta-galactosidase gene** for detection.
- 🧬 **DNA ligase** is used to join the insulin DNA and plasmid, creating **recombinant plasmids**.

➤ **Introduction to bacteria:**

- 🧬 The recombinant plasmids are inserted into *E. coli* bacterial cells through **heat shock** or electroporation transforming the bacteria.
- 🧬 Each plasmid is introduced into separate bacterial strains to produce the A and B chains independently.

➤ **Selection of transformed bacteria:**

- 🧬 Bacteria are grown on **antibiotic agar plates**, allowing only those with recombinant plasmids (containing antibiotic resistance genes) to survive.
- 🧬 Successful incorporation of the insulin genes is confirmed using **beta-galactosidase activity**.

➤ **Expression and protein extraction:**

- 🧬 Transformed bacteria are cultured, producing **fusion proteins** consisting of beta-galactosidase and insulin chains.
- 🧬 Fusion proteins are extracted, and enzymes are used to **cleave** the insulin chains (A and B) from the beta-galactosidase portion.

➤ **Purification and assembly:**

- 🧬 The insulin A and B chains are purified and then chemically combined.

➤ **Disulfide bonds** are formed between the chains, resulting in functional human insulin ready for medical use.

[1.7.5] Explain the significance of beta-galactosidase, other reporter genes, & antibiotic-resistance genes in the selection of transformed, recombinant bacterial cells

➤ **Selecting Bacteria:**

- 🧬 Identify bacteria with recombinant plasmids using antibiotic selection and reporter genes.
- 🧬 Three groups: without plasmid, with non-recombinant plasmid, and with recombinant plasmid.
- 🧬 Antibiotic resistance indicates transformed bacteria.
- 🧬 Reporter gene function changes indicate recombinant plasmids, often through colour change.
- 🧬 Selected bacteria colonies produce the desired protein for purification.

➤ **Function of b-gal:**

- 🧬 The gene for human insulin will be inserted next to a gene coding for the production of beta-galactosidase protein. This will create a fusion protein, which allows for the detection of successful gene insertion and ensures the survival of the insulin peptide when expressed by the bacteria. Detection of gene insertion is done when the fusion protein allows the plated X-gal substrate to turn blue.


Sub-Section: Questions

INSTRUCTION: 21 Marks. 21 Minutes Writing.



Question 44 (1 mark)

The diagram below is a map of a bacterial plasmid showing ORI, the origin of DNA replication, and selected restriction endonuclease sites.



One plasmid was mixed with the restriction enzymes EcoRI, BamHI and HincII.

Which of the following shows the number of restriction sites that have been cut and the resulting number of DNA fragments produced?

	Number of Restriction Sites Cut	Number of DNA Fragments Produced
A.	3	3
B.	3	4
C.	4	4
D.	4	5

Space for Personal Notes

Question 45 (13 marks)

The process of making insulin using bacteria is given below (some parts simplified). This process avoids the previous need to extract it from animals, commonly pigs.

1. A target gene is isolated and cut using a restriction enzyme.
2. A plasmid with ampicillin resistance and a reporter gene is selected and also cut with the same restriction enzyme.
3. Mixed together with DNA ligase.
4. Using a transformation method, delivered to bacteria.
5. Ampicillin applied to the colony.
6. Bacteria that do not express reporter genes are selected and placed in culture.
7. Insulin collected and purified.

- a.** Describe the action of restriction enzymes and explain which one will you use in this experiment. (3 marks)

They cut DNA. Sticky Ends, they allow for specificity in the insertion of DNA.

- b.** Explain two methods that can be used to insert the plasmid into the bacterium containing the insulin gene. (2 marks)

Heat Shock – involves the solution being included with Ca^{2+} ions, placed on ice. It is then heated to 37-42°C, and then placed back on ice to increase membrane permeability.
Electroporation - involves passing an electric current through the solution containing bacteria and the recombinant plasmids to increase membrane permeability.

c. What is the purpose of treating the bacteria with ampicillin? (2 marks)

The ampicillin will select between the bacteria who have taken up the plasmid or not. The bacteria with the plasmid will have resistance, and therefore will survive, and those that do not have the plasmid will not, and hence will die. Therefore, we can select for bacteria that have been transformed.

d. Describe the function of the reporter gene. (2 marks)

The reporter gene codes for a protein product that is easily identifiable. Given when the recognition sites for the plasmid are, if the gene of interest has been successfully inserted, then the reporter gene will be inactive. Therefore, the presence of the reporter genes product or not, can be used to distinguish and select for transformed bacteria that have taken up a RECOMBINANT plasmid.

e. With reference to both ethical implications and social implications, discuss the advantages of this approach to making insulin above the alternatives. (4 marks)

- The principle of respect – no longer using animals means that they will have their autonomy and welfare respected during the process of making insulin.
- Non maleficence – not only will this confer benefits to using pure human insulin vs animal derivatives, the process of extracting insulin from bacteria produces significantly less harm compared to the animals.
- This process is much more cheaper, allowing lower cost insulin, improving access for many more people.
- Justice – related to the additional point above, but also individuals with religious beliefs will be able to access this with no group unfairly discriminated anymore.

Space for Personal Notes

Question 46 (7 marks)

Gene technology has enabled human insulin to be artificially manufactured within bacteria and then purified for medicinal use by type I diabetics. The steps involved in the procedure from extraction to purification are listed below in the incorrect order.

- Insertion of the modified plasmids into bacteria.
- Amplification of the human insulin gene.
- Isolation of DNA from a human cell.
- Extraction of the insulin protein.
- Selection of the genetically modified bacteria.
- Mixing the restricted plasmid and the restricted human insulin gene together.

- a. Complete the table below by placing the correct step of the insulin purification procedure from the list above in the correct order. (2 marks)

Step 1	Isolation of DNA from a human cell.
Step 2	Amplification of the human insulin gene,
Step 3	Mixing the restricted plasmid and the restricted human insulin gene together.
Step 4	Insertion of the modified plasmids into bacteria.
Step 5	Selection of the genetically modified bacteria.
Step 6	Extraction of the insulin protein.

- b. Discuss how the restricted plasmid and restricted human gene would successfully anneal together when they are combined. (2 marks)

The same restriction enzyme should be used for both the plasmid and the human gene so that they have complementary sticky ends.

When added together with ligase, the complementary sticky ends will have a chance of annealing together with the human gene connected to the plasmid in a ring of DNA.

- c. Use the following terms to discuss the selection of bacteria that have been genetically modified. (2 marks)

bacterial antibiotic resistance gene restriction enzyme binding site

Plasmids (sometimes) contain **bacterial antibiotic resistance genes**, as well as **restriction enzyme binding sites** that are usually separate from each other.

After the uptake of the modified plasmid by bacteria, the bacteria are exposed to the antibiotic the plasmid will confer resistance to. The bacteria surviving are the colonies that have been genetically modified.

- d. Is the manufacture of genetically engineered insulin by the bacteria an example of a genetically modified organism or an example of a transgenic organism? (1 mark)

It is both genetically modified as well as transgenic. The bacteria is genetically modified as it has had its genome altered artificially to achieve a desired outcome. It is also transgenic because DNA from a human has been transferred into the bacterial genome via a genetically altered plasmid.

Space for Personal Notes



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

VCE Biology $\frac{3}{4}$

Free 1-on-1 Support



Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

<u>1-on-1 Video Consults</u>	<u>Text-Based Support</u>
➤ Book via bit.ly/contour-biology-consult-2025 (or QR code below). ➤ One active booking at a time (must attend before booking the next).	➤ Message +61 440 137 387 with questions. ➤ Save the contact as "Contour Biology".

Booking Link for Consults

bit.ly/contour-biology-consult-2025



Number for Text-Based Support

[+61 440 137 387](tel:+61440137387)