# **CONTOUREDUCATION**

Website: contoureducation.com.au | Phone: 1800 888 300

Email: hello@contoureducation.com.au

## VCE Chemistry ½ Models of Atoms [1.1]

Workbook

#### Outline:



#### **Atom Structure**

Pg 2-8

- Atoms Recap
- Rutherford's Gold Foil experiment

#### Bohr's Model of the Atom

Pg 9-26

- Bohr's Key Ideas
- Movement of Electrons Between Energy Levels
- Emission Spectrum
- Electron Configuration
- Shell Model Diagrams

#### Schrödinger's Model of the Atom

Pg 27-39

- Key features of Schrodinger's Model
- Representation of Schrödinger Electron Configurations

#### **Atypical Electron Configurations**

Pg 40-50

- Ionic and Excited State Configurations
- Chromium and Copper
- Condensed Electron Configuration

## **Learning Objectives:**

- CH12 [1.1.1] Describe the Composition of an Atom, and write the Isotopic Symbol of an Element/Ion & use it to identify an Element's/Ion's Atomic and Mass Number
  - mission
- CH12 [1.1.2] Describe Bohr's Model of the Atom & draw Shell Model diagrams & apply Emission Spectra to Bohr's Model of the Atom
- CH12 [1.1.3] Explain Schrodinger's Model of the Atom and identify differences between his Model and Bohr's Model
- CH12 [1.1.4] Write Electron Configurations of Elements and Ions, in both Ground and Excited States, using both Bohr and Schrodinger Models (including Cu and Cr exceptions and Condensed Notation)




## Section A: Atom Structure

## **Sub-Section**: Atoms Recap




#### **Sub-Atomic Particles**

<u>Protons</u>	<u>Neutrons</u>	<u>Electrons</u>
[Positive] / [Negative] / [Neutral]	[Positive] / [Negative] / [Neutral]	[Positive] / [Negative] / [Neutral]
Found in:	Found in:	Found in:
Size:	Size:	Size:



#### **Atoms and Molecules**



- > Atomic Number:
- Mass Number:
- Isotopic Symbol:

 $\frac{A}{Z}E$ 

#### <u>Isotopes</u>

What are the following?




Isotopic Symbol	Worded Representation
<sup>12</sup> <sub>6</sub> C	
13 C	


- Isotope Definition:
  - (same] / [different] number of **protons**
  - (same] / [different] number of **neutrons**


#### <u>lons</u>

Consider Carbon (Atomic Number 6)



- Overall Charge: \_\_\_\_\_
- If the carbon atom gains or loses an electron:





Cations	Anions
[Positive] / [Negative] Charge	[Positive] / [Negative] Charge
[Gains] / [Loses] Electrons	[Gains] / [Loses] Electrons







#### Question 1 Walkthrough.

What is the mass number of Uranium which contains 92 protons and 146 neutrons?

#### Question 2 Walkthrough.

How many protons, neutrons and electrons are in the following atom?

## Try some questions!



#### **Question 3**

Which of the following is not true regarding atomic composition?

- **A.** Atoms are mostly empty space.
- **B.** Electrons are in constant motion around the nucleus.
- C. The nucleus takes up a large portion of an atom's size due to its weight.
- **D.** Electrons are roughly 1800 times lighter than neutrons.

#### **Question 4**

Which of the following is true?

- **A.** An element can be made up of multiple atoms.
- **B.** An element can be made up of more than one type of atom.
- C. An atom can have multiple nuclei, as long as they are identical.
- **D.** F is an example of a molecule.

#### **Question 5**

- **a.** What is the mass number of tin, which contains 50 protons and 70 neutrons?
- **b.** Find the mass number of aluminium, which has an atomic number of 13 and 14 neutrons.

#### **Question 6**

For each of the following, state whether the substance is a/an **atom**, **molecule** or **ion**. If it is an ion, specify whether it is a **cation** or **anion**.

a. Mg

**d.** Cl<sup>-</sup>

**b.** H<sub>2</sub>0

**e.** CO<sub>2</sub>

 $\mathbf{c}$ .  $\mathbf{F}_2$ 

f. Na<sup>+</sup>



#### **Question 7**

For each of the following, state how many **neutrons & electrons** each atom has, and state whether it is a **cation/ anion/ neutral** atom.

a. <sup>19</sup><sub>9</sub> F<sup>-</sup>

**c.**  $^{12}_{6}$  C

**b.**  $^{39}_{19}$  K<sup>+</sup>

**d.** <sup>13</sup><sub>6</sub> C

#### **Question 8 Additional Question.**

Give an example of an atom, element, and molecule made from chlorine Cl.

#### **Question 9 Additional Question.**

State whether  $SO_4^{\ 2^-}$  would be classified as an element, atom, molecule or ion. (Hint: Multiple answers can be correct).

#### **Question 10 Additional Question.**

State the number of protons, electrons and neutrons a carbon atom with a mass number of 13 will have, respectively.

Let's have a look at how this model of the atom was discovered!






### **Sub-Section**: Rutherford's Gold Foil experiment




#### History: Ernest Rutherford's Gold Foil Experiment


Ernest Rutherford was famous for his experiment using a sheet of gold in 1908.



#### **Exploration:** Rutherford's Gold Foil Experiment







- Some of the particles reflected back. Would you expect this result? [Yes]/ [No]
- Most of the particles passed through. Would you expect this result? [Yes]/ [No]
- **Conclusion**: This proves that atoms are primarily made of \_\_\_\_\_\_.





#### Rutherford's Gold Foil Experiment



- Alpha particles were fired at a thin gold sheet.
- Some particles reflected back, but most particles passed through.
- The majority of an atom is empty space.

## Try a question!



#### **Question 11**

Circle the correct alternative from the following:

- **A.** Rutherford's model of the atom explains how electrons exist in shells.
- **B.** Electrons are negatively charged and have the same size as protons.
- **C.** Protons and neutrons are called nucleons because they have similar masses.
- **D.** Neutrons do not have any charge but are found in almost all atoms' nuclei.

#### Key Takeaways



- Atoms are made up of three subatomic particles.
- Protons are positively charged and are found in the nucleus.
- ✓ Neutrons have no charge and are also found within the nucleus.
- Electrons are negatively charged and significantly smaller in size and mass than nucleons.
- Rutherford's Gold Foil experiment saw him firing alpha particles at a thin gold sheet, where some of the particles reflected back, but the majority passed through, proving that atoms are mostly composed of empty space.
- ✓ Atoms are identified by their atomic number.
- Atoms can gain or lose electrons to form ions.



#### **Key Takeaways**



- ✓ The mass number is the number of nucleons present in an atom.
- ✓ Its formula is given by:

## A = Z + number of neutrons

✓ Isotopic Symbol Representation:



- An equal number of protons, but a different number of neutrons.
- ✓ Therefore, different mass numbers.



## Let's have a look at another model of the atom!



## Section B: Bohr's Model of the Atom

<u>Discussion:</u> What did Bohr propose in his model of the atom?



**History:** Bohr's Model of the Atom








## **Sub-Section**: Bohr's Key Ideas



#### **Exploration**: Bohr's Model





- Key Ideas:
  - G Electrons existed at \_\_\_\_\_\_ energy levels (called \_\_\_\_\_\_).
  - G Electrons could \_\_\_\_\_\_ between energy levels.

#### **Electron Shell**



- Definition:
  - A \_\_\_\_\_\_ in the electron cloud of an atom at a \_\_\_\_\_ energy level, whereby an electron can be found.





## **Sub-Section: Movement of Electrons Between Energy Levels**




## How do electrons move between energy levels?



#### **Exploration:** Electron movement between Shells

- What charge does the nucleus have? Positive or negative? (Label Below)
- What charge does an electron have? Positive or negative? (Label Below)
- A/an [attractive]/ [repelling] force acts between the nucleus and the electrons. (Label Below)
- The electrons will want to move [towards]/ [away] from the nucleus.
- Therefore, if an electron was already in the second electron shell, it would move to the [first]/ [third] shell. (Label Below)







#### Analogy: Ball

Think about a ball held in the air which is let go.






Ball



- Which way will the ball go?
- > If we imagine the Earth like an atom:



The first electron shell is called the \_\_\_\_\_\_, as it acts similarly to the ground on the Earth!

#### **Ground State**



- Electrons always try to go the \_\_\_\_\_ energy level possible.
- The first electron shell is known as the ground state.




## How do we get electrons to move up in an electron shell?



Discussion: If we have a ball sitting on the ground, how do we get it to move up in the air?






## <u>Exploration</u>: Electrons moving up energy levels




How to move to another higher energy level from the ground state? (Label Below)





However, what must the electron eventually do? (Label Below)



Discussion: What type of energy is this energy released as?






#### **Analogy**

After eating a lot of sugar, you might be super hyper for a while:



- But eventually, you crash and return to your original energy.
- When energy is inputted, electrons are excited to a [higher]/ [lower] energy level.







#### **Extension:** LED Lighting



A large majority of light sources we see today, use this to emit light, light sources like LEDs and fluorescent lighting all produce light due to electrons going down to a lower energy level.





## Try a question!



Question 12
Outline the 2 major stages that occur when energy is provided to an atom.
<del></del>

#### **Question 13 Additional Question.**

Bohr's model suggests which of the following?

- **A.** Electrons orbit the nucleus randomly within a cloud at continuous energy levels.
- **B.** Electrons exist in shells within the nucleus.
- C. Atoms have shells which are discrete energy levels.
- **D.** Electrons are fixed to a particular shell at all times.

Space for	Personal	Notes
-----------	----------	-------




## **Sub-Section:** Emission Spectrum




## What does the light emitted look like?



#### **Exploration**: Returning back to the Ground State



- Each path the electron takes on the way down has a different \_\_\_\_\_\_.
- This energy difference leads to different \_\_\_\_\_\_ of light being emitted.
- Different elements release \_\_\_\_\_\_ spectra of light when their electrons are excited.





ONTOUREDUCATION
-----------------

VCE Chemistry ½ Questions? Message +61 440 137 304



Each element has a \_\_\_\_\_\_ spectra of light when their electrons return to the ground state of the ground s after being excited.



<u>Discussion:</u> What happens if we input so much energy that the electron is excited past the last energy shell level?



**NOTE**: The energy required to completely remove the first electron from an atom is called its



ALSO NOTE: The idea of first ionisation energy will be covered in the next lesson!

## Try a question for yourself!



#### **Question 14**

State whether the following statements about the electron shell model of the atom are true or false.

	True	False
Electrons can exist between two energy levels.		
When electrons occupy the lowest possible energy level, the atom is in the ground state.		
The energy difference between $n = 2$ and $n = 3$ is greater than the energy difference between $n = 3$ and $n = 4$ .		
Electrons in atoms in excited states return to the next lowest energy level.		
An electron that moves between $n = 5$ and $n = 2$ emits higher energy light than an electron that falls between $n = 4$ and $n = 2$ .		




## **Sub-Section**: Electron Configuration



#### **Context**



We know Bohr's Model looks something like this and can have electrons in each shell.




But how many electrons fit into each energy level?

#### **Exploration**: Maximum number of electrons in each shell



Which shell can hold more electrons between the first and third shell?



- As the electron shell number increases, the maximum number of electrons per shell
   \_\_\_\_\_\_ as the shells become larger.
- > Rule:

## **Maximum Number of Electrons**:



Maximum number of electrons that each shell can hold is:



## Maximum Number of Electrons: $2 \times n^2$

## Try a question!



#### **Question 15**

How many electrons can each of the following shells hold? (Hint: use the  $2n^2$  formula)

a. Shell 1

c. Shell 3

b. Shell 2

d. Shell 4

NOTE: Electron shells are filled in order from the nucleus, with the innermost shells fully filled before moving on.



#### Valence Shell / Valence Electrons



- Definition:
  - Valence Outer.
  - Valence Shell Outer-most electron shell.
  - Valence Electrons Electrons in \_\_\_\_\_\_ electron shell.







#### Question 16 Walkthrough.

Write out the electron configuration for each of the following elements:

- **a.** C (carbon) (6 electrons)
- **b.** Na (sodium)

**NOTE:** Sometimes, we need to look at the **periodic table** to find the number of electrons present.



#### **REMINDER:** Don't forget!

Assuming that the atom has a neutral charge, the number of electrons is equal to the atomic number!

#### **Active Recall**



Shell Number	Maximum Number of Electrons
1	
2	
3	
4	



## Now try a question for yourself!



#### **Question 17**

Write out the electron configuration for each of the following elements:

a. N (Nitrogen)

c. Ar (Argon)

b. S (Sulphur)

d. K (Potassium)

#### **Misconception**



"The electron configuration of elements such as potassium (K) is 2, 8, 9."

#### TRUTH:

- ➤ The \_\_\_\_\_ states the valence shell can hold a maximum of \_\_\_\_ electrons.
- If the electron shell can hold more than 8 electrons, if it is the valence shell, electrons are added to the next electron shell instead.
- As such, potassium (K) has an electron configuration of:

#### **Octet Rule**



- Definition:
  - The valence electron shell can only hold a maximum of \_\_\_\_\_ electrons.



## Try another similar question!



$\sim$	4 •	40
( )11	estion	18

Write out the electron configuration for calcium, which has 20 electrons.

L	
	Space for Personal Notes




## **Sub-Section: Shell Model Diagrams**



#### **Context**

Electron configurations of atoms are represented by **shell model diagrams**.



## Shell model diagrams




They show the number of electrons which are present in each electron shell of an atom.

## Let's have a look at a question together!



#### Question 19 Walkthrough.

Draw the shell model diagram for Calcium (Ca).





### REMINDER: Don't forget!

0

> Pair up the electrons where possible!


## Try some questions for yourself!




#### **Question 20**

Draw the shell model diagram for each of the following elements:

a. Nitrogen (N)



**b.** Chlorine (Cl)







#### Key Takeaways

- ✓ Key Ideas from Bohr's Model:
- ☑ Electrons exist in **discrete** energy levels (called **shells**).
- ☑ Electrons can **move** between energy levels.
- ✓ When energy is inputted, electrons are **excited** to a **higher** energy level.
- Electrons eventually drop back down as they always want to be in the **lowest** energy state whereby energy is released, in the form of **light**, which is unique to each element.
- ☑ Electron shells are filled in order from the nucleus, with the **innermost** shells fully filled before moving on.
- $\checkmark$  Each shell can hold up to  $2n^2$  electrons, where n is the shell number.
- ☑ The octet rule states that the valence electron shell can only hold a maximum of **eight** electrons.
- ☑ Shell model diagrams show the number of electrons within each **shell**.

Space	for	Personal	Notes
Space	101	i Ci Sullai	140163



## Section C: Schrödinger's Model of the Atom

#### **Context**



- Bohr's Model has several shortcomings:
  - igcirc Cannot explain why electron shells can only hold  $2n^2$  electrons.
  - Cannot explain why the **fourth shell accepts two electrons before the third shell** is completely filled (we haven't covered this scenario yet but we'll have a look at it soon!).
- Erwin Schrödinger came up with his new revised model of the atom.


Spa	for Personal Notes



## **Sub-Section**: Key Features of Schrödinger's Model




#### History: Schrödinger's Model of the atom





#### Exploration: Schrödinger's Model vs Bohr's model of the atom

- Main difference:
  - Electrons did not orbit the nucleus like planets but instead had wave-like properties like light.
  - e Electrons merely existed \_\_\_\_\_\_ in a region of space, which was called an



The Bohr model The Schrodinger model

He claimed that each orbital can only hold \_\_\_\_\_\_ electrons.

## Definition

#### **Orbital**

- Definition: A region of space in which electrons exist \_\_\_\_\_\_, not in fixed discrete energy levels.
- Feature: Can hold up to 2 electrons.

## **Exploration**: Subshells

- Electron shells can be broken down into \_\_\_\_\_\_.
- Each subshell has a different \_\_\_\_\_ and orientation in 3D space. These include the:
  - e orbital.
  - e \_\_\_\_\_ orbital.
  - e \_\_\_\_\_ orbital.
  - \_\_\_\_\_ orbital.


**NOTE**: These subshells are quoted in this order as they are arranged in \_\_\_\_\_\_ size.




## Let's have a look at some of these subshells in-depth!

## **CONTOUREDUCATION**

#### **Exploration**: s-orbital







The Schrodinger model

#### Exploration: p-orbital

- The p-orbitals basically look like two blown-up balloons attached to each other.
- $\blacktriangleright$  Aligned with either the x, y or z-axes.



The s and p orbitals fitted together:







**NOTE**: The *z*-axis exists in 3D space, as we live in a 3-dimensional world!




ALSO NOTE: Electrons are assumed to be in constant motion, so their exact location within an orbital is

## Extension: Shape of d-orbitals (not assessed)



 $\blacktriangleright$  d-orbitals start to look a bit more complicated! There are five d-orbitals:



#### **Orbital capacity**



There are four types of orbitals, s, p, d and f.

Type of Orbital	Number of Orbitals
S	
р	
d	
f	

Active Recall: How many	alactrone ca	n fit into	aach af	those.	orbitale?
Active Recall. How illarly	elections ca	וו וונ ווונט	eacii oi	נוובאב י	ni ni raiz:





#### How do orbitals relate to subshells and shells?



#### **Exploration**: Shells, Subshells and Orbitals

- The first electron shell is small, contains \_\_\_\_ -orbitals.
- The second electron shell is larger, contains s-orbitals and \_\_\_\_ -orbitals.
- The third electron shell is even larger, contains s-orbitals, p-orbitals and\_\_\_\_\_\_ -orbitals.
- Conclusion: As the electron shell increases, it can fit more orbitals!

Shell number (n)	Number of subshells	<u>Subshell</u> symbol	Number of Orbitals	Maximum number of electrons in the subshell	Total Number of electrons in the shell
1	1	S			
2	2	S			
	2	р			
3	3	S			
		р			
		d			
4		S			
	4	р			
		d			
		f			

Space for Personal Notes			





#### **Subshells**



- The number of subshells in a given shell matches the shell number.
- For example, shell 3 can hold \_\_\_\_\_ subshells.
- Example:

<u>Subshell</u>	<u>Orbital</u>	<u>Electrons</u>
3s	1	2
3p	3	6
3d	5	10

## Try some questions!



#### **Question 21**

Select the correct alternative from the following:

- **A.** The 1<sup>st</sup> shell can hold s and p orbitals.
- **B.** The d orbital can hold 5 electrons.
- **C.** The p subshell can hold 6 electrons.
- **D.** The 2s subshell has 2 orbitals.





#### **Question 22**

Which of the following is NOT correct regarding the Schrodinger and Bohr Models of the atom?

- **A.** Bohr's model explains the existence of shells.
- **B.** Schrodinger's model states that electrons are in fixed positions within orbitals.
- **C.** Schrodinger's model is the newer one out of the two, explaining more properties of atoms.
- **D.** Bohr's model can be linked to emission spectra of elements.

Question 23 Additional Question.
State how many orbitals and electrons the 5d subshell can hold, respectively:
Orbitals:
Electrons:

Electrons:		
П		
Space for Personal Notes		







How do we write electron configurations according to Schrödinger's model?

## Schrödinger Electron Configurations




> Representation:

#### **Exploration**: Energy Levels of Sub-shells



Each of the subshells has slightly different energy levels.



➤ Electrons are filled up from the \_\_\_\_\_ energy level to the \_\_\_\_\_ energy level.



#### Discussion: 4s and 3d Subshell comparison



- The 4s subshell has a [higher]/[lower] energy level than 3d.
- Therefore, the 4s subshell will be filled [before]/[after] the 3d subshell.

TIP: A good way to remember the hierarchy of energy levels of subshells is through the following:























Each subshell has different energy levels.



For electron configuration, start from the lower energy levels and fill up to higher energy levels.





## Active Recall: The following subshells can hold:



Type of Orbital	Number of Orbitals	Number of Electrons
S	1	
р	3	
d	5	
f	7	

# Let's try a question together!



#### Question 24 Walkthrough.

Write Schrödinger's electronic configuration for Potassium (K).

**REMINDER:** Don't forget the 4s subshell fills **before** the 3d subshell



**NOTE**: The third electron shell starts to fill up while the fourth electron shell remains at one/two electrons!



TIP: Write Schrödinger's electronic configuration using the periodic table!





# Try some questions for yourself!



#### **Question 25**

Write Schrodinger's electronic configuration for each of the following elements:

a. Cl

c. V

**b.** K

d. Fe

NOTE: There are two ways to write 3d and 4s orbitals in electron configurations:



- 4s before 3d.
- 3d before 4s.
- As both are valid, follow what your teacher wants!

**Question 26 Additional Question.** 

Write Schrodinger's electronic configuration for each of the following elements:

**a.** 0


**b.** Al



#### **Key Takeaways**



- An orbital is a region of space in which electrons exist randomly, not in fixed discrete energy levels.
- ☑ Each orbital can hold up to 2 electrons.
- There are 4 different types of orbitals: s, p, d and f, which respectively hold d, d, d electrons.
- lacktriangle Each shell n contains n subshells. For example, shell 2 contains 2 subshells: 2s and 2p.
- Representation:



- When writing out the electron configuration of elements, start from the **lower** energy levels and fill up to higher energy levels.
- ✓ 4s subshell fills before 3d subshell.





# Section D: Atypical Electron Configurations

## **Sub-Section:** Ionic and Excited State Configurations



## **Exploration**: Excited state configurations

- Default electron configurations of atoms in their ground state, meaning that the electrons fill up the lowest energy levels first.
- If an electron is excited, it will go to a [higher]/[lower] energy state.
- Example: Magnesium

$$1s^2$$
,  $2s^2$ ,  $2p^6$ ,  $3s^1$ ,  $6s^1$ 

[ground] / [excited] state

How do we know?

## **Excited state electron configurations**



If electrons are in higher energy levels without lower ones being filled then	Excited state (e.g., $4^{th}$ shell filled before $3^{rd}$ ).
If lower energy subshells are not filled but higher ones are being filled then	Excited state (e.g., 3p orbital is filled but not 3s).







# Try some questions!

Qu	Question 27							
a.	Find the Schrodinger electron configuration of an aluminium ion, Al <sup>3+</sup> .							
b.	The following electron configuration represents an atom or an ion. Is it in an excited state?							
	$1s^22s^22p^63s^1$							
	[Yes] / [No]							
c.	Is $1s^22s^22p^63s^2\ 3p^34s^1$ an excited state configuration?							
	[Yes] / [No]							



## **Sub-Section**: Chromium and Copper



#### **Context**



There are two specific exceptions you need to be aware of to the above rules in Chemistry ½.

**Exploration**: Chromium and Copper



Expected configurations:

Chromium  $1s^22s^22p^63s^23p^63d^44s^2$ 

Copper  $1s^22s^22p^63s^23p^63d^94s^2$ 

Actual Configurations:

Chromium  $1s^22s^22p^63s^23p^6$ 

Copper  $1s^22s^22p^63s^23p^6$ 

- > Result: Gives chromium a \_\_\_\_\_\_ d-subshell and copper a \_\_\_\_\_ d-subshell.
- **Stability**: Partly filled subshells < Half-filled subshells < Fully filled subshells.
- The combination of half-filled/filled subshell ( $3d^5$  and  $4s^1$ ) is **[more] / [less]** stable than one filled subshell and one partly-filled subshell ( $3d^4$  and  $4s^2$ ).

## **Chromium and Copper: Atypical Electron Configurations**



Their Shrodinger electron configurations are:

Chromium  $1s^22s^22p^63s^23p^63d^54s^1$ 

 $Copper\ 1s^22s^22p^63s^23p^63d^{10}4s^1$ 

Due to an increased stability in these forms.





## Practice this idea!



Qu	Question 28									
a.	Write the full Schrödinger's electron configuration for Copper (Cu).									
b.	Rewrite this in Bohr's electron configuration.									

**NOTE:** As an electron from the fourth electron shell jumps back to the third shell, there is only one electron left in the fourth shell.





# **Sub-Section**: Condensed Electron Configuration



The periodic table arranges all of the chemical elements in terms of



#### Periodic table of the elements

1 # 1.0 Hydrogen																	2 He 4.0 Helium
3 Li 6.9 Lithiam	4 Be 9.0 Beryllian		Au Symbol of element B C N 0 F N Relative atomic mass 197.0 10.8 12.0 14.0 16.0 19.0 20												10 Ne 20.2 Neon		
11 Na 23.0 Sodiam	12 Mg 24.3 Magnesiam											13 Al 27.0 Alaminiam	14 Si 28.1 Silicon	15 P 31.0 Phosphorus	16 5 32.1 Sulfur	17 Cl 35.5 Chlorine	18 Ar 39.9 Argon
19 K 39.1 Potassium	20 Cm 40.1 Calcium	21 Sc 45.0 Scandium	22 Ti 47.9 Titanium	23 V 50.9 Vanadium	24 Cr 52.0 Chromium	25 Ma 54.9 Manganese	26 Fe 55.8 Iron	27 Co 58.9 Cobalt	28 Ni 58.7 Nickel	29 Cu 63.5 Copper	30 Zn 65.4 Zinc	31 Ga 69.7 Gallium	32 Ge 72.6 Germanium	33 As 74.9 Arsenic	34 Se 79.0 Seleniam	35 By 79.9 Byomine	36 Kr 83.8 Krypton
37 Rb 85.5 Rubidium	38 Sr 87.6 Strontinm	39 88.9 9Hvium	40 Zv 41.2 Zirconium	41 Nb 929 Niobium	42 Mo 96.0 Molybdenum	43 Tc (98) Technetium	44 Ru 101.1 Ruthenium	45 Rh 1029 Rhodiam	46 PJ 106.4 Palladium	47 Ag 107.9 Silver	48 Cd 112.4 Cadmium	49 In 114.8 Indian	50 Sn 118.7 Tin	51 56 121.8 Antimony	52 Te 127.6 Tellaviam	53   126.9  odine	54 Xe 131.3 Xenon
55 Ca 132.9 Caesiam	56 Ba 137.3 Barium	57-71 Lanthanoids	72 Hr 178.5 Hafninm	73 Ta 180.9 Tantalam	74 W 183.8 Tungsten	75 Re 186.2 Rhenium	76 0s 190.2 Osmium	77  r 192.2  ridium	78 PH 195.1 Platinum	79 Au 197.0 Gold	80 Hg 200.6 Mercary	81 TI 204.4 Thalliam	82 Pb 207.2 Lead	8.3 Bi 209.0 Bismath	84 Po (210) Polonium	85 At (210) Astatine	86 Ra (222) Radon
87 fr (223) Francium	88 Ra (226) Radina	89-103 Actinoids	104 Rf (261) Rutherfordium	105 Db (262) Dubnium	106 Sg (266) Seaborgium	107 Bh (264) Bohrium	108 Hs (267) Hassium	109 M+ (268) Meitneviam	110 Ds (271) Darmstadtiun	111 Rg (272) Roentgenium	112 Cn (285) Coperniciam	113 Nk (280) Nikonium	114 Fl (289) Flerovium	115 Mc (289) Moscovium	116 Lv (292) Livermorium	117 Ts (294) Tennessine	118 Og (294) Oganesson

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La	Ce	Pr	NJ	Pm	Sm	Eq	GJ	Tb	Dy	Ho	Er	Tm	96	Lu
138.9	140.1	140.9	144.2	(145)	150.4	152.0	157.3	158.9	162.5	164.9	167.3	168.9	173.1	175.0
Lanthanum	Cerium	Praseodymium	Neodymium	Promethium	Samariam	Europium	Gadolinium	Terbium	Dysprosium	Holmiam	Erbium	Thalian	9Herbium	Latetium
89	90	91	92	93	94	45	96	97	98	44	100	101	102	103
Ac	Th	Pa	U	Np	Pu	An	Cm	Bk	CF	Es	Fm	MJ	No	Lv
(227)	232.0	231.0	238.0	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(262)
Actinium	Thorium	Protactinium	Uranjum	Neptunium	Plutonium	Americian	Curium	Berkelium	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Laurencium

The value in brackets indicates the mass number of the longest-lived isotope

- There are 7 horizontal rows in the periodic table called \_\_\_\_\_\_\_.
- The **period** number also represents how many \_\_\_\_\_\_ the element has.





## **Exploration**: Condensed Electron Configuration



$$1s^22s^22p^63s^23p^2$$

- Condensed electronic configuration of silicon:
- Examples:
  - Starting with:

$$1s^22s^22p^63s^23p^2$$

[Ne]3s <sup>2</sup> 3p <sup>2</sup>	[Mg]3p <sup>2</sup>
[Correct]/[Incorrect]	[Correct]/[Incorrect]

## **Condensed Electron Configuration**



- **Use**: To 'get rid' of entire shells-worth of notation, for tidiness and efficiency.
- Notation: \_\_\_\_\_ in square brackets, as these are the elements with \_\_\_\_\_ which may be condensed.



# Let's have a look at a question together!



#### Question 29 Walkthrough.

Write the **condensed** electron configuration of the following elements:

a. Boron (B)

**b.** Magnesium (Mg)

# Try some for yourself!



#### **Question 30**

Write the **condensed** electron configuration of the following elements:

a. Lithium (Li)

**b.** Potassium (K)







## **Contour Check**

Learning Objective: [1.1.1] Describe the Composition of an Atom, and write the Isotopic Symbol of an Element/Ion & use it to identify an Element's/Ion's Atomic and Mass Number

#### **Study Design**

the definitions of elements, isotopes and ions, including appropriate notation: atomic number; mass number; and number of protons, neutrons and electrons

Key Takeaways
☐ Atoms are made up of three particles.
Protons arecharged and are found in the
□ Neutrons have charge and are also found within the
□ Electrons are charged and significantly in size and mass than nucleons.
Rutherfold's gold experiment saw him firing <b>alpha particles</b> at a <b>thin gold sheet</b> , where some of the particles <b>reflected</b> back, but the majority, proving that atoms are mostly comprised of
Atoms are identified by their
Atoms can gain or lose electrons to form
☐ The mass number is the number of present in an atom.
□ Isotopic symbol Representation:
<del></del>
Isotopes are two or more of the same elements with number of <b>protons</b> , but number of <b>neutrons</b> , and therefore, mass numbers



# <u>Learning Objective</u>: [1.1.2] Describe Bohr's Model of the Atom & draw Shell Model diagrams & apply Emission Spectra to Bohr's Model of the Atom

#### Study Design

the periodic table as an organisational tool to identify patterns and trends in, and relationships between, the structures (including shell and subshell electronic configurations and atomic radii) and properties (including electronegativity, first ionisation energy, metallic and non-metallic character and reactivity) of elements

	Key Takeaways
Ke	y Ideas from Bohr's Model:
	Electrons exist in energy levels (called)
	Shell model diagrams show the number of electrons within each
	Electrons can between energy levels
	When energy is inputted, electrons are to a energy level.
	Electrons eventually drop back down - as they always want to be in the energy state - whereby energy is released, in the form of which is to each element



# <u>Learning Objective</u>: [1.1.3] Explain Schrodinger's Model of the Atom and identify differences between his Model and Bohr's Model

#### **Study Design**

the periodic table as an organisational tool to identify patterns and trends in, and relationships between, the structures (including shell and subshell electronic configurations and atomic radii) and properties (including electronegativity, first ionisation energy, metallic and non-metallic character and reactivity) of elements.

Key Takeaways
An is a region of space in which electrons exist, not in fixed discrete energy levels
Each orbital can hold up to electrons
There are 4 different types of orbitals: which respectively hold electrons
Each shell n contains n subshells. For example, shell 2 contains subshells: 2s and



<u>Learning Objective</u>: [1.1.4] Write Electron Configurations of Elements and Ions, in both Ground and Excited States, using both Bohr and Schrodinger Models (including Cu and Cr exceptions and Condensed Notation)

### Study Design

the periodic table as an organisational tool to identify patterns and trends in, and relationships between, the structures (including shell and subshell electronic configurations and atomic radii) and properties (including electronegativity, first ionisation energy, metallic and non-metallic character and reactivity) of elements

Key Takeaways
Electron shells are filled in order from the nucleus, with the shells fully filled before moving on.
Each shell can hold up to electrons, where n is the shell number
The octet rule states that the valence electron shell can only hold a maximum of electrons.
Schrodinger representation:
$1s^22s^22p^63s^2$
When writing out the electron configuration of elements, start from the energy levels
4s subshell fills before subshell
If electrons are found to be in higher energy levels – <b>shells or subshells</b> - without the lower ones being filled first, you can assume that they are in the state.
Cu and Cr Shrodinger electron configurations are
Chromium:
<b>€</b> Copper:
This is due to an in these forms
Condensed notation always has a in square brackets, as these are the elements with full outer shells



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

# VCE Chemistry ½

# Free 1-on-1 Support

### Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

1-on-1 Video Consults	<u>Text-Based Support</u>
<ul> <li>Book via <a href="bit.ly/contour-chemistry-consult-2025">bit.ly/contour-chemistry-consult-2025</a> (or QR code below).</li> <li>One active booking at a time (must attend before booking the next).</li> </ul>	<ul> <li>Message <u>+61 440 137 304</u> with questions.</li> <li>Save the contact as "Contour Chemistry".</li> </ul>

Booking Link for Consults
bit.ly/contour-chemistry-consult-2025



Number for Text-Based Support +61 440 137 304

