# **CONTOUREDUCATION**

Website: contoureducation.com.au | Phone: 1800 888 300

Email: hello@contoureducation.com.au

# VCE Chemistry ½ Intermolecular Bonding [1.8]

Workbook

# Outline:

Intermolecular Bonding
Types of Bonding
Dispersion Forces
Dipole-Dipole Attraction
Hydrogen Bonding
Comparing Intermolecular Bonding

# **Learning Objectives:**

CH12 [1.8.1] - Explain how dispersion forces form & identify molecules that can exhibit them
 CH12 [1.8.2] - Explain how dipole-dipole attractive forces form & identify molecules that can exhibit them
 CH12 [1.8.3] - Explain how hydrogen bonds form & identify molecules that can exhibit them
 CH12 [1.8.4] - Apply intermolecular bonding to compare molecules' melting & boiling points



# Section A: Intermolecular Bonding

# **Sub-Section**: Types of Bonding



#### **Context**

Covalent bonds, ionic bonds, and metallic bonds are examples of intramolecular bonds.



# What even is a bond?



#### **Chemical Bond**

- Definition:
  - A chemical bond is a strong force of \_\_\_\_\_\_.
- There are two main categories of chemical bonds covered in VCE Chemistry. They include:
  - e \_\_\_\_\_Bonds.
  - G \_\_\_\_\_Bonds



<u>Discussion:</u> What is the difference between intramolecular and intermolecular bonding?

# **C**ONTOUREDUCATION

#### **Exploration**: Intramolecular vs Intermolecular Bonds


- Consider 3 water molecules:
- Type of bond between 0 and H:

[Covalent] / [lonic] / [Metallic]

Location of covalent bond:

[Within] / [Between] water molecule


- As such, what type of bond is it? (Label Below)
- ▶ What bonds exist **between** each of the molecules of water? (Label Below)



# Intramolecular vs Intermolecular Bonding



- Intramolecular bonding happens \_\_\_\_\_ molecules.
- Intermolecular bonding happens \_\_\_\_\_ molecules.



**TIP:** You can think of **inter**molecular as going between molecules as **inter**national flights go **between** countries.





# **Sub-Section**: Dispersion Forces






<u>Activ</u>	ve Recall: What is a dipole?	?
-		

# **Exploration**: Dispersion Forces



- Inside atoms, electrons exist in a region of space called the \_\_\_\_\_\_
  - The electrons inside this electron cloud move around \_\_\_\_\_\_\_.
- Consider Beryllium:
  - The electrons *could* be distributed \_\_\_\_\_\_. (Label Below)
  - However, as they move randomly, they could also be distributed \_\_\_\_\_\_. (Label Below)
  - When the electrons are distributed **unevenly**, what charges will exist in the atom? (Label Below)



Electron Movement:

[Fixed Positions] / [Constant Motion]

• Will they remain in these positions for a long time?


[Yes] / [No]




Dipole Formation:

# [Temporary] / [Permanent]

- Consider two molecules, both of which have a temporary, uneven distribution of electrons. (Label Below)
- What type of intermolecular bond will form between the two molecules? (Label Below)








# **Dispersion Forces**



- Definition:
  - When two or more **temporary** or \_\_\_\_\_\_ dipoles come together, arising from the **asymmetrical** distribution of electrons.
  - Arises due to electrons randomly moving in molecules and temporarily group together on one side.





<u>Discussion</u>: Are dispersion forces a strong type of intermolecular bond?



**Discussion: Which molecules have dispersion forces?** 



**NOTE:** Dispersion forces are the \_\_\_\_\_\_ type of intermolecular bonding that exists, as they are **[temporary] / [permanent]** and last for short periods of time.




**ALSO NOTE:** Dispersion forces occur in \_\_\_\_\_ molecules.

# Are dispersion forces equally strong in every molecule?



# **Exploration**: Strength of Dispersion Forces




[More] / [Less] electrons


Will a molecule with more or less electrons have stronger dispersion forces?

[More] / [Less] electrons

What will this look like? (Label Below)



G





# **Features of Dispersion Forces**



- Weakest type of intermolecular bonding due to temporary dipoles.
- Occurs in all molecules.
- Atoms/molecules with more electrons have a greater chance for temporary dipoles to be formed and have stronger dispersion forces.

# Let's have a look at a question together!



Questi	on 1 (2 marks) Walkthrough.				
Explain whether hydrogen gas (H <sub>2</sub> ) or fluorine gas (F <sub>2</sub> ) will form stronger intermolecular bonds.					

**TIP:** A quick way to figure out which molecule will have stronger dispersion forces is by checking which one has a greater atomic number/mass number!





# Your Turn!

<b>Y</b>

Question 2			
State which molecule of the pair has stronger dispersion forces.			
<b>a.</b> Methane (CH <sub>4</sub> ) vs Hydrogen Gas (H <sub>2</sub> )	<b>b.</b> Nitrogen (N <sub>2</sub> ) vs Bromine (Br <sub>2</sub> )		
Question 3 (2 marks)			
Navindu is investigating intermolecular bonding and of Explain how dispersion forces form in molecules.	doesn't quite understand how dispersion forces work.		
	doesn't quite understand how dispersion forces work.		



**REMINDER:** Some molecules naturally exist diatomically.



#### Question 5 (1 mark) Additional Question.

Select which of the following correctly explains the difference in intermolecular bonding strength between  $CO_2$  and  $N_2$ .

- A.  $CO_2$  has stronger bonding as it is polar whereas  $N_2$  is non-polar.
- **B.** CO<sub>2</sub> is a larger molecule, making the bonds heavier and thus stronger.
- **C.** There are more electrons present in CO<sub>2</sub>, which increases the chance of temporary dipoles to form, thereby increasing the strength of the dispersion forces.
- $\mathbf{D}$ .  $\mathrm{CO}_2$  is heavier and as such can form permanent dipoles.



# **Sub-Section**: Dipole-Dipole Attraction



# Let's take a look at another type of intermolecular bonding!



Discussion: What type of molecules contain permanent dipoles?



# **Exploration**: Hydrogen Chloride (HCI) Molecules



- Two adjacent HCl molecules:
- What is the more electronegative atom?

[Chlorine] / [Hydrogen]

- What are the partial charges within each of the HCl molecules? (Label Below)
- What will form in each HCl molecule? \_\_\_\_\_\_ (Label Below)
- What will occur between the adjacent ends of the two molecules? (Label Below)
- What do we call this intermolecular bond formed? (Label Below)








#### **Dipole-Dipole Attraction**



- Definition:
  - Dipole-Dipole Attractions are an intermolecular bond that occurs between two dipoles.



NOTE: Non-polar molecules can only undergo the weaker dispersion forces.



Can dipole-dipole attractions form between any polar molecules or do the molecules have to be diatomic?

#### Exploration: Chloromethane (CH<sub>3</sub>Cl)



➤ How can the C — H bond be classified?

[Chlorine] / [Hydrogen]

- Which atom is more electronegative?
- What will the partial charges and dipoles look like? (Label Below)
- What type of intermolecular bond will exist between the two molecules? (Label Below)





**NOTE:** Regardless of how the polar molecule is arranged, some parts of the polar molecule can act as a dipole, and thus dipole-dipole interactions can form.




<u>Discussion:</u> What type of intermolecular bonding is stronger? <u>Dispersion forces</u> or <u>dipole-dipole</u>?



# **Features of Dipole-Dipole Attractions**



Dipole-dipole attractive forces also exist in polar molecules that are not diatomic:



Dipoles in polar molecules are \_\_\_\_\_\_, these are \_\_\_\_\_ than **temporary** dispersion forces.

**NOTE:** The **more polar** the molecule is, the \_\_\_\_\_\_ the dipole-dipole attraction will be.





# Let's have a look at a question together!



#### Question 6 Walkthrough.

For each of the following molecules:

- i. Draw the Lewis structure.
- ii. Draw any relevant polarity arrows.
- iii. State the strongest type of intermolecular bonding they can exhibit.
- **a.**  $0_2$

**b.** CO<sub>2</sub>

c. HF

**REMINDER:** Don't forget If the polarity arrows cancel due to the molecule being symmetric, the molecule is **non-polar** overall, despite having polar bonds.





# Your Turn!



Qu	esti	on 7 (2 marks)
Fly	nn i	s investigating two of the most abundant chemicals used in society, methane (CH <sub>4</sub> ) and water (H <sub>2</sub> 0).
a.	Dra	aw the Lewis structures of:
	i.	Water
	ii.	Methane
b.	Sta	te the polarity of:
	i.	Water
	ii.	Methane
c.	He	nce or otherwise, explain whether water or methane will have stronger intermolecular bonding. (2 marks)



Question 8 (2 marks)				
Explain whether Hydrofluoric acid (HF) or Hydrochloric acid (HCl) will have stronger intermolecular bonding.				
Question 9 (1 mark)				
Which of the following best compares the intermolecular	bonding of HCl versus Br <sub>2</sub> ?			
<b>A.</b> Both molecules are non-polar but Br <sub>2</sub> has stronger dis	spersion forces.			
<b>B.</b> HCl is polar whereas Br <sub>2</sub> is non-polar so HCl will have dispersion forces.	e dipole-dipole forces which are stronger than Br <sub>2</sub> 's			
C. Br <sub>2</sub> 's dispersion forces will overpower HCl's dipole-dipole bonds, and so bromine has stronger intermolecular bonding.				
<b>D.</b> Both are polar but HCl has a larger electronegativity difference so it has stronger dipole-dipole bonding.				
<b>NOTE</b> : You are not expected to know the cases in which dispersion forces overpower dipole-dipole forces of attraction, but the idea is very important!				
Question 10 Additional Question.				
For each of the following, list <b>all</b> the types of intermolecular forces present between the molecules.				
a. Hydrochloric acid (HCl)	c. Ammonia (NH <sub>3</sub> )			
<b>b.</b> Methane (CH <sub>4</sub> )	<b>d.</b> Methanol (CH <sub>3</sub> OH)			



**REMINDER:** Don't forget that every molecule contains dispersion forces!



# **Sub-Section:** Hydrogen Bonding



# Let's have a look at the final type of intermolecular bonding!



### Exploration: Hydrofluoric Acid (HF) Bonding

- Consider adjacent HF molecules:
  - What will happen to the shared pair of electrons and their partial charges? (Label Below)
  - What force will exist between the F on the left and the H on the right? (Label Below)
  - What is the electronegativity difference between H and F?
  - What will happen to the hydrogen atom? What will the fluorine atom now be able to do? (Label Below)















Hence, is this a strong or weak dipole-dipole bond?

[Strong] / [Weak]

This type of bonding is given a special name, which is called a \_\_\_\_\_\_



<u>Discussion:</u> How does the strength of hydrogen bonding compare with dispersion forces and dipole-dipole attractions?






# Features of Hydrogen Bonding



- Definition:
  - Hydrogen bonds are a **special type of dipole-dipole bond** which are very strong. They are the \_\_\_\_\_\_\_ type of intermolecular bond due to:
  - The large \_\_\_\_\_\_
  - The \_\_\_\_\_ which is **small in nature**, allows molecules to approach close to it, as it is left exposed to the large electronegativity difference.



Discussion: What other highly electronegative elements can bond with H to form a hydrogen bond?



# **Exploration**: Hydrogen Bond Formation


- Hydrogen Bonds can form when there are hydrogens that are covalently bonded to one of the three following atoms:
  - **e**
  - **@**
  - **@** \_\_\_\_\_

# **CHONTOUREDUCATION**

- Consider ammonia (NH<sub>3</sub>) and water (H<sub>2</sub>0) molecules:
  - ls there a H atom **covalently** bonded to one of F, O, or N?

[Yes] / [No]

- What does this look like on NH<sub>3</sub> and H<sub>2</sub>O? (Label Below)
- Is that same H atom able to form a dipole-dipole attraction with a F, O, or N atom on an adjacent molecule?
  [Yes] / [No]
- What will this look like on NH<sub>3</sub> and H<sub>2</sub>O? (Label Below)




TIP: Remember Hydrogen has to be covalently bonded to one of FON (Fluorine, Oxygen or Nitrogen).



**NOTE:** It is actually the  $\_\_\_$  of electrons on the adjacent molecule's F, O or N that the hydrogen atom is attracted to a hydrogen bond!



<u>Discussion:</u> Is the following a hydrogen bond?







### **Misconception**



" ... and any one of Fluorine, Oxygen or Nitrogen"

#### TRUTH:


Hydrogen Bonds can only form between one of FON and a hydrogen which is \_\_\_\_\_\_ bonded to one of FON

What is the issue with the example in the discussion? (Label Above)


# **Hydrogen Bond Formation**



Hydrogen has to be covalently bonded to one of FON (Fluorine, Oxygen or Nitrogen):



► It can occur with any F, O or N atom on a neighbouring molecule, and hydrogen which is covalently bonded to one of FON:






# Let's have a look at a question together!




#### Question 11 Walkthrough.

State whether or not the following molecules can form hydrogen bonds.

a. Ammonia (NH<sub>3</sub>)



**b.** Methanal



**TIP:** When trying to determine whether a substance can form hydrogen bonds with itself, simply check there is a H atom covalently bonded to one of F, O or N!





### Recall!



<u>Active Recall:</u> Rank the 3 intermolecular bonds from strongest to weakest, and state the condition(s) for each of them to form.




Your Turn!

#### **Question 12**


State the strongest type of intermolecular bonding in the following molecules.

**a.** Methanol (CH<sub>3</sub>OH)






**b.** CHCl<sub>3</sub>




 $\mathbf{c}$ .  $SO_2$ 



### **Question 13 Additional Question.**

List **all** the intermolecular bonds that can exist between the following molecule:



CH12 [1.8] - Intermolecular Bonding - Workbook




# **Misconception**

"Hydrogen bonds have to occur in molecules which only have a  $\,H-F,H-O\,$  or  $\,H-N\,$  bond

#### Truth:

It can occur with any F, O, or N atom on a neighbouring molecule and a hydrogen which is covalently bonded to one of FON.

- Consider Methanal and water:
  - Where would a hydrogen bond exist? (Label Below)
  - What is different about the 0 in methanal? (Label Below)



Is this still a hydrogen bond?

[Yes] / [No]

**NOTE:** Understanding intermolecular bonding, especially hydrogen bonding, is something that is essential for ¾ Chemistry as well!





# Let's have a look at another example together!

7
Ĭ

	Question	14	Walkthrough.
--	----------	----	--------------

State the strongest type of intermolecular bonding that will exist between the following molecules:

- **a.** Methanol ( $CH_3OH$ ) and carbon dioxide ( $CO_2$ )
- **b.** Methane  $(CH_4)$  and nitrogen  $(N_2)$

# Try a few more questions!




Question 15 (3 marks)				
Explain why ammonia can form hydrogen bonds. Use a diagram to support your answer.				




### **Question 16**

State the strongest type of intermolecular bonding in the following molecules.


a. Ethanoic Acid




**b.** Methanamine (CH<sub>3</sub>NH<sub>2</sub>)



**c.** Methyl Methanoate (CH<sub>3</sub>COCH<sub>3</sub>)









NOTE: We'll cover the naming of these organic compounds later on in the year!



#### **Question 17 Additional Question.**

Draw two water molecules, explaining and labelling the strongest intermolecular bonding between them.





# **Sub-Section:** Comparing Intermolecular Bonding



# Let's now compare these bonds!



# **Comparing Intermolecular Bonding**



	<del></del>	<del></del>
Weakest Intermolecular Bond.	Stronger Intermolecular Bond.	Strongest Intermolecular Bond.
Occurs in <b>all</b> molecules.	Only occurs in polar molecules.	Only occurs between one of FON and a hydrogen which is covalently bonded to FON.
Formed by temporary/instantaneous dipoles which form temporary partial charges.	Formed from permanent dipoles due to polarity of molecules which results in electrostatic attraction.	Formed from the high electronegativity between hydrogen (H) and one of FON which exposes the hydrogen atom to even stronger electrostatic attraction.

Share	101	Personal	Morez





# Let's have a look at a question together!



Question 18 (5 marks) Walkthrough.			
Matthew is investigating hydrochloric acid (HCl) and water (H <sub>2</sub> 0).			
a. Draw the Lewis structures for each of these molecules. (2 marks)			
<b>b.</b> Hence or otherwise, explain whether hydrochloric acid (HCl) or water (H <sub>2</sub> O) will have stronger intermolecular bonding. (3 marks)			
Sans for Daniel Makes			
Space for Personal Notes			



### Your Turn!




Question 19 (6 marks)

State which of the following has stronger intermolecular bonding, and briefly justify your answer.

**a.** CH<sub>2</sub>Cl<sub>2</sub> vs HF. (2 marks)

**b.** Ethyl ethanoate vs Ethane. (2 marks)





c. Methanal vs Hydrochloric acid. (2 marks)





#### **Question 20**

Which of the following substances exhibits the strongest hydrogen bonding?

- **A.** CH<sub>4</sub> (Methane)
- **B.**  $H_2O$  (Water)
- C. HF (Hydrogen fluoride)
- **D.** NH<sub>3</sub> (Ammonia)

#### **Question 21**

Select the statement that describes the two requirements for hydrogen bonding.

- **A.** A hydrogen atom covalently bonded to any atom and a non-bonding pair of electrons on any atom.
- **B.** A hydrogen atom covalently bonded to an oxygen, a nitrogen or a fluorine atom and a non-bonding pair of electrons on any other atom.
- **C.** A hydrogen atom covalently bonded to any atom and a non-bonding pair of electrons on a neighbouring oxygen, nitrogen or fluorine atom.
- **D.** A hydrogen atom covalently bonded to an oxygen, nitrogen or fluorine atom and a non-bonding pair of electrons on a neighbouring oxygen, nitrogen or fluorine atom.

#### **Question 22 Additional Question.**

Identify which of the following molecules would form hydrogen bonds with water.

a. Ethanol, CH<sub>3</sub>CH<sub>2</sub>OH

**d.** Ethanoic acid, CH<sub>3</sub>COOH

**b.** Ammonia, NH<sub>3</sub>

e. Tetrachloromethane, CCl<sub>4</sub>

c. Phosphine, PH<sub>3</sub>

**f.** Ethanal, CH<sub>3</sub>CHO



### **Key Takeaways**



- ✓ Intramolecular bonding happens within molecules.
- ✓ Intermolecular bonding happens **between** molecules.
- ☑ **Dispersion forces** occur when two or more **temporary** or **instantaneous** dipoles come together, arising from the **asymmetrical** distribution of electrons
- Dispersion forces occur as electrons are **randomly** moving in molecules, and so they may **temporarily group closely together** on one side
- ☑ Dispersion forces are the **weakest** type of intermolecular bonding that exist, and occur in **all** molecules.
- Atoms/molecules with more electrons have a greater chance for temporary dipoles to be formed, and as such, have stronger dispersion forces
- ☑ Dipole-Dipole Attractions are an intermolecular bond that occurs between two dipoles.
- While dispersion forces can occur in **all** molecules, dipole-dipole attractions only occur in **polar** molecules.
- ☑ Dipoles in polar molecules are **permanent**, these forces are **stronger** than temporary dispersion.
- ✓ Hydrogen bonds are a **special type of dipole-dipole bond.** They are the **strongest** type of intermolecular bond due to:
  - The large electronegativity difference.
  - The **hydrogen atom** which is **small in nature** allows molecules to approach close to it as it is left exposed from the large electronegativity difference.
- For a H-bond to form:
  - A hydrogen atom has to be **covalently bonded** to one of **FON** (Fluorine, Oxygen or Nitrogen):



That same H atom can form a dipole-dipole bond with any F, O or N atom on a neighbouring molecule

<u>Dispersion Forces</u>	<u>Dipole-dipole</u>	<u>Hydrogen Bonds</u>
Weakest Intermolecular Bond.	Stronger Intermolecular Bond.	Strongest Intermolecular Bond.
Occurs in all molecules.	Only occurs in polar molecules.	Only occurs between one of FON and hydrogen which is covalently bonded to FON.
Formed by temporary/instantaneous dipoles which form temporary partial charges.	Formed from permanent dipoles due to polarity of molecules which results in electrostatic attraction.	Formed from the high electronegativity between hydrogen (H) and one of FON which exposes the hydrogen atom to even stronger electrostatic attraction.

Space for Personal Notes		



# Section B: Boiling and Melting Points

#### **Context**




Substances can generally exist in one of three physical states: (Label Below)

# Particle Arrangement in Phases of Matter




Particle are packed tightly together in a fixed arrangement.

Particles can vibrate but not move.



Particles are closed together with no distinct arrangement.

Particles can move and slide around each



Particles are free floating with no distinct arrangement.

Particles move and collide with each other.

# **Boiling & Melting Points**



- Boiling is the state change between \_\_\_\_\_\_ and \_\_\_\_\_ and \_\_\_\_\_.
- Melting is the state change between \_\_\_\_\_\_ and \_\_\_\_\_.


<u>Discussion:</u> What makes something a solid, liquid or gas at a given temperature?





### **Exploration:** Changing State of Matter





- When substances melt (go from solid to liquid), what happens to the intermolecular bonds?
- When substances boil (go from liquid to gas), what happens to the intermolecular bonds?

<u>Discussion:</u> If a substance has stronger intermolecular bonds, is it <u>more</u> or <u>less</u> likely to melt?



# **Changing States of Matter**



- Substances exist either as a solid, liquid or gas at a given temperature.
- State depends on the \_\_\_\_\_\_ of intermolecular bonds.

Melting (Solid to Liquid)	Intermolecular bonds are
Boiling (Liquid to Gas)	Intermolecular bonds are

<b>⇔</b> ONTOU	JREDU	ICATION
1011101		

VCE Chemistry ½ Questions? Message +61 440 137 304

NOTE: Intramolecular bonds (covalent bonds) are	_ disrupted when a phase change takes
place!	•

# A

# Let's have a look at a question together!

Question 23 (4 marks) Walkthrough.				
Explain whether water or hydrogen gas (H <sub>2</sub> ) is likely to have a higher melting point.				

#### Sample Response: Melting/Boiling Point



- Substance X has stronger intermolecular bonding because \*insert reasoning\*.
  - lt undergoes \*insert type of bond\*, which is a stronger intermolecular bond than \*insert Substance Y's bonding\*.
  - lt is a larger molecule, and therefore, can exhibit stronger dispersion forces than Substance Y.
- As such, more **thermal energy** is required to vibrate and **weaken** (**for melting point**) / **break** (**for boiling point**) the intermolecular bonds for Substance *X*.
- This leads to a higher melting/boiling point for Substance X.





#### Your Turn!



Question 24 (4 marks)				
Explain whether hydrogen chloride (HCl) or oxygen gas (O <sub>2</sub> ) has a higher boiling point.				

# REMINDER: Don't forget!



- Melting point only requires intermolecular bonds to be weakened.
- Boiling point only requires intermolecular bonds to be completely broken.

#### **Question 25**

Which statement best explains the difference in bond strengths **within** a water molecule compared to **between** water molecules?

- **A.** The covalent bonds within water molecules are stronger than the hydrogen bonds between water molecules
- **B.** Hydrogen bonds within water molecules are stronger than covalent bonds between water molecules, leading to water's low boiling point.
- **C.** The covalent and hydrogen bonds in water have similar strengths, which accounts for water's high melting point.
- **D.** Hydrogen bonds between water molecules are non-existent, resulting in water's unique physical properties.



#### **Question 26 Additional Question.**

Which statement correctly explains the difference in boiling points between ammonia (NH<sub>3</sub>), phosphine (PH<sub>3</sub>), and arsine (AsH<sub>3</sub>)?

- **A.** Arsine has the highest boiling point due to strong hydrogen bonding.
- **B.** Ammonia has a higher boiling point than phosphine and arsine due to hydrogen bonding.
- C. Phosphine and arsine have higher boiling points than ammonia because they are heavier.
- **D.** All have similar boiling points due to similar dispersion forces.



# Try one more question to bring everything together!

#### Question 27 (6 marks)

Luque is a funny lad, and thus tries to investigate two molecules – sulphur dioxide  $(SO_2)$  and carbon dioxide  $(CO_2)$ .

The structure of sulphur dioxide is shown below, which does not obey the Octet Rule. (Outside the study design.)



- **a.** Draw the polarity arrows **and** partial charges on the sulphur dioxide molecule provided. (1 mark)
- **b.** Draw the Lewis Structure of carbon dioxide  $(CO_2)$ . (1 mark)



VCE Chemistry ½ Questions? Message +61 440 137 304

c. With reference to their structure and bonding, state and explain which molecule is most likely to have a higher boiling point. (4 marks)		
Sn	pace for Personal Notes	
<b>اد</b> ا	race for reisonal notes	





# **Contour Checklist**

Learning Objective: [1.8.1] - Explain how dispersion forces form & identify molecules that can exhibit them

#### Study Design

"the relative strengths of intramolecular bonding (covalent bonding) and intermolecular forces (dispersion forces, dipole-dipole attraction and hydrogen bonding)"

#### **Key Takeaways**

- Dispersion forces: when two or more \_\_\_\_\_\_ or instantaneous dipoles come together, arising from the [symmetrical] / [asymmetrical] distribution of electrons.
- This intermolecular bond occurs because electrons are **[fixed] / [randomly moving]** in molecules, and so they may temporarily group closely together on one side.
- ☐ Dispersion forces are the **[strongest] / [weakest]** type of intermolecular bonding that exists.
- □ Dispersion forces occur in **[polar] / [non-polar] / [all] / [specific]** molecules.
- Atoms/molecules with more electrons have a [greater] / [same] / [lesser] chance for temporary dipoles to be formed, and as such, have [stronger] / [identical] / [weaker] dispersion forces.



\_\_\_\_\_ molecules.

□ <u>Lea</u>	arning Objective: [1.8.2] -	Explain how dipol	e-dipole attractive	forces
	form & identify	molecules that car	exhibit them	

# **Study Design**

"the relative strengths of intramolecular bonding (covalent bonding) and intermolecular forces (dispersion forces, dipole-dipole attraction and hydrogen bonding)"

### **Key Takeaways**

	are an intermolecular bond that occurs between <b>two dipoles</b> .
П	While dispersion forces can occur in <b>all</b> molecules, dipole-dipole attractions only occur in

As dipoles in polar molecules are [temporary] / [permanent], these	forces are [stronger] /
[weaker] than dispersion forces, wherein the dipoles are only tem	porarily formed.



Learning Objective: [1.8.3] - Explain how hydrogen bonds form & identify molecules that can exhibit them							
Study Design  "the relative strengths of intramolecular bonding (covalent bonding) and intermolecular forces (dispersion forces, dipole-dipole attraction and hydrogen bonding)"  Key Takeaways  Hydrogen bonds are a special type of dipole-dipole bond. They are thetype of intermolecular bond out of the three major intermolecular bonds. This is due to:							
				□ The large			
				☐ The atom which is [small] / [large] in nature allows molecules to approach close to it as it is left exposed from the large electronegativity difference.			
				□ For a H-bond to form:			
A hydrogen atom has to bebonded to one of FON (Fluorine, Oxygen or Nitrogen):							
That same H atom can form a <b>dipole-dipole</b> bond with <b>any F, O or N atom</b> on a							



# □ <u>Learning Objective</u>: [1.8.4] - Apply intermolecular bonding to compare molecules' melting & boiling points

### Study Design

"physical properties of molecular substances (including melting points and boiling points and nonconduction of electricity) with reference to their structure and bonding"

	Key Takeaways					
	☐ Intramolecular bonding - happens [within] / [between] molecules.					
	☐ Intermolecular bonding - happens [within] / [between] molecules.					
Substances exist either as a			a given temperature.			
☐ Their state depends on the <b>strength</b> of its			bonds.			
	is the phase change from Solid → Liquid: Intermolecular bonds are					
$\hfill \Box$ is the phase change from Liquid $\rightarrow$ Gas: Intermolecular bonds are						
	<b>Weakest</b> Intermolecular Bond. Str	ronger Intermolecular Bond.	Intermolecular Bond.			
	Occurs in <b>all</b> molecules. Onl	ly occurs in polar molecules.	Only occurs between one of FON and a hydrogen which is covalently bonded to FON.			
		ormed from dipoles due to polarity of nolecules which results in electrostatic attraction.	Formed from the high electronegativity between hydrogen (H) and one of which exposes the hydrogen atom to even stronger electrostatic			

attraction.



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

# VCE Chemistry ½

# Free 1-on-1 Support

# Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

1-on-1 Video Consults	Text-Based Support
<ul> <li>Book via bit.ly/contour-chemistry-consult- 2025 (or QR code below).</li> <li>One active booking at a time (must attend before booking the next).</li> </ul>	<ul> <li>Message +61 440 137 304 with questions.</li> <li>Save the contact as "Contour Chemistry".</li> </ul>

Booking Link for Consults
bit.ly/contour-chemistry-consult-2025



Number for Text-Based Support +61 440 137 304

