



Website: [contoureducation.com.au](http://contoureducation.com.au) | Phone: 1800 888 300  
Email: [hello@contoureducation.com.au](mailto:hello@contoureducation.com.au)

VCE Chemistry  $\frac{3}{4}$   
Spontaneous Redox Reactions [0.6]  
Workshop Solutions

Error Logbook:



Mistake/Misconception #1		Mistake/Misconception #2	
Question #:	Page #:	Question #:	Page #:
Notes:		Notes:	
Mistake/Misconception #3		Mistake/Misconception #4	
Question #:	Page #:	Question #:	Page #:
Notes:		Notes:	

## Section A: Recap (5 Marks)




### Learning Objective: [1.7.1] - Apply the ECS to Predict Spontaneous Reactions

- Net Ionic Equation Definition: A balanced full equation with spectator ions omitted.
- Spectator Ion: Compound which is present but does not participate in the reaction.

Reduction Reaction	Oxidation Reaction
[Forward] / [Reverse] reaction on ECS.	[Forward] / [Reverse] reaction on ECS.

Oxidants	Reductants
Positioned on the [left] / [right] side.	Positioned on the [left] / [right] side.

Strongest Oxidants	Strongest Reductants
Positioned [top] / [bottom] - [left] / [right].	Positioned [top] / [bottom] - [left] / [right].







➤ Steps to predicting spontaneous reaction:

1. Split all species into cations and anions. Some cations/anions are inert.
2. Locate all species on the electrochemical series. Draw a vertical line to split oxidants and reductants apart.
3. Draw a mini-electrochemical series version
4. Find the strongest oxidant and strongest reductant.
5. Check for downhill gradient.
6. Write out half-equations.

➤ When multiple oxidants/reductants are present, the strongest oxidant reacts with strongest reductant.


➤ The four ions which appear on both sides of the electrochemical series:

-  Fe<sup>3+</sup>(aq)/Fe<sup>2+</sup>(aq)
-  Sn<sup>4+</sup>(aq)/Sn<sup>2+</sup>(aq)
-  Pb<sup>2+</sup>(aq)/Pb(s)
-  Mn<sup>2+</sup>(aq)/Mn(s)

**Question 1 (3 marks) Walkthrough.**

Some copper metal is dipped into a solution which contains copper nitrate, tin (II) nitrate, zinc chloride and aluminium bromide.

Find the overall reaction which takes place.





### Learning Objective: [1.7.2] - Identify Differences Between Direct & Indirect Redox Reactions & Features of ECS

- Standard Electrode Potential Definition: Method to measure electromotive force (EMF).
- Standard Hydrogen Electrode (SHE):  $\text{H}^+(\text{aq})/\text{H}_2(\text{g})$  which has  $E^0 =$  0.00 V.
- The electrochemical series does not predict the rate of reaction.

<u>Direct Contact Spontaneous Redox Reaction</u>	<u>Indirect Contact Spontaneous Redox Reaction</u>
<u>Chemical → Thermal</u>	<u>Chemical → Electrical</u>



### Learning Objective: [1.7.3] - Find Strongest Oxidants/Reductants by Constructing Your Own ECS

- Electrochemical series ordered from [lowest → highest] / highest → lowest  $E^0$  value.

<u>Strongest Oxidant</u>	<u>Strongest Reductant</u>
<u>[Highest]</u> / [Lowest] $E^0$ value.	[Highest] / <u>[Lowest]</u> $E^0$ value.

- Creating electrochemical series yourself steps:
  1. Draw a vertical line to separate oxidants and reductants.
  2. Using information, place oxidants/reductants on this mini electrochemical series.

<u>Spontaneous Reactions</u>	<u>Non-Spontaneous Reactions</u>
[Positive] / <u>[Negative]</u> gradient.	<u>[Positive]</u> / [Negative] gradient.

3. Write the conjugate version of the oxidant/reductant.
4. Repeat for each piece of information.

Space for Personal Notes

**Question 2 (2 marks) Walkthrough.**

There are three unknown substances, P, Q and R. The following half-equations are given, but their  $E^0$  values are not given.

Reaction
$P^{2+}(aq) + 2e^- \rightleftharpoons P(s)$
$Q^{2+}(aq) + 2e^- \rightleftharpoons Q(s)$
$R^{2+}(aq) + 2e^- \rightleftharpoons R(s)$
$S^{2+}(aq) + 2e^- \rightleftharpoons S(s)$

It is known that when P is mixed into a solution of  $R^{2+}$ , no observable reaction occurs.

It is also known that when Q is mixed into a solution containing  $R^{2+}$ , no observable reaction occurs.

When S and  $Q^{2+}$  are combined, a reaction occurs.

When  $S^{2+}$  and P are combined, a reaction occurs.

Rank the three metals in terms of their decreasing oxidant strength.

► Decreasing oxidant strength: \_\_\_\_\_  $P^{2+}, S^{2+}, Q^{2+}, R^{2+}$  \_\_\_\_\_

Space for Personal Notes

## Section B: Warm Up (13.5 Marks)

INSTRUCTION: 13.5 Marks. 9 Minutes Writing.



### Question 3 (0.5 marks)

What is the strongest reductant out of the following chemicals?




### Question 4 (2 marks)


Cobalt (II) nitrate has cadmium metal dipped inside of it.

a.


- i. Write the reduction reaction which takes place. (0.5 marks)



- ii. Write the oxidation reaction which takes place. (0.5 marks)



- b. Write the **full balanced ionic equation**. (1 mark)





Space for Personal Notes

**Question 5** (3 marks)

For each of the following, determine whether a reaction will occur or not. If there is a reaction, write the relevant reduction and oxidation reactions.



- a. A solid nickel rod (Ni) dipped into a solution containing tin (II) nitrate. (1 mark)

Reaction is: **Spontaneous** / [Non-Spontaneous]



- b. A solution containing hydrofluoric acid (HF) is mixed with a strip of calcium metal. (1 mark)

Reaction is: **Spontaneous** / [Non-Spontaneous]



- c. A solution containing iron (II) nitrate and zinc metal. (1 mark)

Reaction is: [Spontaneous] / **Non-Spontaneous**

No reaction.

Space for Personal Notes

**Question 6** (3 marks)

A solution of iron (III) fluoride has a nickel rod placed inside of it.

- a. Write the half-reactions which take place: (2 marks)

Reduction:  $\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}$

Oxidation:  $\text{Ni} \rightarrow \text{Ni}^{2+} + 2\text{e}^-$

- b. Write the overall reaction which takes place. (1 mark)

$\text{Ni} + 2\text{Fe}^{3+} \rightarrow 2\text{Fe}^{2+} + \text{Ni}^{2+}$

**Question 7** (1 mark)

Solution I - 1.0 M NaCl

Solution II - 1.0 M CuCl<sub>2</sub>

Solution III - 1.0 M MgCl<sub>2</sub>

Which solution or solutions above will react with Zn powder?





- A. Solution I only.
- B. Solution II only.**
- C. Solutions I and III only.
- D. Solutions I, II and III.

Space for Personal Notes



**Question 8** (4 marks)

The following half-equations are given:



- a. State the weakest oxidant and the weakest reductant.

Weakest Oxidant			
Weakest Oxidant		Weakest Reductant	
$\text{B}^{2+}(\text{aq})$		$\text{A}(\text{s})$	

- a. A solution of  $\text{B}^{2+}(\text{aq})$  is mixed with some  $\text{D}(\text{s})$ . Will a reaction occur? Explain why/why not, and if there is a reaction, write the overall reaction which takes place. (2 marks)

No –  $\text{B}^{2+}$  is a weaker oxidant than  $\text{D}$

- b. A solution of  $\text{A}^{2+}(\text{aq})$  is mixed with some  $\text{C}(\text{s})$ . Will a reaction occur? Explain why/why not, and if there is a reaction, write the overall reaction which takes place. (2 marks)

Yes –  $\text{A}^{2+} + \text{C} \rightarrow \text{C}^{2+} + \text{A}$

Space for Personal Notes

## Section C: Ramping Up (11 Marks)

INSTRUCTION: 11 Marks. 8 Minutes Writing.



### Question 9 (1 mark)

Some strips of the metals, iron, zinc and silver were placed in separate beakers, each containing 1.0 M nickel (II) sulfate solution in water at 25°C.

What is expected to occur over time?

- A. Ni will be deposited in all of the beakers.
- B. Ni will not be deposited in any of the beakers.
- C. A reaction will occur only in the beakers containing Ag.
- D. A reaction will occur only in the beakers containing Fe and Zn.**

### Question 10 (2 marks)

A solution containing silver bromide is mixed with a solution of tin (II) chloride.

State whether a reaction will occur or not. If yes, write the overall reaction which takes place. If not, explain why no reaction will occur.

\_\_\_\_\_

\_\_\_\_\_

Yes –  $2\text{Ag}^+(\text{aq}) + \text{Sn}^{2+}(\text{aq}) \rightarrow \text{Sn}^{4+}(\text{aq}) + 2\text{Ag}(\text{s})$

\_\_\_\_\_

\_\_\_\_\_


Space for Personal Notes

**Question 11** (6 marks)


A solution of iron (II) nitrate was placed in a beaker with  $\text{Ag}_2\text{SO}_4$ .

a.


- i. Write the reduction half-equation. (1 mark)



- ii. Write the oxidation half-equation. (1 mark)



- iii. Hence, write the overall equation. (1 mark)



- b. It is then noted that the beaker has a very high pH. Would this realisation have any effect on the reaction(s) taking place? Explain your answer, writing any relevant half-equations to justify your answer. (2 marks)

Yes, as  $\text{OH}^-$  is present, which is a stronger reductant than  $\text{Fe}^{2+}$ .

Therefore, oxidation half-equation would now be  $4\text{OH}^-(\text{aq}) \rightarrow \text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^-$

- c. If, instead of  $\text{Ag}_2\text{SO}_4$ , the beaker had  $\text{CdSO}_4$ , outline what effect this would have on the reaction(s) taking place. (1 mark)

No reaction would be able to occur as oxidant is no longer higher than reductant.

Space for Personal Notes

**Question 12** (2 marks)

Nethaya is similarly given five metals and 1 M solutions of nitrates of the metals.

The metals are labelled A, B, C, D and E, and the solutions are labelled  $A^{2+}$ ,  $B^{2+}$ ,  $C^{2+}$ ,  $D^{2+}$  and  $E^{2+}$ .

The student carries out several experiments and the results obtained are listed below.

- Metal A reacts with  $B^{2+}$  spontaneously.
- Metal C becomes coated with another metal when placed in each of solutions  $A^{2+}$ ,  $B^{2+}$ ,  $D^{2+}$ , but not with  $E^{2+}$ .
- When metal A is dipped into a solution of  $D^{2+}$ , no reaction takes place.

Rank each of the 5 metals in order of increasing  $E^0$  values.

	$B^{2+}$	B	
	$A^{2+}$	A	
	$D^{2+}$	D	
	$C^{2+}$	C	
	$E^{2+}$	E	

- Increasing  $E^0$  values: E, C, D, A, B

Space for Personal Notes

## Section D: Getting Trickier I (12 Marks)

INSTRUCTION: 12 Marks. 10 Minutes Writing.



### Question 13 (1 mark)

Samples of four metals (W, X, Y and Z) were each placed in separate solutions containing the cations  $W^{2+}$ ,  $X^{2+}$  and  $Y^{2+}$ . If a reaction occurred, a tick was placed in the appropriate cell of the results table shown below.

		Metal			
		W	X	Y	Z
Solution	$W^{2+}$			✓	✓
	$X^{2+}$	✓		✓	✓
	$Y^{2+}$				✓

Which of the following shows the metals in order of **decreasing** reductant strength?

A. Z, Y, W, X

B. Y, W, X, Z

C. X, W, Y, Z

D. Z, X, W, Y

#### Question 7 A



W reacts with  $X^{2+}$ . W is therefore a stronger reductant than X. Y reacts with  $W^{2+}$  and  $X^{2+}$ . Y is therefore a stronger reductant than W and X. Z reacts with  $W^{2+}$ ,  $X^{2+}$ , and  $Y^{2+}$ . Z is therefore a stronger reductant than W, X and Y. The order of reductant strength is therefore  $Z > Y > W > X$ .

Space for Personal Notes


**Question 14** (4 marks)

Johaann, who is a bored Contour student, is chilling at home, when he decides to get a solution of iron (III) chloride and bubbles hydrogen sulphide ( $\text{H}_2\text{S}$ ) into the solution.

Hydrogen sulphide can react according to the following reaction:




- a. Write the half-equation for the oxidation reaction which takes place. (1 mark)




b.

- i. Write the balanced equation for the overall reaction which takes place. (1 mark)



- ii. Write the balanced equation for the overall reaction which takes place in alkaline conditions. (1 mark)



- c. As the reaction takes place, explain how the pH will change as the reaction proceeds. (1 mark)


pH will

**Question 15** (7 marks)


Lithium-ion batteries are becoming very commonplace and useful to society. Lithium can oxidise into  $\text{Li}^+$ , but a major issue is that Li is an extremely reactive metal.

Siggy, who does not study VCE chemistry, suggests to her friend, Nishi, to simply react Li with something random to oxidise it into the desired  $\text{Li}^+$ . Nishi naively decides to place Li in a solution of aluminium hydroxide, and a violent explosion is observed.

- a. Write the two half-equations for this reaction. (2 marks)



- b. Hence, write the overall equation. (1 mark)



- c. With reference to specific chemicals, explain why an explosion is observed. (2 marks)

$\text{H}_2(\text{g})$  is produced, which is flammable. It reacts with  $\text{O}_2(\text{g})$  in the air as follows:  $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{g})$

- d. If a strong oxidant such as  $\text{Au}^+(\text{aq})$  were to be reacted with lithium metal, would the desired lithium ions be produced **safely**? Justify your answer. (2 marks)

Still no. Although  $\text{Au}^+$  is a stronger oxidant than  $\text{H}_2\text{O}(\text{l})$ , the moment the Li touches the surface of the water, it will explode;  $\text{Au}^+(\text{aq})$  won't get a chance to prevent water from spontaneously reacting with Li as the  $\text{Au}^+$  is inside the water itself.

Space for Personal Notes

## Section E: Getting Trickier II (13 Marks)

INSTRUCTION: 13 Marks. 12 Minutes Writing.



### Question 16 (1 mark)

A student reacted 4 metals (A, B, C and D) with 1 M solutions of their corresponding ion. A table of results was set up with a tick placed against any reaction that occurred.

Metal/Solution	$A^{2+}$	$B^{+}$	$C^{2+}$	$D^{3+}$
A			✓	✓
B	✓		✓	✓
C				✓
D				

The strongest reductant and weakest oxidant respectively are:

A. B and  $B^{+}$ .

B. D and  $D^{3+}$ .

C. B and  $D^{3+}$ .

D. D and  $B^{+}$ .

The strongest reductant is B as it is oxidised by all of the solutions. The weakest oxidant is  $B^{+}$  as it is reduced by none of the reductants.

Space for Personal Notes




**Question 17** (5 marks)


Renee wants to test her chemistry and thus decided to bubble fluorine gas into a solution of silver (I) chloride.

- a. State the strongest oxidant and reductant present. (1 mark)

Strongest reductant: \_\_\_\_\_




Strongest oxidant: \_\_\_\_\_




- b. Write the half-equations which take place. (2 marks)

Reduction half-equation: \_\_\_\_\_



Oxidation half-equation: \_\_\_\_\_



c.

- i. Bubbles are observed as the reaction takes place. Explain this observation with reference to the products formed. (1 mark)

$\text{O}_2(\text{g})$  is produced as water is oxidised (a lot of students will erroneously say  $\text{Cl}_2(\text{g})$ )

- ii. List another possible observation. (1 mark)

Decrease in pH as  $\text{H}^+(\text{aq})$  is produced.

Space for Personal Notes

**Question 18** (7 marks)

Stainless steel items are very popular in the hospitality industry. Steel is made up mostly of iron, carbon, and small amounts of cobalt and manganese.

a.

- i. If a steel pan were to be filled with water, would a reaction be expected to occur? Justify your answer. (2 marks)

Yes, as the oxidant ( $\text{H}_2\text{O}$ ) is higher on the ECS than the strongest reductant (Mn).

- ii. Why is a reaction often not observed in practice? (1 mark)

Rate of reaction is too slow.

b.

- i. Now a mixture of  $\text{ZnSO}_4(\text{aq})$  and  $\text{Ni}(\text{NO}_3)_2(\text{aq})$  solutions are tossed into the steel pan. Write the overall equation taking place. (1 mark)



- ii. If the  $\text{Ni}(\text{NO}_3)_2(\text{aq})$  runs out after some time, causing another reaction to take place. Write the half-equation for the new reaction which takes place. Explain your answer. (2 marks)

$\text{Zn}^{2+}$  is now strongest oxidant so it would react with Mn(s):  
 $\text{Zn}^{2+}(\text{aq}) + \text{Mn}(\text{s}) \rightarrow \text{Zn}(\text{s}) + \text{Mn}^{2+}(\text{aq})$

- iii. After a prolonged time, both the  $\text{Ni}(\text{NO}_3)_2(\text{aq})$  and the Mn(s) in the steel pan ran out. Explain what would be observed to occur now. (1 mark)

Reactions would cease as strongest oxidant ( $\text{Zn}^{2+}(\text{aq})$ ) is lower on ECS than strongest reductant (Fe(s)).

*Let's take a BREAK!*



Space for Personal Notes

## Section F: VCAA-Level Questions I (13 Marks)

INSTRUCTION: 13 Marks. 30 Seconds Reading. 12 Minutes Writing.



### Question 19 (13 marks)




Inspired from VCAA Chemistry Exam 2019

<https://www.vcaa.vic.edu.au/Documents/exams/chemistry/2019/NHT/2019chem-nht-w.pdf#page=24>

Energy can be produced in a variety of ways, including from galvanic cells, fuel cells and gas-fired power stations. Each of these methods suits particular applications.


Galvanic cells are methods of energy production that are based on redox reactions, similar to the reaction that would occur in Set-up A shown below. Set-up A consists of a beaker with a strip of Zinc, Zn, in a solution of lead (II) nitrate,  $\text{Pb}(\text{NO}_3)_2$ .

#### Set-up A




a. For the reaction which occurs in set-up A,

i. Identify the oxidising agent. (1 mark)



ii. Write the oxidation half-equation for this reaction. (1 mark)



iii. Provide the location of the two products for this system. (1 marks)

$\text{Zn}^{2+}(\text{aq})$  ions will be found along the lead nitrate solution.  
Pieces of  $\text{Pb}(\text{s})$  will be found on the strip of zinc.

b. Due to the transfer of electrons in this reaction, identify:

i. The sign of change in chemical enthalpy of the system. (1 mark)

Negative.


ii. The energy transformations. (1 mark)

Chemical energy is converted to heat energy.

c. State the specific properties of reactants present in this system that allow a reaction to occur and explain why this is the case. State any assumptions made in using the electrochemical series to predict the reactions occurring in this example. (3 marks)

$\text{Pb}^{2+}(\text{aq})$  has a stronger reduction potential than the  $\text{Zn}^{2+}/\text{Zn}$  conjugate redox pair, thus allowing a spontaneous redox reaction to occur. The reaction in the system must be spontaneous as no external energy is provided to force a reaction. SLC is assumed as this is how the ECS was constructed.

A similar setup to **part a.** is shown below.



All substances are placed together simultaneously in an already acidic environment which leads to the heating of the beaker.

The  $\text{Cr}_2\text{O}_7^{2-}(\text{aq})/\text{Cr}^{3+}(\text{aq})$  conjugate redox pair is known to have an  $E^\circ$  value of  $+1.36 \text{ V}$ .

The  $\text{SO}_4^{2-}(\text{aq})/\text{SO}_3^{2-}(\text{aq})$  is known to have a  $E^\circ$  value of  $-0.94 \text{ V}$ .

**d.** As they mix together, a reaction begins to occur.

**i.** Write the half-equations which occur for (2 marks)

Reduction: — Reduction:  $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14\text{H}^+(\text{aq}) + 6\text{e}^- \rightarrow 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$  —

Oxidation: — Oxidation:  $\text{SO}_3^{2-}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{SO}_4^{2-}(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^-$  —

**ii.** Write the overall reaction which occurs. (1 mark)

Overall:  $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 8\text{H}^+(\text{aq}) + 3\text{SO}_3^{2-}(\text{aq}) \rightarrow 2\text{Cr}^{3+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l}) + 3\text{SO}_4^{2-}(\text{aq})$

**e.** The primary reductant is measured and observed to have run out, however, a reaction keeps occurring, leading to a decrease in pH. Write the overall equation that occurs. (2 marks)

Oxidation:  $2\text{H}_2\text{O}(\text{l}) \rightarrow \text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^-$

Overall:  $2\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 16\text{H}^+(\text{aq}) \rightarrow 8\text{H}_2\text{O}(\text{l}) + 4\text{Cr}^{3+}(\text{aq}) + 3\text{O}_2(\text{g})$

## Section G: Multiple Choice Questions (7 Marks)

INSTRUCTION: 7 Marks. 7 Minutes Writing.



*The following information relates to the next two questions.*

A number of experiments were conducted using various metals (Zn, Cu, Cr and Cd) and solutions of their ions. The results are shown in the table below.

<u>Experiment</u>	<u>Result</u>
Cadmium and copper (I) nitrate solution.	Copper metal deposited.
Cadmium and zinc nitrate solution.	No reaction.
Chromium and cadmium nitrate solution.	Cadmium metal deposited.

### Question 20 (1 mark)

From the results in the table, the relative reducing strength of three of the metals can be deduced. Beginning with the weakest reductant, the order of increasing reductant is:

A.  $\text{Cd} < \text{Zn} < \text{Cr}$

B.  $\text{Cu} < \text{Cr} < \text{Cd}$

C.  $\text{Zn} < \text{Cd} < \text{Cu}$

**D.  $\text{Cu} < \text{Cd} < \text{Cr}$**

These deductions can be made from the experiments: I – Cd is a stronger reductant than Cu; II – Zn is a stronger reductant than Cd; III – Cr is a stronger reductant than Cd. It is not clear which of Zn or Cr is the stronger reductant. So the order in alternative **D** is correct with Zn being omitted.

### Question 21 (1 mark)

Which additional experiment must be conducted to place all four metals ion order of their reducing strength?

A. Zinc and copper (I) nitrate solution.

**B. Chromium and zinc nitrate solution.**

C. Copper and cadmium

D. Cadmium and chromium nitrate solution.

**Question 14 B**




To determine which of Zn or Cr is the stronger reductant, a further experiment using chromium ions with zinc, or zinc ions with chromium, will need to be undertaken.


**Question 22** (1 mark)

*Inspired from VCAA Chemistry Exam 2008*

<https://www.vcaa.vic.edu.au/Documents/exams/chemistry/2008chem2-w.pdf#page=9>

The following reactions occur spontaneously as written.






Using this information, predict which one of the following pairs of reactants will react spontaneously.

- A.  $\text{Co}(\text{s}) + \text{Fe}^{2+}(\text{aq})$
- B.  $\text{Cr}^{2+}(\text{aq}) + \text{Fe}^{2+}(\text{aq})$
- C.  $\text{Cr}^{2+}(\text{aq}) + \text{Pb}^{2+}(\text{aq})$**
- D.  $\text{Pb}(\text{s}) + \text{Co}^{2+}(\text{aq})$

**Question 23** (1 mark)

The following reactions occur spontaneously



Using this information, predict which one of the following pairs of reactants will react spontaneously.

- A.  $\text{Co}(\text{s})$  and  $\text{Ce}^{3+}(\text{aq})$ .
- B.  $\text{Cr}^{3+}(\text{aq})$  and  $\text{Hg}(\text{l})$ .
- C.  $\text{Co}^{2+}(\text{aq})$  and  $\text{Ce}^{4+}(\text{aq})$ .
- D.  $\text{Hg}^{2+}(\text{aq})$  and  $\text{Cr}^{2+}(\text{aq})$ .**

Space for Personal Notes





**Question 24** (1 mark)

Inspired from VCAA Chemistry Exam 2020

<https://www.vcaa.vic.edu.au/Documents/exams/chemistry/2020/2020chem-w.pdf#page=15>

Consider the following half-equation.



It is also known that:

➤  $\text{ClO}_2(\text{g})$  will oxidise  $\text{HI}(\text{aq})$ , but not

➤  $\text{Fe}^{3+}(\text{aq})$  will oxidise  $\text{HI}(\text{aq})$ , but not

Based on this information, which of the



**A.**  $\text{Cl}_2(\text{g})$  and  $\text{I}_2(\text{aq})$ .

**B.**  $\text{Cl}_2(\text{g})$ , but not  $\text{ClO}_2(\text{g})$ .

**C.**  $\text{ClO}_2(\text{g})$  and  $\text{Cl}_2(\text{g})$ , but not  $\text{I}_2(\text{aq})$ .

**D.**  $\text{Cl}_2(\text{g})$ ,  $\text{ClO}_2(\text{g})$  and  $\text{I}_2(\text{aq})$ .

According to information from the electrochemical series



Based on the information supplied in the question,  $\text{ClO}_2/\text{ClO}_2^-$  is located above

$\text{Fe}^{3+}/\text{Fe}^{2+}$  but below  $\text{Cl}_2/\text{Cl}^-$ , i.e. the order of the redox pairs in terms of decreasing oxidising agent strength / increasing reducing agent strength) is:

➤  $\text{Cl}_2(\text{g})/\text{Cl}^-(\text{aq})$ ,

➤  $\text{ClO}_2(\text{g})/\text{ClO}_2^-(\text{aq})$

➤  $\text{Fe}^{3+}(\text{aq})/\text{Fe}^{2+}(\text{aq})$

➤  $\text{I}_2(\text{s})/\text{I}^-(\text{aq})$

Hence,  $\text{Fe}^{2+}(\text{aq})$  will be oxidised by  $\text{Cl}_2(\text{g})$  and  $\text{Cl}^-(\text{aq})$  but not by  $\text{I}_2(\text{s})$ .

Space for Personal Notes

Use the following information to answer the questions below.

Various reagents were mixed in separate flasks as shown in the table below.

Flask 1	Flask 2	Flask 3	Flask 4
$\text{Cu}(\text{NO}_3)_2(\text{aq}) + \text{Sn}$	$\text{Ag}^+(\text{aq}) + \text{Cd}$	$\text{Fe}^{3+}(\text{aq}) + \text{NaCl}(\text{aq})$	$\text{I}_2 \text{ solution} + \text{Cu}$

**Question 25** (1 mark)

A reaction is likely to occur in:

**A.** Flasks 1 and 2 but not in flask 3

**B.** Flasks 1 and 3 but not in flask 2

**C.** Flask 2 but not in flasks 1 and 3

**D.** Flask 3 but not in flasks 1 and 2.

**Question 25**

**A**

$\text{Cu}^{2+}$  ions are stronger oxidising agents than  $\text{Sn}^{2+}$  ions and so flask 1 contains the stronger oxidising agent and stronger reducing agent (Sn); therefore a reaction is predicted.  $\text{Ag}^+$  ions are stronger oxidising agents than  $\text{Cd}^{2+}$  ions and so flask 2 contains the stronger oxidising agent and stronger reducing agent (Cd); therefore a reaction is predicted.  $\text{Fe}^{3+}$  ions are weaker oxidising agents than  $\text{Cl}_2$ . Flask 3 contains the weaker oxidising agent and weaker reducing agent ( $\text{Cl}^-$ ); therefore no reaction is predicted.

**Question 26** (1 mark)

Using the electrochemical series, a reaction is predicted to occur in flask 4. However, no reaction had occurred by the time any reactions took place in the other flasks. Which one of the following is the most likely reason to explain this?

**A.** The iodine was in a different state

**B.** The enthalpy change for the reaction was small

**C.** An alloy of copper and zinc was used

**D.** The products are formed much more slowly than products in the other reactions.

**Question 26**

**D**

The state of the iodine should not make a large difference in predictions, provided that iodine molecules were available for reaction. **A** is incorrect. There is a difference of 0.20 V ( $+0.54 - (+0.34)$ ) in the  $E^\circ$  values for the two half-reactions. An exothermic reaction ( $\Delta H < 0$ ) is predicted to occur as the stronger oxidant ( $\text{I}_2$ ) is mixed with the stronger reductant (Cu). **B** is also incorrect. The alloy of copper is still a source of copper atoms which should react with iodine according to the electrochemical series. **C** is incorrect. The electrochemical series does not predict the rate of reactions and so it is most likely that the predicted reaction is slow. Hence no reaction was evident in a short period of time. **D** is the required answer.

Space for Personal Notes

## Section H: VCAA-Level Questions II (10 Marks)

INSTRUCTION: 10 Marks. 30 Seconds Reading. 9 Minutes Writing.




### Question 27 (10 marks)


Gaseous hydrogen peroxide is bubbled into aqueous chloride ions, all at SLC.

a. A reaction is begins to occur.

i. Write the balanced half-equation for the reduction reaction. (1 mark)



ii. Write the overall reaction that takes place. (2 marks)



iii. Describe one observation expected as this reaction continues. (1 mark)

- Bubbling due to oxygen gas forming.
- Heat being released due to the redox reaction being exothermic.

iv. After being left undisturbed for a long time, 40.0 L of gaseous products are collected at SLC. Calculate the mass of hydrogen peroxide which must have reacted. (2 marks)

$$\begin{aligned} n(\text{O}_2) &= \frac{V}{V_m} = \frac{40.0}{24.8} = 1.61 \text{ mol} \\ n(\text{H}_2\text{O}_2) &= 2 \times n(\text{O}_2) = 2(1.61) = 3.23 \text{ mol} \\ m(\text{H}_2\text{O}_2) &= n \times Mr = 3.23 \times (2 + 32) = 54.8 \text{ g} \end{aligned}$$

- v. Even after 20 minutes of the system being set up, there is no observable indication of the reaction described in **part i-iii.** occurring. Explain this phenomenon. (1 mark)


The reaction is occurring but extremely slowly and thus the observations are negligible. This is because the electrochemical series does not dictate the rate of the reaction.

- b. When manganese dioxide,  $\text{MnO}_2$ , is added to the hydrogen peroxide, the reaction occurs more rapidly. If the products are filtered at the end of the reaction, the  $\text{MnO}_2$  can be recovered and reused.

Explain the role of the  $\text{MnO}_2$ . (1 mark)

It is a catalyst to speed up the rate of reaction.

Hydrogen peroxide can also react with concentrated nitric acid according to the following equation:



- c. Identify whether the nitric acid acts as the oxidant or the reductant. Justify your answer with reference to oxidation numbers. (2 marks)

Oxidant with a decrease in oxidation number of from +5 to +4.  
In  $\text{H}_2\text{O}_2$  the oxidation number of oxygen is -1. In  $\text{O}_2$  the oxidation number of oxygen is 0, an increase of 1. Increase in oxidation number is an oxidation process, hence  $\text{H}_2\text{O}_2$  is the reductant. The oxidation number of nitrogen decreases from +5 to +4, with acidified  $\text{NO}_3^-$  being the oxidant.


Space for Personal Notes

## Section I: Extension Questions (9 Marks)

### Question 28 (4 marks)

A sample of four unlabelled metals and 1 M solutions of the nitrate of the metals. The metals are labelled A, B, C, D and the solutions are labelled  $A^{2+}$ ,  $B^{2+}$ ,  $C^{2+}$ ,  $D^{2+}$ .

An experiment was carried out and the following observations were made:




- Metal A remains unchanged in all solutions.
- Metal B becomes coated with another metal when placed in solutions of  $D^{2+}$  and  $A^{2+}$  only.
- Metal C becomes coated with another metal when placed in solutions of  $A^{2+}$ ,  $B^{2+}$  and  $D^{2+}$ .

a. Consider the ranking obtained above.

- i. State the strongest oxidant and the strongest reductant. (1 mark)

Strongest oxidant =  $A^{2+}(aq)$   
 Strongest reductant = C(s)

- ii. Write the overall reaction between the species identified in **part a.i.** (1 mark)



b. Would a reaction occur between  $B^{2+}$  and D(s)? Justify your answer. (2 marks)

No, a reaction will not occur as D is a weak reductant and  $B^{2+}$  is a weak oxidant.


Space for Personal Notes

**Question 29** (5 marks)


Rochelle mixes a solution of  $\text{Zn}^{2+}(\text{aq})$ ,  $\text{Co}^{2+}(\text{aq})$ ,  $\text{Fe}^{2+}(\text{aq})$  and  $\text{Sn}^{2+}(\text{aq})$  into a beaker of pure deionised water.

Rochelle then adds a stick of pure calcium into the mixture.


- a. Identify the oxidation reaction in this situation. (1 mark)



- b. Identify the reduction reaction in this situation. (1 mark)



- c. After some time, a new reduction equation starts in the beaker. Identify the new reduction equation that would occur. (1 mark)



- d. Are these reactions exothermic or endothermic? Identify the difference between the two. (2 marks)

Exothermic. Exothermic reactions release energy into the environment and are usually from spontaneous reactions, whereas endothermic reactions absorb energy and are usually from non-spontaneous reactions.

Space for Personal Notes

VCE Chemistry  $\frac{3}{4}$

## Free 1-on-1 Support



### Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

<u>1-on-1 Video Consults</u>	<u>Text-Based Support</u>
<ul style="list-style-type: none"><li>➤ Book via <a href="https://bit.ly/contour-chemistry-consult-2025">bit.ly/contour-chemistry-consult-2025</a> (or QR code below).</li><li>➤ One active booking at a time (must attend before booking the next).</li></ul>	<ul style="list-style-type: none"><li>➤ Message <a href="tel:+61440137304">+61 440 137 304</a> with questions.</li><li>➤ Save the contact as "Contour Chemistry".</li></ul>

### Booking Link for Consults

[bit.ly/contour-chemistry-consult-2025](https://bit.ly/contour-chemistry-consult-2025)



### Number for Text-Based Support

[+61 440 137 304](tel:+61440137304)