



Website: [contoureducation.com.au](http://contoureducation.com.au) | Phone: 1800 888 300

Email: [hello@contoureducation.com.au](mailto:hello@contoureducation.com.au)

## VCE Chemistry $\frac{3}{4}$ Rates of Reaction [2.6] Workbook

### Outline:



#### Collision Theory

Pg 02-04

- Introduction to Collision Theory
- Correct Orientation

#### Frequency of Collisions

Pg 5-25

- Concentration and Pressure
- Proportion of Successful Collisions
- Inert Gas
- Surface Area

#### Proportion of Successful Collisions

Pg 26-42

- Sufficient Energy
- Temperature
- Catalyst

#### Measuring Rates of Reaction

Pg 43-53

- Measuring Rates of Reaction
- In-Depth Graphical Analysis

### Learning Objectives:

- ❑ CH34 [2.6.1] - Explain How Factors Increase Frequency of Collisions
- ❑ CH34 [2.6.2] - Explain How Temperature & Catalyst Affect the Proportion of Successful Collisions
- ❑ CH34 [2.6.3] - Graph Differences in Rate & Yield



## Section A: Collision Theory

### Sub-Section: Introduction to Collision Theory

*What is the rate of reaction?*


#### Rate of Reaction

➤ Definition: how \_\_\_\_\_ something is reacting.



*Before we look at rates of reactions, how do reactions occur in the first place?*

Discussion: What requirements need to be fulfilled for two reactants to react with each other?



- \_\_\_\_\_
- \_\_\_\_\_
- \_\_\_\_\_

**NOTE:** The idea of 'sufficient energy' will be explored later in the booklet!




*Let's first look at the idea of 'Correct Orientation'.*

## Sub-Section: Correct Orientation

### Exploration: Correct Orientation

➤ Consider the following reaction:




- You don't have to know **what** the correct orientation is!
- You only need to know that **there is** a particular correct orientation.

Space for Personal Notes




### Analogy: Jigsaw Puzzle

- Consider a jigsaw puzzle:



*What if you were blind & you couldn't feel the ridges of the puzzle piece?*

- Consider this scenario:



- How might you get the two pieces to fit together? \_\_\_\_\_
- How can success be achieved quicker? \_\_\_\_\_
- **Example:** Consider a **10 %** chance of a successful collision, and there are **10 collisions / sec.**

| <u>Total Frequency of Collisions</u> | <u>Expected Frequency of Successful Collisions</u> |
|--------------------------------------|--|
| <b>10 collisions/s</b>               |  |
| <b>100 collisions/s</b>              |  |

- Effect of the increased total frequency of collisions on the frequency of **successful** collisions:
- [increases] / [decreases] / [stays constant]



Space for Personal Notes

## Section B: Frequency of Collisions

*We need to increase the frequency of total collisions. But how do we do this?*

### Exploration: Increasing Frequency of Collisions

- Consider bumper cars at the carnival:



- Scenario: You have the **capability** to alter the **setup** of the bumper cars at the carnival.
- How can the **frequency of collisions** be increased?

| Bumper Cars | Collisions between Reactants |
|-------------|------------------------------|
|             |                              |
|             |                              |
|             |                              |

- The amount of **particles** ( $n$ ) and **volume** ( $V$ ) is related to \_\_\_\_\_ of the reacting species.


## Sub-Section: Concentration and Pressure



### Concentration and Pressure


➤ **Definition:** How \_\_\_\_\_ a mixture is.

| Concentration                                    | Pressure   |
|--|--|
| Used for [aqueous (aq)] / [gaseous (g)] mixtures | Used for [aqueous (aq)] / [gaseous (g)] mixtures |



### Exploration: Decrease in Volume

➤ Consider the **volume** of a gas mixture is **decreased**.



➤ The pressure of gas mixture: [increases] / [decreases] / [no change]

➤ Relationship:

$$P \propto \frac{1}{V}$$



### Exploration: Increase in Volume

- Consider a cup of orange juice which has water added to it.



- Concentration of Orange Juice: [increases] / [decreases] / [no change]

### Extension: Mathematical Relationship of Concentration/Pressure on Moles ( $n$ ) and Volume ( $V$ )



| Concentration                                       | Pressure  |
|---|---|
| $C = \frac{n}{V} \rightarrow C \propto \frac{n}{V}$ | $P = \frac{nRT}{V} \rightarrow P \propto \frac{n}{V}$ |

### Concentration on Rate



- To increase the frequency of **successful** collisions with **correct orientation**, the frequency \_\_\_\_\_ collisions must be [increased] / [decreased].
- Concentration/Pressure can be increased by:

| Amount ( $n$ )                              | Volume ( $V$ )                              |
|---|---|
| [increase] / [decrease] amount of particles | [increase] / [decrease] volume of container |

*Let's look at a question together!*


**Question 1 (3 marks) Walkthrough.**

The rate of reaction of a particular gaseous mixture was attempted to be altered by adding more reactant gas particles in same volume.

**a.** State what happens to the rate of reaction. (1 mark)

---

**b.** Explain your answer in **part a.** (2 marks)

---

---

---

---

---

**NOTE:** The hardest part of these questions is [identifying what happens to rate] / [explaining why], so be sure to explain every detail!


**Sample Response: Increase in Concentration or Pressure**


- When concentration/pressure is increased, **particles are moved:** [closer together] / [further apart]
- Frequency of total collisions: [increases] / [decreases]
- Frequency of fruitful/successful collisions with **correct orientation** collisions: [increases] / [decreases]
- Overall rate of reaction: [increases] / [decreases]





### Explanation Flow Chart for Frequency of Collisions



*Key Feature* → \_\_\_\_\_ *Freq. Collisions*  
 → *Frequency of* \_\_\_\_\_ *Collisions with Correct Orientation*  
 → *Rate of Reaction*

*Let's now have a look at a sample response from a student, and let's grade it together!*

#### **Question 2 (3 marks) Mark, Improve.**

How does increasing the volume of a gaseous mixture affect the rate of reaction?

#### **Student's Response:**

-  *As the volume of the gaseous mixture is increased, particles drift further apart, resulting in a lower concentration of the gaseous reactants.*
-  *This decreases the number of successful collisions with the correct orientation, thereby decreasing the rate of reaction.*

**Mark (/3) :** \_\_\_\_\_

**Comments:**

---



---



---

**Space for Personal Notes**



### Misconception

*"Increasing concentrations increase the number of collisions."*

*Truth: When responding, you need to relate to either the:*

- The number of collisions/unit time (e.g. number of collisions per second or per minute).
- The **frequency** of collisions.

**REMINDER:** Link the **total frequency** of collisions first, before linking to the **frequency of successful collisions** with correct orientation!



*Your Turn! (Write Full Sentences)*



### Question 3

Matthew drops a sheet of magnesium metal into a beaker containing 1.0 M sulphuric acid which will react to form magnesium sulphate and hydrogen gas.

State the effect on the rate of reaction if the 1.0 M sulphuric acid is replaced with 2.0 M sulphuric acid. Justify your answer with reference to collision theory.

---



---



---



---



---

Space for Personal Notes



*To save time, you can shorthand this one!*

**Question 4 (3 marks)**

In a solution mixture of hydrochloric acid and sodium hydroxide are mixed together. A large volume of water is added to the mixture.

Explain whether the reaction will take a longer or shorter time to react to completion.

---



---



---



---



---

**Question 5 Additional Question.**

If the volume of a reaction vessel is doubled for a gaseous reaction, what happens to the frequency of successful collisions per unit time?

- A.** It increases because the molecules have more space.
- B.** It decreases because the concentration of gas molecules decreases.
- C.** It remains unchanged because only temperature affects successful collisions.
- D.** It increases because molecules gain more kinetic energy.

**Space for Personal Notes**

**Question 6 Additional Question.**

If the number of successful collisions per second in a gaseous reaction is doubled, what happens to the reaction rate?

- A. It remains the same because total collisions matter more than successful ones.
- B. It decreases because more molecules are colliding incorrectly.
- C. It increases as it means the total number of collisions per second has also doubled.
- D. It doubles because successful collisions are directly linked to the rate of reaction.

**Space for Personal Notes**


## Sub-Section: Proportion of Successful Collisions

**Discussion:** How does increasing the concentration/pressure change the proportion or probability of fruitful/successful collisions?

[increases] / [decreases] / [no change]

### Analogy: Darts

➤ Scenario: You were **blindfolded** playing darts and you had a **10% chance** of hitting a **bullseye**.



|  | <u>10 Darts Thrown Per Minute</u> | <u>100 Darts Thrown Per Minute</u> |
|--|-----------------------------------|------------------------------------|
| Expected Number of Bullseyes Hit Per Minute: |                                   |                                    |
| Number of Bullseyes Hit                      | [greater] / [lesser]              | [greater] / [lesser]               |
| Probability of Hitting a Bullseye            |                                   |                                    |

Space for Personal Notes



### Proportion/Probability of Successful Collisions for Concentration/Pressure Change

- As the **frequency** of total collisions is increased, the **frequency** of successful collisions is increased!


***DO NOT MENTION PROBABILITY/PROPORTION  
OF SUCCESSFUL COLLISIONS***

- Probability/Proportion of Successful Collisions will be properly covered later!

*Try some questions!*


#### Question 7 (5 marks)

There are two beakers of 1000 mL each which contain differing concentrations of ethene ( $C_2H_4$ ). In beaker A, 1.0 M concentration of  $C_2H_4$  is used, and in beaker B, 2.0 M concentration is used.



One drop of 0.100 mol bromine liquid ( $Br_2(l)$ ) is dropped into each beaker and the time taken for a reaction to occur is recorded.

The reaction that takes place is as follows:



- a. State which beaker will require the most amount of time for the brown colour to fade, giving justification for your response. (3 marks)

---



---




---



---

The experiment is then repeated with a slightly different setup.

- b. This time, beakers *A* and *B* both have a 1.00 M concentration of ethene ( $C_2H_4$ ), however, different amounts of bromine ( $Br_2$ ) is added to each beaker, as shown in the diagram below.



- i. Given that the volume of water in the beakers is still 1000 mL, state which beaker will have a greater concentration of bromine ( $Br_2$ ) added.

---

- ii. Explain which beaker will require the most amount of time for the colour to fade. (2 marks)

---



---



---



---

**NOTE:** As we need to wait for all the bromine to react, adding more bromine results in a quicker rate of reaction, but as more bromine needs to react, the overall time required for the brown colour to fade is longer.



Space for Personal Notes

## Sub-Section: Inert Gas



### Context

- Sometimes, an inert gas is added to the mixture.

*Let's have a look at what happens when we add an inert gas to a gas reaction chamber!*



Discussion: What are some examples of inert gases?



Space for Personal Notes





### Analogy: Bumper Cars

- Consider the following bumper cars with **just your friends** compared to **with random people**:

| Scenario #1: Just Friends  | Scenario #2: Friends & Strangers                        |
|--|---|
| <p>Scenario #1</p> <p style="text-align: center;">Vs</p> <p style="text-align: center;">Friends      Strangers</p> |   |
| Overall Concentration of Cars: [greater] / [lesser]  | Overall Concentration of Cars: [greater] / [lesser]     |
| Overall Frequency of Bumps: [greater] / [lesser]   | Overall Frequency of Bumps: [greater] / [lesser]        |
| Concentration of Friends: [greater] / [lesser] / [same]  | Concentration of Friends: [greater] / [lesser] / [same] |

- In scenario #2 v/s Scenario #1:

➤ Concentration of Friends: [greater] / [lesser] / [same]

- Frequency of Bumps between Friends: [greater] / [lesser] / [same]


Space for Personal Notes

Let's link to a real example!



### Exploration: Inert Gas in Reaction Chamber

- Consider the following reaction:



**Overall:**

- Overall Concentration of Container: [increases] / [decreases] / [stays same]
- Overall Frequency of Collisions: [increases] / [decreases] / [stays same]

**Just between NO and O<sub>2</sub>:**

- Concentrations of NO and O<sub>2</sub>: [increases] / [decreases] / [stays same]
- Frequency of collisions between NO and O<sub>2</sub>: [increases] / [decreases] / [stays same]
- Rate of reaction between NO and O<sub>2</sub>: [increases] / [decreases] / [stays same]
- Conclusion:

⚙ While overall concentration has increased,

⚙ The \_\_\_\_\_ of NO and O<sub>2</sub> has [increased] / [decreased] / [stayed same]

⚙ The **frequency of collisions** between them [increase] / [decrease] / [stays same]

➤ Rate of reaction: [increase] / [decrease] / [stays same]


**NOTE:** Instead of 'partial concentration', as the mixture is a gaseous mixture, the term 'partial pressure' is used!





### Partial Pressure/Partial Concentration



- **Definition:** The pressure or concentration of \_\_\_\_\_ in a mixture of substances.
- **Example:** Consider 1.00 L of water which has had 3.00 mol each of substance A and B added to it.



1.00 L water

-  The overall concentration is: \_\_\_\_\_
-  The concentration of only substance A is: \_\_\_\_\_

### Effect of Inert Gas on Rate of Reaction



- When an inert gas is added, the **overall pressure**: [increases] / [decreases] / [stays same]
- Partial pressure of reactants: [increases] / [decreases] / [stays same]
- Frequency of collisions between reactants: [increases] / [decreases] / [stays same]
- Rate of reaction: [increases] / [decreases] / [stays same]

Space for Personal Notes



**Extension:** Won't the inert gas get in the way of the collisions?

| No Inert Gas | Inert Gas |
|--------------|-----------|
|              |           |


- For every collision that is '**missed**' because an inert gas was in the way, a **new collision** that wouldn't have happened will occur!

*Your Turn!*



**Question 8** (3 marks)

Consider a reaction vessel containing nitrogen monoxide and oxygen which react in the following equation:



Neon gas is then added to the reaction chamber.

- a. Identify the effect of the addition of neon gas on the overall pressure of the reaction vessel. (1 mark)

---

- b. Explain what would happen to the rate of reaction. (2 marks)

---



---



---



---

Sub-Section: Surface Area



*How does surface area affect the rate of reaction?*



Analogy: Cutting Cake



➤ Consider a full v/s cut cake.



➤ Which cake has a greater surface area exposed?

[Full Cake] / [Cut Cake]


Space for Personal Notes

*Similar thing applies to chemicals!*



### Exploration: Surface Area

➤ Consider the following scenarios:



Scenario #1 : Solid lump

Scenario #2 : Broken down pieces

➤ In scenario #2 (broken down pieces):

- 🔄 Surface Area: [greater] / [lesser] / [same]
- 🔄 Frequency of Collisions: [greater] / [lesser] / [same]
- 🔄 Rate of Reaction: [greater] / [lesser] / [same]

Space for Personal Notes



Sample Response: Surface Area

- Cutting/dividing substance into thin powder [increases] / [decreases] surface area.
- Contact between reactants: [increases] / [decreases]
- Total Frequency of Collisions: [increases] / [decreases]
- Frequency of fruitful/successful collisions with correct orientation collisions: [increases] / [decreases]
- Overall rate of reaction: [increases] / [decreases]

*Try some questions!*




**Question 9** (5 marks)


Two experiments were conducted to investigate various factors that affect the rate of reaction between calcium carbonate and diluted hydrochloric acid. The two experiments are summarised in the diagrams below.

- a. Identify the rate-determining factor that is investigated in **Experiment 1**. (1 mark)

Experiment 1



- b. In **Experiment 2**, will the rate of reaction be faster in beaker *A* or beaker *B*? Explain your selection in terms of collision theory. (2 marks)




---

---

---

---

- c. Why is the following statement **incorrect**?

‘Collision theory states that all collisions between reactant particles will result in a chemical reaction.’  
(2 marks)

---

---

---

### Question 10 Additional Question.

If the surface area of a solid reactant is increased, what happens to the rate of reaction?

- A. The rate increases because more reactant particles are exposed to collisions.
- B. The rate decreases because molecules move more slowly.
- C. The rate remains unchanged because only temperature affects the reaction rate.
- D. The reaction stops because equilibrium is reached faster.



**Question 11 Additional Question.**

A student is conducting different experiments to investigate factors affecting reaction rates. Which of the following changes would result in an increased reaction rate?

- A. A reaction between hydrogen gas ( $\text{H}_2$ ) and oxygen gas ( $\text{O}_2$ ) occurs in a sealed container. The student increases the volume of the container while keeping the temperature constant.
- B. Adding argon gas to hydrogen and oxygen gas combustion reaction vessel.
- C. A reaction between magnesium ( $\text{Mg}$ ) and hydrochloric acid ( $\text{HCl}$ ) is performed. Instead of using a single strip of magnesium metal, the student uses finely powdered magnesium.
- D. A reaction between calcium carbonate ( $\text{CaCO}_3$ ) and hydrochloric acid ( $\text{HCl}$ ) is performed. The student removes some of the calcium carbonate from the flask, keeping all other conditions the same.

Space for Personal Notes

## Section C: Proportion of Successful Collisions

### Sub-Section: Sufficient Energy


#### Context

➤ Successful collisions are based on two factors:

- Colliding with the **correct orientation**.
- Colliding with **sufficient energy**.

Discussion: How much energy is sufficient energy?

#### Activation Energy




- **Definition:** Energy required to be \_\_\_\_\_ to break the pre-existing bonds before a reaction can proceed.
- **For Successful Reactions:** Energy \_\_\_\_\_ than the activation energy must be inputted.

*How can the proportion of particles which can overcome the activation energy be increased?*

**Exploration:** Factors that affect the energy required to overcome the activation energy.

- Consider a boy trying to run over a hill with a particular height (activation energy).



- How can we increase the chances of him being able to go completely over the hill?


| <u>Changes in Hill Scenario</u> | <u>Changes in Reactions</u> |
|---------------------------------|-----------------------------|
|                                 |                             |
|                                 |                             |

Space for Personal Notes


## Sub-Section: Temperature

### REMINDER

- Temperature is a type of kinetic energy!



Low temperature




High temperature

- At higher temperatures, particles move \_\_\_\_\_.

### Analogy: Usain Bolt

- Consider the two scenarios:

Usain Bolt



Average people



- Who will collide with greater force? [Usain Bolt] / [average people]
- Who is more likely to break their bones upon collision? [Usain Bolt] / [average people]

**Breaking Bones (Humans) ↔ Breaking Bonds (Chemicals)**

*Let's link this to an actual collision!*



### Exploration: Linking to an Actual Collision



### *Scenario: Temperature is Increased*

- Average kinetic energy: [increases] / [decreases]
- Impact upon collision: [more forceful] / [less forceful]
- Probability for sufficient energy to overcome the activation energy: [increases] / [decreases]
- Rate of reaction: [increases] / [decreases]

**NOTE:** Increasing temperature has increased the **proportion** of successful collisions as more particles have sufficient energy to react!

**ALSO NOTE:** While 'proportion/probability' should not have been mentioned before, it **should be mentioned now!**

### Exploration: Frequency of Collisions

- At greater temperatures, **particles move with greater speed.**
- What happens to the frequency of collisions? [more] / [less] frequent collisions.

Space for Personal Notes



### Effect of Temperature on Rate

➤ Increasing the temperature has a **dual** effect on the rate of reaction:

- It increases the \_\_\_\_\_ of particles with **sufficient energy** to react.
- It increases the \_\_\_\_\_ of successful collisions with the **correct orientation**.

*Which factor has the greatest effect on the rate of reaction?*

Discussion: Does energy upon collision or frequency of collisions affect the rate of reaction more?

[energy upon collision] / [frequency of collisions]

### Analogy: Punching

➤ Imagine people punching each other.



➤ Imagine someone punching someone with a **1% chance of knocking them out**.

| <u>Increasing Frequency of Collisions</u>                                 | <u>Increasing Energy Upon Collision</u>                                       |
|---|---|
| Punching someone <b>multiple times</b> with <b>1%</b> chance of knockout. | Punching someone <b>harder</b> with a <b>10%</b> chance of knocking them out. |



### Exploration: Sufficient Energy to React vs Frequency of Collisions

- Suppose a scenario with **1000 particles**, whereby **10%** of them have sufficient energy to react.
- How many particles out of **1000** have **sufficient energy** to react? \_\_\_\_\_
- Let's consider two scenarios:
  - 🔗 **Scenario #1:** The frequency of collisions is doubled.
  - 🔗 **Scenario #2:** The proportion of successful collisions is doubled.
  - 🔗 **Scenario #3:** Both the frequency & proportion of successful collisions are doubled.

| <u>Original</u>  | <u>Scenario #1: Increase Frequency of Collisions</u>  | <u>Scenario #2: Increase Proportion of Successful Collisions</u>   | <u>Scenario #3: Increase in Both</u>  |
|--|---|--|---|
| 10% of particles react.  | 10% of particles react at double the rate.  | 20% of particles react.  | 20% of particles react at double the rate.  |
| <div style="text-align: center;"> <b>1000</b><br/> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> </div> | <div style="text-align: center;"> <b>1000</b><br/> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="text-align: center;">🔗</div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> </div> | <div style="text-align: center;"> <b>1000</b><br/> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> </div> | <div style="text-align: center;"> <b>1000</b><br/> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="text-align: center;">🔗</div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> ↓<br/> <div style="border: 1px solid black; width: 50px; height: 20px; margin: 5px auto;"></div> </div> |

- We see that a greater proportion of particles results in a greater rate of reaction!

Space for Personal Notes



### Greatest Effect on Rate of Reaction

- Greatest effect on the rate of reaction: [frequency of collisions] / [energy upon collision]
- When explaining the rate of reaction for temperature, \_\_\_\_\_ factors must be referenced!



### Sample Response: Effect of Temperature

- **Overall:** Increasing temperature [increases] / [decreases] average kinetic energy.
- **Energy Upon Collision:**
  - ⚙ Reacting particles collide with [greater] / [lesser] force.
  - ⚙ Probability of colliding with sufficient energy to overcome the activation energy.
- **Frequency of Collisions:**
  - ⚙ Proportion/probability of successful/fruitful collisions. [increases] / [decreases] / [same]
  - ⚙ Average moving speed of particles: [increases] / [decreases] / [same]
  - ⚙ Total frequency of collisions: [increases] / [decreases] / [same]
  - ⚙ Frequency of successful collisions: [increases] / [decreases] / [same]
- **Overall Rate:** [increases] / [decreases]

Space for Personal Notes



**Question 12 (3 marks) Mark, Improve.**

How does increasing the temperature of a gaseous mixture affect the rate of reaction?

**Student's Response:**

✚ *As the temperature of a gaseous mixture is increased, the kinetic energy of the reactants is increased, resulting in a greater proportion of particles which can successfully react.*

✚ *Additionally, particles move at a greater speed, resulting in more frequent collisions. This results in a greater rate of reaction.*

**Mark (/3) :** \_\_\_\_\_

**Comments:**


---



---



---

**Energy Upon Collision Flow Chart**


\_\_\_\_\_ Energy → Greater \_\_\_\_\_  
 → Greater \_\_\_\_\_ for Collisions with \_\_\_\_\_  
 \_\_\_\_\_ → Greater Rate

**NOTE:** As the increase in the proportion of particles with sufficient energy has the \_\_\_\_\_ effect on the rate of reaction, it is best to mention this point \_\_\_\_\_.



**ALSO NOTE:** '\_\_\_\_\_ Kinetic Energy' must be referenced and not 'kinetic energy'.

Space for Personal Notes

**Question 13** (5 marks)

Explain the following phenomena using collision theory:

- a. The chemicals mixed by a panel beater make body filler harden faster on a hot day than on a cold day. (3 marks)

---

---

---

---

---

---

---

- b. Nail polish remover evaporates faster if you shake your fingers than if you don't. (2 marks)

---

---

---

---

Space for Personal Notes

**Question 14 Additional Question.**

What factor is the major reason for the increased rate of reaction when temperature is increased?

When the temperature of a chemical reaction is increased, the rate of reaction also increases. Which of the following factors has the greatest effect in explaining this increase?

- A. The reactant molecules move faster, leading to a greater frequency of collisions.
- B. The total number of reactant molecules increases, providing more particles to react.
- C. The increase in temperature increases the concentration of the reactants, making collisions more likely.
- D. The proportion of reactant molecules with energy greater than or equal to the activation energy increases, leading to more successful collisions.

Space for Personal Notes

## Sub-Section: Catalyst




### Catalyst

- **Definition:** A substance that \_\_\_\_\_ the rate of a chemical reaction without itself being \_\_\_\_\_.
- **How:** Catalysts alter the rate of reaction by providing an \_\_\_\_\_ reaction pathway with a \_\_\_\_\_ activation energy.

*How does a catalyst do this? What does this even mean?*

### Analogy: Paper

- Imagine trying to tear a piece of paper with one hand...





- This is how a reaction takes place normally.
- **Catalyst is present:** It simply acts as another \_\_\_\_\_.
- **Catalyst Purpose:** Helps to \_\_\_\_\_ the other end of the paper, allowing us to tear the paper much more easily.



### Exploration: How a Catalyst Works

➤ Consider the following reaction:



- Reactants **adsorb** onto the iron surface. They form **temporary** and **partial** intermolecular bonds with the catalyst.
- As reactants **attach**: Covalent bonds within their molecules \_\_\_\_\_.
- Activation Energy: [increases] / [decreases] / [stays same]
- Rate of Reaction: [increases] / [decreases] / [stays same]


Space for Personal Notes



## Catalysts

- A catalyst provides an alternative reaction pathway with lower activation energy by forming **temporary** and **partial intermolecular bonds** with the reacting particles.
- **Activation Energy:** [increases] / [decreases] / [stays same]
- **Rate of Reaction:** [increases] / [decreases] / [stays same]
- **Catalyst Before vs After:** \_\_\_\_\_.

## Exploration: Catalysts - Energy Profile Diagram




- Overall change in enthalpy ( $\Delta H$ ) with catalyst: [increases] / [decreases] / [no change]

Space for Personal Notes



### Analogy: Math Test

- We see that **increasing temperature** and using a **catalyst** both **increase the proportion of particles** with sufficient energy to overcome the activation energy.
- **Setup:** Imagine a math test with an **average score** of **60%** and a **high distinction** being **90%**.

 Do a lot of people get a high distinction? [yes]/[no]

| <u>Scenario #1: Increasing Temperature</u>  | <u>Scenario #2: Using Catalyst</u>  |
|---|---|
| Increases Average Kinetic Energy.   | Lowers Activation Energy.   |
| Average Test Score: <b>60% → 75%</b>  | Average Test Score: <b>60%</b>  |
| High Distinction Requirement: <b>90%</b>  | High Distinction Requirement: <b>90% → 75%</b>                                    |
| Proportion of People with High Distinction:<br>[increases] / [decreases] / [same] | Proportion of People with High Distinction:<br>[increases] / [decreases] / [same] |

- Temperature and adding a catalyst both increase the proportion of successful reactions, just in **different ways!**

### Sample Response: Effect of the Addition of a Catalyst



- The addition of a catalyst **lowers the activation energy** by providing an **alternative reaction pathway**.
- Proportion of particles with **sufficient energy** to overcome this **new activation energy** is **increased**.
- Probability/proportion of **fruitful/successful collisions** is **increased**, increasing the overall rate of reaction.

Space for Personal Notes

Try some questions!




Question 15 (1 mark)








Inspired from VCAA Chemistry Exam 2015

<https://www.vcaa.vic.edu.au/Documents/exams/chemistry/2015/2015chem-w.pdf#page=6>

The oxidation of Sulfur dioxide is an exothermic reaction. The reaction is catalysed by Vanadium(V) Oxide.



Which one of the following energy profile diagrams correctly represents both the catalysed and the uncatalysed reaction?



Space for Personal Notes




**Question 16** (1 mark)

*Inspired from VCAA Chemistry Exam 2017*

<https://www.vcaa.vic.edu.au/Documents/exams/chemistry/2017/2017chem-w.pdf#page=2>

A catalyst:

- A. Slows the rate of reaction.
- B. Ensures the reaction is exothermic.
- C. Increases frequency of collisions between reacting particles.
- D. Provides an alternative pathway for the reaction with lower activation energy.

**NOTE:** A catalyst merely increases the proportion of particles that can successfully react - it does not affect the frequency of collisions!


**Question 17**

An experiment is done twice – once with a catalyst and once without a catalyst. When the experiment is run with a catalyst, the rate of reaction is seen to be greater. Explain why this is the case.

---



---



---



---

Space for Personal Notes


**Question 18 (1 mark) Additional Question.**

*Inspired from VCAA Chemistry Exam 2019*

<https://www.vcaa.vic.edu.au/Documents/exams/chemistry/2019/2019chem-w.pdf#page=5>

5 mL of Ethanol,  $\text{CH}_3\text{CH}_2\text{OH}$ , undergoes combustion in a test tube with a diameter of 1 cm. This experiment is performed in a fume cupboard. The temperature in the fume cupboard is 20°C. Which one of the following actions will reduce the rate of reaction?

- A. Perform the experiment in a test tube with a diameter of 2 cm.
- B. Increase the temperature in the fume cupboard to 25°C.
- C. Increase the volume of the Ethanol to 7 mL.
- D. Mix 2 mL of a dilute solution of Sodium hydroxide, NaOH, with the Ethanol.

**NOTE:** Sodium hydroxide doesn't do anything, and thus is effectively the same as adding water!



Space for Personal Notes





## Section D: Measuring Rates of Reaction

### Sub-Section: Measuring Rates of Reaction



#### Context

- Generally, rates of reaction are measured by monitoring changes which can be \_\_\_\_\_.
- In total, there are 4 main methods we use to measure the rate of reaction. They include observing the change over a period of time:

-  \_\_\_\_\_
-  \_\_\_\_\_
-  \_\_\_\_\_
-  \_\_\_\_\_


#### Space for Personal Notes

*Let's have a look at each of the methods in depth!*



### Exploration: Method 1 - Volume of Gas Evolved

- **Conditions:** Reaction produces a \_\_\_\_\_.
- **Setup:** Attach a **gas syringe** to capture the gaseous products.
- **Consider the reaction:**



*What does the graph look like?*

| <u>Volume against Time</u> | <u>Rate Against Time</u> |
|----------------------------|--------------------------|
| <p>Volume</p> <p>Time</p>  | <p>Rate</p> <p>Time</p>  |

**NOTE:** The rate of reaction is indicated by the \_\_\_\_\_ of the graph!




**ALSO NOTE:** The **rate vs time graph** is simply just the \_\_\_\_\_ of the volume vs time graph!

**Exploration: Method 2 - Change in Mass**



- **Conditions:** Reaction produces a **gas**.
- In the following acid-carbonate reaction, there are **gaseous products**:



- **Mass of the reaction vessel over time:** [increases] / [decreases] / [no change]
- **Reasoning:** \_\_\_\_\_

*What does the graph look like?*

| Mass Against Time | Rate Against Time |
|-------------------|-------------------|
|                   |                   |





### Exploration: Method 3 - Change in Temperature

- Almost every single reaction has a change in \_\_\_\_\_, whether it be **exothermic** or **endothermic**!
- Consider the following reaction:



- Type of Reaction: [endothermic] / [exothermic]
- Temperature Change: [increases] / [decreases] / [no change]

| Temperature Against Time   | Rate Against Time   |
|--|---|
|  |  |

Space for Personal Notes



### Exploration: Method 4 - Change in pH

- Most reactions which involve acids or bases will have a change in \_\_\_\_\_ over time.
- Consider the following reaction:



- **Acidity/Basic Changes:** [more] / [less] - [acidic] / [basic]
- **pH Change:** [increases] / [decreases] / [no change]
- As the pH scale is **logarithmic**, you will not be asked to plot the graph for pH!

### Method to Measure Rate of Reaction



| <u>Method Of Measuring Rate</u>     | <u>Conditions</u>                             |
|-------------------------------------|---|
| Change in Volume (Gas Syringe)      | Gaseous Products                              |
| Change in Mass (Weighing Scale)     | Gaseous Products                              |
| Change in pH (pH metre/indicator)   | H <sup>+</sup> or OH <sup>-</sup> used/formed |
| Change in Temperature (thermometer) | Reaction is endothermic/exothermic            |

Space for Personal Notes

## Sub-Section: In-Depth Graphical Analysis



**Active Recall:** How is the rate of reaction indicated in a graph?


\_\_\_\_\_



*Let's look at a question together!*


### Question 19 Walkthrough.

Nitric acid is added to a 10 g granular zinc, which reacts according to the following equation:




On the axes provided below, draw the new graph observed if:

- a. A catalyst was added.



- b. The zinc metal was used in large chunks rather than broken down into granular pieces.





**NOTE:** As both scenarios result in the same amount of product forming, the graph \_\_\_\_\_ at the same point!




*Your turn!*




### Question 20

Now consider the following reaction, which occurs at SLC in the presence of a manganese dioxide catalyst.




On the axes provided below, draw the new graph observed if:

- a. The reaction is carried out at 40°C.



- b. The manganese dioxide catalyst is removed.



*Now, let's consider a scenario where the amount of acid is changed as well!*


**Exploration: 1.0 M vs 2.0 M HCl as the Limiting Reagent**

- Consider a solution where there is 10.0 mL of 1.0 M and 2.0 M of HCl and an excess amount of zinc metal.



| Setup  | 10 mL of 1.0 M HCl   | 10 mL of 2.0 M HCl   |
|--|----------------------|----------------------|
| Rate of reaction                               | [slower] / [quicker] | [slower] / [quicker] |
| $n(\text{HCl})$                                |                      |                      |
| Mass of hydrogen gas ( $\text{H}_2$ ) produced | [greater] / [lesser] | [greater] / [lesser] |
| Change in the mass of the mixture              | [greater] / [lesser] | [greater] / [lesser] |

*There is a difference in the rate of reaction and the amount of  $\text{H}_2\text{(g)}$  produced! What does this look like on the graph?*




Rate vs Yield in Graphs


original

difference


rate  $\uparrow$ , yield - constant




rate  $\downarrow$ , yield - constant




rate  $\uparrow$ , yield  $\uparrow$




rate  $\downarrow$ , yield  $\downarrow$



rate - constant, yield  $\uparrow$



rate - constant, yield  $\downarrow$




*Your turn!*




### Question 21

Now consider a similar situation but now hydrochloric acid is now the **limiting reagent**.




Assuming the 'original' reaction used 10 mL of 1.0 M hydrochloric acid, compared to the 'original' graph, how would the new graph look like if:

- a. 10 mL of 2.0 M hydrochloric acid was used instead.



- b. 10 mL of 0.5 M hydrochloric acid was used instead.



Space for Personal Notes


**Question 22** (1 mark)

Inspired from VCAA Chemistry Exam 2016


<https://www.vcaa.vic.edu.au/Documents/exams/chemistry/2016/2016chem-amd-w.pdf#page=14>

A student set up an experiment to test the effect of different factors on the rate and extent of the reaction between a strong acid and marble chips (Calcium carbonate,  $\text{CaCO}_3$ ). In each trial, the mass of the flask and its contents were measured every 30 seconds from the instant the reactants were mixed.

The strong acid used was Hydrochloric acid, HCl. The equation for the reaction is as follows.



The results of the two trials were graphed on the same axes and are shown below.



In Trial 2, the student must have:

- A. Heated the 0.5 M HCl before adding it to the flask.
- B. Doubled the volume of 0.5 M HCl added to the flask.
- C. Used 100 mL of 0.5 M  $\text{H}_2\text{SO}_4$  instead of 100 mL of 0.5 M HCl.
- D. Used the same mass of marble but crushed it into a powder.



## Contour Checklist

- ☐ **Learning Objective: [2.6.1] - Explain how factors increase the frequency of collisions**

### Key Takeaways

| <u>Concentration</u>                              | <u>Pressure</u>                                   |
|---|---|
| Used for [aqueous (aq)] / [gaseous (g)] mixtures. | Used for [aqueous (aq)] / [gaseous (g)] mixtures. |

- ☐ To increase the frequency of **successful** collisions with **correct orientation**, the frequency \_\_\_\_\_ collisions must be [increased] / [decreased].

- ☐ **Concentration/Pressure** can be **increased** by:

| <u>Amount (<math>n</math>)</u>               | <u>Volume (<math>V</math>)</u>               |
|--|--|
| [increase] / [decrease] amount of particles. | [increase] / [decrease] volume of container. |

- ☐ **Increase in Concentration or Pressure**

- ☐ When concentration/pressure is increased, **particles move:** [closer together] / [further apart]
- ☐ **Frequency of total collisions:** [increases] / [decreases]
- ☐ **Frequency of fruitful/successful collisions with correct orientation** collisions: [increases] / [decreases]
- ☐ **Overall rate of reaction:** [increases] / [decreases]

- ☐ **Increase in Concentration or Pressure Flow Chart**

**Key Feature** → \_\_\_\_\_ **Freq. Collisions**  
 → **Frequency of** \_\_\_\_\_ **Collisions with Correct Orientation**  
 → **Rate of Reaction**

### □ Effect of Inert Gas on Rate of Reaction

- When an inert gas is added, the **overall pressure**: [increases] / [decreases] / [stays same]
- Partial pressure of reactants: [increases] / [decreases] / [stays same]
- Frequency of collisions between reactants: [increases] / [decreases] / [stays same]
- Rate of reaction: [increases] / [decreases] / [stays same]

### □ Surface Area

- Cutting/dividing substance into thin powder [increases] / [decreases] surface area.
- **Contact between reactants**: [increases] / [decreases]
- **Total frequency of collisions**: [increases] / [decreases]
- **Frequency of fruitful/successful collisions with correct orientation collisions**: [increases] / [decreases]
- **Overall rate of reaction**: [increases] / [decreases]

## □ Learning Objective: [2.6.2] - Explain how temperature & catalyst affect the proportion of successful collisions

### Key Takeaways

- Greatest effect on the rate of reaction: [frequency of collisions] / [energy upon collision]
- **Effect of Temperature on Rate (Sample Response)**
  - **Overall**: Increasing temperature [increases] / [decreases] average kinetic energy.

| <u>Energy Upon Collision</u>   | <u>Frequency of Collision</u>  |
|--|--|
| Reacting particles collide with [greater] / [lesser] force   | Average moving speed of particles: [increases] / [decreases] / [same]  |
| Probability of colliding with sufficient energy to overcome the activation energy [increases] / [decreases] / [same] | Total frequency of collisions: [increases] / [decreases] / [same]      |
| Proportion/probability of successful/fruitful collisions [increases] / [decreases] / [same]                          | Frequency of successful collisions: [increases] / [decreases] / [same] |

- ☐ Energy Upon Collision Flow Chart:

\_\_\_\_\_ Energy → Greater  
 → Greater \_\_\_\_\_ for Collisions with  
 \_\_\_\_\_ → Greater Rate

- ☐ **Catalysts** are substances that \_\_\_\_\_ the rate of a chemical reaction without itself being \_\_\_\_\_.
- ☐ Catalysts alter the rate of reaction by providing an \_\_\_\_\_ reaction pathway with a \_\_\_\_\_ activation energy.
- ☐ Catalyst Sample Response:
- ☐ A catalyst provides an alternative reaction pathway with lower activation energy by forming **temporary** and **partial intermolecular bonds** with the reacting particles.
  - ☐ **Activation Energy:** [increases] / [decreases] / [stays same]
  - ☐ **Rate of Reaction:** [increases] / [decreases] / [stays same]
  - ☐ Catalyst Before vs After: \_\_\_\_\_.




□ **Learning Objective: [2.6.3] - Graph differences in rate & yield**

**Key Takeaways**

| Method Of Measuring Rate            | Conditions                         |
|-------------------------------------|------------------------------------|
| Change in Volume (Gas Syringe)      | Gaseous Products                   |
| Change in Mass (Weighing Scale)     | Gaseous Products                   |
| Change in pH (pH meter/indicator)   | $H^+$ or $OH^-$ used/formed        |
| Change in Temperature (thermometer) | Reaction is endothermic/exothermic |

□ **Rate vs Yield in Graphs**





Website: [contoureducation.com.au](https://contoureducation.com.au) | Phone: 1800 888 300 | Email: [hello@contoureducation.com.au](mailto:hello@contoureducation.com.au)

VCE Chemistry  $\frac{3}{4}$

# Free 1-on-1 Support



## Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

| <u>1-on-1 Video Consults</u>   | <u>Text-Based Support</u>   |
|--|---|
| <ul style="list-style-type: none"><li>➤ Book via <a href="https://bit.ly/contour-chemistry-consult-2025">bit.ly/contour-chemistry-consult-2025</a> (or QR code below).</li><li>➤ One active booking at a time (must attend before booking the next).</li></ul> | <ul style="list-style-type: none"><li>➤ Message <a href="tel:+61440137304">+61 440 137 304</a> with questions.</li><li>➤ Save the contact as "Contour Chemistry".</li></ul> |

## Booking Link for Consults

[bit.ly/contour-chemistry-consult-2025](https://bit.ly/contour-chemistry-consult-2025)



## Number for Text-Based Support

[+61 440 137 304](tel:+61440137304)