CONTOUREDUCATION

Website: contoureducation.com.au | Phone: 1800 888 300

Email: hello@contoureducation.com.au

VCE Mathematical Methods ½ Quadratics [1.3]

Workbook

Outline:

 Basics of Quadratics Factorising Quadratics Perfect Squares Difference of Squares Completing the Square 	Pg 2-11	Graphs of Quadratic Equations ➤ Parabola and Symmetry ➤ Graphing Quadratics ➤ Finding a Rule of a Quadratic From a	Pg 18-27 a Graph
 Quadratic Equations Solving by Factorisation Quadratic Formula Discriminant 	Pg 12-17	 Advanced Algebra of Quadratics Quadratic Inequalities Hidden Quadratics 	Pg 28-34

Learning Objectives:

MM12 [1.3.1] - Find factorised form of quadratics.
 MM12 [1.3.2] - Find solutions and the number of solutions to quadratic equations.
 MM12 [1.3.3] - Graph and find rules from the graph of quadratic equations.
 MM12 [1.3.4] - Solving Quadratic Inequalities and hidden quadratics.



Section A: Basics of Quadratics

Sub-Section: Factorising Quadratics



Let's quickly revise how we factorised quadratics!



Context: Factorising

- Reversing the process of algebraic expansion is known as factorisation.
- When we write a quadratic as the product of two linear terms, we say we factor the quadratic.
 - What does factorising allow us to do?

Definition

Factorising Quadratics

$$y = (x - a)(x - b)$$

- Steps:
 - 1. Divide by the coefficient of the leading term. (If applicable)
 - **2.** Consider the factors of the constant term.
 - 3. (If Positive Constant Term): See which pair of factors can add up to the coefficient of the x term.
 (If Negative Constant Term): See which pair of factors can subtract from the coefficient of the x term.
 - 4. Construct the linear factors.

Space for Personal Notes





Onection	1	Walkthrough.
Question	1	waikuirougii.

Factorise $x^2 + 3x - 10$.

Active Recall



- Steps:
 - 1. _____ by the coefficient of the leading term. (If applicable)
 - 2. Consider the factors of the ______.
 - **3.** (If Positive Constant Term): See which pair of factors can add up to the coefficient of the *x* term. (If Negative Constant Term): See which pair of factors can subtract from the coefficient of the xterm.
 - **4.** Construct the ______.

Factorise the following expressions:

a.
$$x^2 + 4x + 3$$

b.
$$x^2 - 7x + 12$$

Question 3 Extension.

Express $x^2 - 4kx + 3k^2$ as the product of two linear factors.

Question 4 Walkthrough.

Factorise $2x^2 + 5x - 3$.



Factorise the following expressions:

a.
$$-x^2 + 5x - 6$$

b.
$$-6x^2 + x + 1$$

Question 6 Extension.

Write the expression $-6x^2 + kx + 2k^2$ as the product of two linear factors.



Sub-Section: Perfect Squares



Let's quickly revise perfect squares!



Perfect Squares

$$(a+b)^2 = \underline{\hspace{1cm}}$$

$$(a-b)^2 = \underline{\hspace{1cm}}$$

- Perfect squares are special quadratic expressions that are made up of two identical linear factors.
- In other words, when a linear factor is squared, it becomes a perfect square.

Question 7

Factorise the following expression using the perfect square formula.

$$x^2 - 14x + 49$$



Factorise the following expressions using the perfect square formula.

a.
$$x^2 + 4x + 4$$

b.
$$9x^2 - 12x + 4$$

Question 9 Extension.

Express $4x^2 - 20kx + 25k^2$ as a perfect square.

TIP: Identify the value of a and b using the form above.





Sub-Section: Difference of Squares



Let's quickly revise the difference between squares!



Difference of Squares

$$a^2 - b^2 = \underline{\hspace{1cm}}$$



Question 10 Walkthrough.

Factorise $x^2 - 4$.

Factorise the following expressions:

a. $x^2 - 25$

b. $4x^2 - 9$

Question 12 Extension.

Factorise $3x^2 - 25$.



Sub-Section: Completing the Square



Let's quickly revise completing the square!



Completing the Square



When we complete the square of a quadratic $x^2 + bx + c$, we write it in the form:

$$x^2 + bx + c = (\underline{})^2 - (\frac{b}{2})^2 + c$$

- Steps:
 - 1. We halve the coefficient of x.
 - **2.** Subtract the half of the coefficient of *x* squared outside the square bracket.

Question 13 Walkthrough.

Complete the square for $x^2 - 4x + 7$.

?

Active Recall: Complete the Square Steps

- Steps:
 - 1. We _____ the coefficient of x.
 - 2. Subtract the ______ outside the square bracket.



Complete the square for each quadratic.

a. $x^2 - 6x + 1$

b. $2x^2 + 20x + 3$

Question 15 Extension.

Complete the square for $2x^2 - 4kx + 2k^2 + 3$.

Key Takeaways



- \checkmark A perfect square is in the form of $x^2 + 2ax + a^2 = (x + a)^2$.
- ✓ The difference in squares is in the form of $a^2 b^2 = (a b)(a + b)$.
- Arr Complete the square form of $x^2 + bx + c = \left(x + \frac{b}{2}\right)^2 \left(\frac{b}{2}\right)^2 + c$.



Section B: Quadratic Equations

Sub-Section: Solving by Factorisation

Solving by Factorisation

$$(x-a)(x-b)=0$$

$$x = a \text{ or } b$$

- Steps:
 - 1. Factorise the quadratic.
 - **2.** Equate each factor to 0 and solve for x.

Question 16

Solve each of the following quadratic equations for x:

a.
$$x^2 - 5x + 4 = 0$$

b.
$$2x^2 - 5x - 7 = 0$$



Sub-Section: Quadratic Formula



<u>Discussion:</u> What do we do if the quadratic is not easy to factorise?



The Quadratic Formula



for
$$ax^2 + bx + c = 0$$

x =

Space for Personal Notes





Where does this come from?



Extension: Derivation of Quadratic Formula



- The quadratic formula can be algebraically derived from attempting to complete the square for the general form of a quadratic equation $ax^2 + bx + c = 0$.
 - Since $a \neq 0$, we can divide the whole equation by a to make the coefficient of x^2 equal to 1:

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0$$

Now, complete the square!

$$x^2 + \frac{b}{a}x + \frac{c}{a} =$$
______ = 0

• Rearrange the constant term to the other side!

$$\left(x+\frac{b}{2a}\right)^2=\underline{\hspace{1cm}}$$

Expand the RHS!

$$\left(x+\frac{b}{2a}\right)^2=$$

 \bullet Now, solve for x and see what happens!



NOTE: That was the ultimate satisfaction.



Question 17

Solve each of the following quadratic equations for x.

a.
$$x^2 - 3x - 7 = 0$$

b.
$$2x^2 + 3x - 1$$



Sub-Section: Discriminant

Active Recall: Quadratic Formula



<u>Discussion:</u> What happens if the inside of the root is negative/zero/positive?



The Discriminant



- Definition:
 - \bullet The discriminant, often denoted by Δ (Delta), is the part **inside** the square root of the quadratic formula.

$$discriminant = \Delta = b^2 - 4ac$$

if $\Delta > 0$, there are _____

if $\Delta = 0$, there is ______

if $\Delta < 0$, there are _____

Space for Personal Notes



Determine how many unique roots exist in each of the following quadratic equations:

$$x^2 - 4x + 20 = 0$$

Question 19 Extension.

Find the value(s) of m that makes $3x^2 + 4x = 2m$ have no real solutions.

Key Takeaways



- ✓ We can solve quadratic equations by first factorising.
- Alternatively, we can use the quadratic formula given by $x = \frac{-b \pm \sqrt{b^2 4ac}}{2a}$.
- \checkmark The discriminant is given by $b^2 4ac$ which dictates the number of solutions.




Section C: Graphs of Quadratic Equations

Sub-Section: Parabola and Symmetry

<u>Parabola</u>

- Definition:
 - The shape of the graph of a quadratic is known as a ______.



<u>Discussion:</u> What is the parabola symmetrical to?




Space for Personal Notes



Axis of Symmetry





axis of symmetry:
$$x = -\frac{b}{2a}$$

Question 20

Find the axis of symmetry of each of the following quadratic hence, the coordinate of turning point.

$$y = 2x^2 - 3x + 5$$

NOTE: When a question asks for coordinates, you must mention both the x- and y-value of the point.



<u>Discussion:</u> Given that $ax^2 + bx + c$ has an x-intercept of $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$, what do you notice about the line of symmetry at $x = -\frac{b}{2a}$?



Sub-Section: Graphing Quadratics



Turning Point Form



The turning point form of a quadratic is given by:

$$y = a(x - h)^2 + k$$

turning point = _____

The turning point form is obtained by **completing the square**.

<u>Discussion:</u> Can every quadratic be put into turning point form? (Does every quadratic have a turning point?)




Space for Personal Notes



Question 21 Walkthrough.

Sketch the graph of the following quadratic equation, labelling the coordinates of all axes intercepts and the turning point.


$$y = (x+1)^2 - 1$$



Question 22

Sketch the graph of the following quadratic equation, labelling the coordinates of all axes intercepts and the turning point.

$$y = -\left(x + \frac{3}{2}\right)^2 - 4$$






<u>Discussion:</u> What would the discriminant of $-\left(x+\frac{3}{2}\right)^2-4=0$ equation be? (Graph from part b. above.)

Question 23 Extension.

Sketch the graph of the following quadratic equation, labelling the coordinates of all axes intercepts and the turning point. Begin by putting the equation into turning point form.

$$y = -x^2 + 6x - 5$$



Intercept Form



The x-intercept form of a quadratic is given by:

$$y = a(x - b)(x - c)$$

x-intercepts: (b, 0) and (c, 0)


 \blacktriangleright The axis of symmetry is located exactly in the middle of the two x-intercepts.



Question 24 Walkthrough.

Sketch the graph of the following quadratic equation, labelling the coordinates of all axes intercepts, as well as the turning point.


$$y = x^2 - 3x + 2$$



Question 25

Sketch the graph of the following quadratic equation, labelling the coordinates of all axes intercepts, as well as the turning point.

$$y = -x^2 + 4x + 5$$






NOTE: When a is negative, the x-intercepts stay the same, but the **shape** of the parabola becomes a **negative** parabola instead.



Question 26 Extension.


Sketch the graph of the following quadratic equation, labelling the coordinates of all axes intercepts, as well as the turning point.

$$y = 4x^2 - 5x - 6$$



Space for Personal Notes







Let's try to do it the other way around!



Finding the Equation of a Quadratic

Form 1: Turning Point Form

$$y = a(x - h)^2 + k$$

- Recommended when a turning point is easy to identify.
- Form 2: *x*-intercept Form


$$y = a(x - b)(x - c)$$

 \bullet Recommended when both x-intercepts are easy to identify.

Question 27 Walkthrough.


Find the equations of the quadratics graphed below. Show your working.

a. Find the equation in turning point form.






b. Find the equation in intercept form.



Question 28


Find the equations of the quadratics graphed below. Show your working.

a.





h



NOTE: Never forget the a coefficient!



Key Takeaways



- \checkmark Every quadratic can be put into the turning point given by $y = a(x h)^2 + k$.
- ✓ Not all quadratic can be put into the *x*-intercept form given by y = a(x b)(x c).
- $\ensuremath{\checkmark}$ We can use x-intercept form or turning point form to find the rule.

Space for Personal Notes



Section D: Advanced Algebra of Quadratics


Sub-Section: Quadratic Inequalities



How can we tackle quadratic inequalities?



Quadratic Inequalities



- For quadratic inequalities, we always _____ the function.
- Steps:
 - **1.** Sketch the function.
 - **2.** See where the *y*-value is within the inequality.
 - **3.** Find the corresponding x-values.

200


<u>Discussion:</u> Why do we look at y-value < 1 if the function < 1?



Question 29 Walkthrough.

Solve $x^2 - 1 > 0$ for x.

Hint: This is the same question as asking, "For what values of x is the graph of $y = x^2 - 1$ greater than 0?"



Active Recall: Quadratic Inequalities

?

- 1. _____ the function.
- 2. See where the _____ value is within the inequality.
- **3.** Find the corresponding _____ values.



Solve each of the following for x:

a.
$$(x-2)(x-3) \ge 0$$

b.
$$2x^2 + 9x + 4 < 0$$



olve $kx^2 + 4x - 4 > 0$ for x , where $k \in R \setminus \{0\}$. Give two answers depending on whether $k > 0$ or $k < 0$	on 31 Extension.	
	$kx^2 + 4x - 4 > 0$ for x, where $k \in R \setminus \{0\}$. Give two answers depending on whether $k > 0$ or $k < 0$	



Sub-Section: Hidden Quadratics



Let's take a look at hidden quadratics!



Hidden Quadratics



Instead of:

$$af(x)^2 + bf(x) + c = 0$$

We can let f(x) = X to have:

$$aX^2 + bX + c = 0$$

Question 32 Walkthrough.

Solve
$$x^4 - 13x^2 + 36 = 0$$
 for x .



a. Solve
$$(x-2)^2 - 7(x-2) + 12 = 0$$
 for x .

b. Solve
$$x - 2\sqrt{x} - 15 = 0$$
 for x .



Question 34 Extension.

Solve $x^4 - 4x^2 - k = 0$ for x, where k is a real number.

Key Takeaways



- ☑ For quadratic inequalities, we always sketch.
- ✓ For hidden quadratics, look for the pattern of something and something squared.





Contour Check

	<u>Learning Objective</u> : [1.1.1] - Find factorised form of quadratics					
	Key Takeaways					
	Perfect square is in the form of					
	Differences of squares are in the form of $a^2 - b^2$					
	Complete the square form of $x^2 + bx + c =$					
<u>Learning Objective</u> : [1.1.2] - Find solutions and number of solutions to quadratic equations						
Key Takeaways						
	We can solve for quadratic equations by first					
	Alternatively, we can use the quadratic formula given by $x = $					
	The discriminant is given by which dictates the number of solutions.					
<u>Learning Objective</u> : [1.1.3] - Graph and find rules from the graph of quadratic equations						
Key Takeaways						
	Key Takeaways					
	Key Takeaways Every quadratic can be put into the turning point given by $y = $					

 \square We can use x-intercept form or turning point form to find the rule.



<u>Learning Objective</u>: [1.1.4] - Solving Quadratic Inequalities and hidden quadratics

Key Takeaways

- For quadratic inequalities, we always ______.
- ☐ For hidden quadratics, look for the pattern of something and something _____



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

VCE Mathematical Methods ½

Free 1-on-1 Support

Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

1-on-1 Video Consults	<u>Text-Based Support</u>
 Book via bit.ly/contour-methods-consult-2025 (or QR code below). One active booking at a time (must attend before booking the next). 	 Message +61 440 138 726 with questions. Save the contact as "Contour Methods".

Booking Link for Consults bit.ly/contour-methods-consult-2025



Number for Text-Based Support +61 440 138 726

