



Website: [contoureducation.com.au](https://contoureducation.com.au) | Phone: 1800 888 300

Email: [hello@contoureducation.com.au](mailto:hello@contoureducation.com.au)

## VCE Mathematical Methods ½

### Transformations [2.4]

#### Workbook

#### Outline:



#### Introduction to Transformations

Pg 2-8

- Image and Pre-Image
- Dilation
- Reflection
- Translation

#### Transformation of Points

Pg 9-20

- Basic Transformation of Points
- The Order of Transformations
- Interpreting the Transformation of Points

#### Transformation of Functions

Pg 21-28

- Applying Transformations to Functions
- Finding the Applied Transformations

#### Learning Objectives:




- MM12 [2.4.1] - Applying  $x'$  and  $y'$  Notation to Find Transformed Points, Find the Interpretation of Transformations and Altered Order of Transformations
- MM12 [2.4.2] - Find Transformed Functions
- MM12 [2.4.3] - Find Transformations From Transformed Function (Reverse Engineering)

## Section A: Introduction to Transformations

### Sub-Section: Image and Pre-Image

*What do we call an original coordinate and a transformed coordinate?*

#### Image and Pre-Image



- The original coordinate is called the Pre-image.
- The transformed coordinate is called the Image.

Pre-Image:  $(x, y)$

Image:  $(x', y')$

#### Question 1

It is known that  $(1, 4)$  transformed into  $(3, 5)$ . State the value of  $x'$  and  $y'$ .

$$\begin{aligned} x' &= 3 \\ \therefore y' &= 5 \end{aligned}$$


**NOTE:** The  $x'$  and  $y'$  notation will be used quite heavily!

## Sub-Section: Dilation



### Exploration: Dilation

► Consider the point below:



► Let's plot the coordinates:

📍 P1: Dilation by a factor 2 from the  $x$ -axis.

📍 P2: Dilation by a factor  $\frac{1}{2}$  from the  $x$ -axis.

📍 P3: Dilation by a factor 2 from the  $y$ -axis.

📍 P4: Dilation by a factor  $\frac{1}{2}$  from the  $y$ -axis.

### Dilation



Dilation by a factor  $a$  from the  $x$ -axis:  $y' = ay$

Dilation by a factor  $b$  from the  $y$ -axis:  $x' = bx$

**Question 2 Walkthrough.**

Find the image  $(x', y')$  after applying the following transformations to  $(x, y)$ .

Dilation by factor 2 from the  $x$ -axis.

$$(x, 2y)$$

Dilation by factor  $\frac{1}{3}$  from the  $y$ -axis.

$$(\frac{1}{3}x, 2y)$$

**Question 3**

Find the image  $(x', y')$  after applying the following transformations to  $(x, y)$ .

Dilation by factor  $\frac{1}{2}$  from the  $x$ -axis.

$$(x, \frac{1}{2}y)$$

Dilation by factor 4 from the  $y$ -axis.

$$(4x, \frac{1}{2}y)$$

**NOTE:** We are applying the transformations on  $(x, y)$  not  $(x', y')$ .




Space for Personal Notes

## Sub-Section: Reflection





### Exploration: Reflection

➤ Consider the point below:



➤ Let's plot the coordinates:

-  P1: Reflection in the  $x$ -axis.  $(x, -y)$
-  P2: Reflection in the  $y$ -axis.  $(-x, y)$

### Reflection



Reflection in the  $x$ -axis:  $y' = -y$


Reflection in the  $y$ -axis:  $x' = -x$

## Sub-Section: Translation





### Exploration: Translation

► Consider the point below:



► Let's plot the coordinates (ignore the scale):

 P1: Translation by 2 units in the negative direction of the  $x$ -axis.

 P2: Translation by 3 units in the negative direction of the  $y$ -axis.

### Translation



Translation by  $c$  units in the positive direction of the  $x$ -axis:  $x' = x + c$

Translation by  $d$  units in the positive direction of the  $y$ -axis:  $y' = y + d$

#### Question 4

Find the image  $(x', y')$  after applying the following transformations to  $(x, y)$ .

Translation by 3 units in the positive direction of the  $x$ -axis.  $(x+3, y)$

Translation by 2 units in the negative direction of the  $y$ -axis.  $(x+3, y-2)$

#### Key Takeaways



- ✓ The transformed point is called the image and is denoted by  $(x', y')$ .
- ✓ The dilation factor is multiplied by the original coordinates.
- ✓ Reflection makes the original coordinates the negative of their original values.
- ✓ Translation adds a unit to the original coordinates.



## Section B: Transformation of Points

### Sub-Section: Basic Transformation of Points

*Let's try to apply all types of transformations to a point!*

#### Question 5 Walkthrough.

Find the image  $(x', y')$  after applying the following transformations to  $(x, y)$ .

Dilation by a factor 2 from the  $x$ -axis.  $(x, 2y)$

Dilation by a factor 4 from the  $y$ -axis.  $(4x, 2y)$

Reflection in the  $x$ -axis.  $(4x, -2y)$

Translation by 2 units in the negative direction of the  $x$ -axis.  $(4x-2, -2y)$

Translation by 3 units in the positive direction of the  $y$ -axis.  $(4x-2, -2y+3)$

### Question 6

Find the image  $(x', y')$  after applying the following transformations to  $(x, y)$ .

Translation by 4 units in the positive direction of the  $x$ -axis.  $(x+4, y)$

Translation by 3 units in the negative direction of the  $y$ -axis.  $(x+4, y-3)$

Dilation by a factor of  $\frac{1}{5}$  from the  $x$ -axis.  $(x+4, \frac{1}{5}(y-3))$

Dilation by a factor of 2 from the  $y$ -axis.  $(2(x+4), \frac{1}{5}(y-3))$

Reflection in the  $x$ -axis.

$$(2(x+4), -\frac{1}{5}(y-3))$$

**NOTE:** Order Matters.



~~Question 7 Extension:~~

Find the image  $(x', y')$  after applying the following transformations to  $(x, y)$ .

Translation by  $a$  units in the negative direction of the  $x$ -axis.  $(x-a, y)$

Translation by  $b$  units in the positive direction of the  $y$ -axis.  $(x-a, y+b)$

Dilation by a factor  $c$  from the  $x$ -axis.  $(x-a, c(y+b))$

Dilation by a factor  $\frac{3}{a}$  from the  $y$ -axis.  $(\frac{3}{a}(x-a), c(y+b))$

Reflection in the  $x$ -axis.  $(\frac{3}{a}(x-a), -c(y+b))$

Space for Personal Notes

## Sub-Section: The Order of Transformations

**Discussion:** From the previous question, what happens when the translation is applied first?

Brackets  
↓  
BODMAS

*What is the order of transformations the same as?*

### The Order of Transformation

Order = BODMAS Order

#### Question 8 Walkthrough.

Consider the point  $(x, y)$  which was transformed into a point  $(3x + 6, y)$  by the transformation  $T$ .

Jennifer thinks the transformation was:

“Translation 6 units in the positive direction of the  $x$ -axis and dilation by a factor of 3 from the  $y$ -axis.”

Meanwhile, David thinks the transformation was:

“Dilation by a factor of 3 from the  $y$ -axis and translation 6 units in the positive direction of the  $x$ -axis.”

Who is correct? And why?

1. Dil 3 from  $y$   
2. 6 right  
↪  $3(x+2)$   
1. 2 right  
2. Dil 3 from  $y$

Question 9

Consider the point  $(x, y)$  was transformed into a point  $(2(x - 5), y)$  by the transformation  $T$ .

Mary thinks the transformation was:

“Translation 5 units in the negative direction of the  $x$ -axis and dilation by a factor of 2 from the  $y$ -axis.”

Meanwhile, Sam thinks the transformation was:

“Dilation by a factor of 2 from the  $y$ -axis and translation 5 units in the negative direction of the  $x$ -axis.”

Who is correct? And why?

1. 5 left  
2. Dil 2 from y

Question 10 Extension.

Consider the point  $(x, y)$  was transformed into a point  $(2ax + 6a, y)$  by the transformation  $T$ .

Jennifer thinks the transformation was:

“A translation by 3 units in the positive direction of the  $x$ -axis, followed by a dilation by a factor  $2a$  from the  $y$ -axis.”

Meanwhile, David thinks the transformation was:

“A dilation by a factor  $2a$  from the  $y$ -axis, followed by a translation by  $3a$  units in the positive direction of the  $x$ -axis.”

Who is correct? And why?

1. Dil  $2a$  from  $y$   
2.  $6a$  right  $\times$

$\rightarrow 2a(x+3)$

1. 3 right  
2. Dil  $2a$  from  $y$

Discussion: If the order is the same as the BODMAS order, how do we change the order of transformations?

Translation First:


Factorised Form

$$x' = 2(x+2)$$

Dilation First:

Expanded Form

$$x' = 2x+4$$



**Question 11 Walkthrough.**

The series of transformations, “a dilation by a factor  $\frac{1}{2}$  from the  $x$ -axis and a translation by 3 units up” yields the same result as the series of transformations, “a translation by  $a$  units up and a dilation by a factor  $b$  from the  $x$ -axis.” Find the values of  $a$  and  $b$ .

$$(x, \frac{1}{2}y)$$

$$(x, \frac{1}{2}y+3)$$

$$(x, y+a)$$

$$(x, b(y+a))$$

$$\frac{1}{2}y+3 = b(y+a)$$

$$\frac{1}{2}(y+6) = b(y+a)$$

$$\therefore a=6, b=\frac{1}{2}$$

Question 12

The series of transformations, "a dilation by a factor 4 from the  $y$ -axis, a reflection in the  $y$ -axis and a translation by 8 units left" yields the same result as the series of transformations, "a translation by  $c$  units right, a reflection in the  $y$ -axis and a dilation by a factor  $d$  from the  $y$ -axis." Find the values of  $c$  and  $d$ .

$$(4x, y)$$

$$(-4x, y)$$

$$(-4x-8, y)$$

$$(-d(x+c), y)$$

$$(x+c, y)$$

$$(-(x+c), y)$$

$$-4x-8 = -d(x+c)$$

$$-4(x+2) = -d(x+c)$$

$$\therefore c=2, d=4$$



Question 13 Extension.

The series of transformations, “a dilation by a factor 2 from the  $y$ -axis, a reflection in the  $y$ -axis, a dilation by a factor 2 from the  $x$ -axis, a translation by 4 units left and a translation by 6 units down”, yields the same result as the series of transformations, “a translation by  $c$  units right, a reflection in the  $y$ -axis, a dilation by a factor  $d$  from the  $y$ -axis, a translation  $k$  units down, and a dilation by a factor  $m$  from the  $x$ -axis.” Find the values of  $c$ ,  $d$ ,  $k$  and  $m$ .

$$(2x, y) \Rightarrow (-2x, y) \Rightarrow (-2x-4, y-6)$$

$$(x+c, y) \Rightarrow (-x+c, y) \Rightarrow (-d(x+c), y) \Rightarrow (-d(x+c), y-k)$$

$$-2x-4 = -d(x+c)$$

$$(-d(x+c), m(y-k))$$

$$-2(x+2) = -d(x+c) \Rightarrow \therefore c=2, d=2$$

$$2y-6 = m(y-k)$$


$$2(y-3) = m(y-k) \Rightarrow \therefore k=3, m=2$$

NOTE: Dilation factors don't change!



Space for Personal Notes

Sub-Section: Interpreting the Transformation of Points



Active Recall: Order of Transformation



Order = BODMAS Order

Question 14 Walkthrough.

Consider the transformation which maps:

$$x' = 2x + 4$$

$$y' = -3(y - 1)$$

Dilate by factor — from  
y/x axis

Translate — units L/R/U/D  
Say it fully

a. State the transformation in DRT (Dilation, Reflection, Translation) order.

1. Dil 2 from y
2. Dil 3 from x
3. Reflection in x
4. 4 right
5. 3 up

Expanded:

$$x' = 2x + 4$$

$$y' = -3y + 3$$

b. State the transformation in the translation first order.

1. 2 right
2. 1 down
3. Reflection in x
4. Dil 2 from y
5. Dil 3 from x

Factorised:

$$x' = 2(x + 2)$$

$$y' = -3(y - 1)$$



**NOTE:** Expanding or factorising changes the order of transformation.

### Question 15

Consider the transformation which maps:

$$x' = 3x + 6$$

$$y' = -2(y + 2)$$

a. State the transformation in DRT (Dilation, Reflection, Translation) order.

1. Dil 3 from y
2. Dil 2 from x
3. Reflection in x
4. 6 right
5. 4 down

$$x' = 3x + 6$$

$$y' = -2y - 4$$


b. State the transformation in the translation first order.

1. 2 right
2. 2 up
3. Reflection in x
4. Dil 2 from x
5. Dil 3 from y

$$x' = 3(x + 2)$$

$$y' = -2(y + 2)$$

Space for Personal Notes



Discussion: Could the order of  $x$  and  $y$  transformations change?

↪ Yes!



### Key Takeaways

- ✓ Transformations should be interpreted when  $x'$  and  $y'$  are isolated.
- ✓ The order of transformation follows the BODMAS order.
- ✓ To change the order of transformations, we either factorise or expand.

Space for Personal Notes

## Section C: Transformation of Functions

### Sub-Section: Applying Transformations to Functions

*Let's now work with Functions!*

#### Transformation of Functions

- The aim is to get rid of the old variables,  $x$  and  $y$ , and have the new variables,  $x'$  and  $y'$ , instead.

REPLACE OLD w/ NEW:

$$y = f(x) \rightarrow y' = f(x')$$

- Steps:

1. Transform the points.
2. Make  $x$  and  $y$  the subjects.
3. Substitute them into the function.

$$f\left(\frac{x'-4}{2}\right)$$

OLD var subject

$$x = \frac{x'-4}{2}$$

$$x' = 2x + 4$$

1. Dil 2 from  $y$
2. 4 right

#### Question 16 Walkthrough.

Apply the transformations given below to  $y = x^2$ .

Reflect in the  $y$ -axis.

$$(-x, y)$$

Translate 1 unit to the right.

$$(-x+1, y)$$

Dilate by a factor of 2 from the  $y$ -axis.

$$(2(-x+1), y)$$

① Find  $x', y'$

$$x' = -2x + 2$$

$$y' = y$$

② Make OLD  $x/y$  the subject

$$x = \frac{x'-2}{-2}$$

$$y = y'$$

③ REPLACE OLD w/ NEW

$$y = x^2$$

$$y' = \left(\frac{x'-2}{-2}\right)^2$$

③.5 Remove dashes

④ Make  $y$  Subject

$$y = \left(\frac{x-2}{-2}\right)^2$$

*Your turn!*



### Active Recall: Transformation of Functions

- The aim is to get rid of the old variables,  $x$  and  $y$ , and have the new variables,  $x'$  and  $y'$ , instead.

$$y = f(x) \rightarrow y' = f(x')$$

- Steps:

1. Transform the points.
2. Make  $x$  and  $y$  the subject.
3. Replace them into the function.

Space for Personal Notes

Question 17

Apply the following transformations to the functions given:

a.  $f(x) = x^2$

Dilation by factor 3 from the  $x$ -axis.  $(x, 3y)$

Reflect in the  $y$ -axis.  $(-x, 3y)$

Translate 3 units to the left.  $(-x-3, 3y)$

Dilate by a factor of 5 from the  $y$ -axis.  $(5(-x-3), 3y)$

$$x' = -5x - 15$$

$$y' = 3y$$

$$x = \frac{x' + 15}{-5}$$

$$y = \frac{y'}{3}$$

$$y = x^2$$

$$\frac{y'}{3} = \left( \frac{x' + 15}{-5} \right)^2$$

$$\therefore y = 3 \left( \frac{x+15}{-5} \right)^2 //$$

b.  $f(x) = \sqrt{x}$

Dilate by a factor of  $\frac{1}{4}$  from the  $y$ -axis.  $(\frac{1}{4}x, y)$

Dilate by a factor of 3 from the  $x$ -axis.  $(\frac{1}{4}x, 3y)$

Translate 4 units to the left.  $(\frac{1}{4}x - 4, 3y)$

Translate 1 unit up.  $(\frac{1}{4}x - 4, 3y + 1)$

Reflect in the  $y$ -axis.  $(-(\frac{1}{4}x - 4), 3y + 1)$

$$x' = \frac{1}{4}x + 4$$

$$y' = 3y + 1$$

$$x = -4(x' - 4)$$

$$y = \frac{y' - 1}{3}$$

$$y = \sqrt{x}$$

$$\frac{y' - 1}{3} = \sqrt{-4(x' - 4)}$$

$$\therefore y = 3\sqrt{-4x + 16} + 1 //$$

**Question 18 Extension.**

Apply the following transformations to  $y = 2^x$ .

Translation by 2 units to the right.  $(x+2, y)$

Reflection in the y-axis.  $(-(x+2), y)$

Dilation by a factor 3 from the y-axis.  $(-3(x+2), y)$

Translation by 3 units up.  $(-3(x+2), y+3)$

A dilation by a factor 2 from the x-axis.  $(-3(x+2), 2(y+3))$

A reflection in the x-axis.  $(-3(x+2), -2(y+3))$

$$\therefore x' = -3x - 6 \Rightarrow x = \frac{x' + 6}{-3}$$


$$\therefore y' = -2y - 6 \Rightarrow y = \frac{y' + 6}{-2}$$

$$\Rightarrow \begin{aligned} y &= 2^x \\ \Downarrow \\ \frac{y' + 6}{-2} &= 2^{\frac{x' + 6}{-3}} \end{aligned}$$

$$\begin{aligned} \therefore y &= -2 \cdot 2^{\frac{x' + 6}{-3}} - 6 \\ &= -2 \cdot 2^{\frac{x' + 3}{-3}} - 6 \end{aligned}$$



## Sub-Section: Finding the Applied Transformations



*Now let's go backwards!*



### Reverse Engineering



#### Steps:

1. Add the dashes (') back to the transformed function.
2. Make  $f( )$  the subject.
3. Equate the LHS of the original and transformed functions to the RHS of the original and transformed functions.
4. Make  $x'$  and  $y'$  the subjects and interpret the transformations.

#### Question 19 Walkthrough.

Find the transformations required for  $y = x^2$  to be transformed to  $y = 3\left(\frac{x+3}{2}\right)^2 + 5$ .

① Put ' back in  $y = x^2$

TF  $(x', y')$

$$y' = 3\left(\frac{x'+3}{2}\right)^2 + 5$$

② Move  $y$  transformations left

$$\frac{y'-5}{3} = \left(\frac{x'+3}{2}\right)^2$$

③ Equate OLD w/ NEW  $x$  &  $y$

$$y = \frac{y'-5}{3}$$

$$x = \frac{x'+3}{2}$$

$$y' = 3y + 5$$

$$x' = 2x - 3$$

④ Make  $x', y'$  the subject & READ

1. Dil 3 from  $x$   
2. 5 up

3. Dil 2 from  $y$   
4. 3 left

*Your turn!*



### Active Recall: Steps for reverse engineering

#### ► Steps:

1. Add the dashes (') back to the transformed function.
2. Make  $f( )$  the Subject.
3. Equate the LHS of the original and transformed functions to the RHS of the original and transformed functions.
4. Make  $x'$  &  $y$  the subjects and interpret the transformations.

Space for Personal Notes

**Question 20**

State a series of transformations (in order) that allow  $f(x)$  to be transformed into  $g(x)$ .

a.  $f(x) = 2(x+1)^2 + 3$  and  $g(x) = 6(x-4)^2 - 3$ .

$$y = 2(x+1)^2 + 3$$

$$y' = 6(x'-4)^2 - 3$$

$$\frac{y-3}{2} = (x+1)^2$$

$$\frac{y'+3}{6} = (x'-4)^2$$

$$\frac{y-3}{2} = \frac{y'+3}{6}$$

$$x+1 = x'-4$$

$$\therefore y' = 3y - 12$$

$$x' = x - 5$$

1. Dil 3 from  $x$

1. 5 left

2. 12 down

b.  $f(x) = 3(x-1)^2$  and  $g(x) = \frac{1}{2}(2x+3)^2 + 1$ .

$$y = 3(x-1)^2$$

$$y' = \frac{1}{2}(2x'+3)^2 + 1$$

$$\frac{y}{3} = (x-1)^2$$

$$2(y'-1) = (2x'+3)^2$$

$$\frac{y}{3} = 2y' - 2$$

$$x-1 = 2x'+3$$

$$y' = \frac{1}{6}y + 1$$

$$x' = \frac{1}{2}x - 2$$

1. Dil  $\frac{1}{6}$  from  $x$

1. Dil  $\frac{1}{2}$  from  $y$

2. 1 up

2. 2 left

Question 21 Extension.

Find a sequence of transformations required for  $y = 2(x - 3)^2 + 4$  to be transformed to  $y = -x^2 - 4x - 9$ .

$$(a+b)^2 = a^2 + 2ab + b^2$$

$$y = 2(x-3)^2 + 4$$

$$\frac{y-4}{2} = (x-3)^2$$

$$\frac{y-4}{2} = -y' - 5$$

$$y' = -\frac{y}{2} - 3$$

1. Dil  $\frac{1}{2}$  from  $x$
2. Reflection in  $x$
3. 3 down

$$\begin{aligned} y &= -(x^2 + 4x) - 9 \\ &= -((x+2)^2 - 4) - 9 \\ &= -(x+2)^2 - 5 \end{aligned}$$

$$-(y'+5) = (x'+2)^2$$

$$x-3 = x'+2$$

$$\therefore x' = x-5$$

1. 5 left

Key Takeaways

- ✓ We transform the coordinates first, then transform the function.
- ✓ To transform the function, replace its old variables with the new ones.
- ✓ To find the transformations, simply equate LHS with RHS after separating the transformations of  $x$  and  $y$ .





## Contour Checklist

- **Learning Objective: [2.4.1] - Applying  $x'$  and  $y'$  Notation to Find Transformed Points, Find the Interpretation of Transformations and Altered Order of Transformations**

### Key Takeaways

- The transformed point is called the image and is denoted by  $(x', y')$ .
- The dilation factor is multiplied to the original coordinate.
- Reflection makes the original coordinates the negative of their original values.
- Translation adds a unit to the original coordinate.
- Transformations should be interpreted when  $x'$  &  $y'$  are isolated.
- The order of transformation follows the BODMAS order.
- To change the order of transformations, we either factorise or expand.

- **Learning Objective: [2.4.2] - Find Transformed Functions**

### Key Takeaways

- To transform the function, replace its old variables with the new one.

□ **Learning Objective: [2.4.3] - Find Transformations From Transformed Function (Reverse Engineering)**

**Key Takeaways**

- To find the transformations, simply equate the LHS & RHS after separating the transformations of  $x$  and  $y$ .



Website: [contoureducation.com.au](https://contoureducation.com.au) | Phone: 1800 888 300 | Email: [hello@contoureducation.com.au](mailto:hello@contoureducation.com.au)

## VCE Mathematical Methods ½

# Free 1-on-1 Consults



### What Are 1-on-1 Consults?

- **Who Runs Them?** Experienced Contour tutors (45 + raw scores and 99 + ATARs).
- **Who Can Join?** Fully enrolled Contour students.
- **When Are They?** 30-minute 1-on-1 help sessions, after school weekdays, and all day weekends.
- **What To Do?** Join on time, ask questions, re-learn concepts, or extend yourself!
- **Price?** Completely free!
- **One Active Booking Per Subject:** Must attend your current consultation before scheduling the next. :)

**SAVE THE LINK, AND MAKE THE MOST OF THIS (FREE) SERVICE!**



### Booking Link

[bit.ly/contour-methods-consult-2025](https://bit.ly/contour-methods-consult-2025)

