

Website: contoureducation.com.au | Phone: 1800 888 300

Email: hello@contoureducation.com.au

VCE Mathematical Methods ½ Circular Functions I [4.1]

Workbook

Outline:

Introduction to Circular Functions

Pg 2-13

→ Radians and Degrees

- Unit Circle
- Period
- Pythagorean Identities
- Exact Values

Symmetry Pg 14-35

Supplementary RelationshipsComplementary Relationships

Solving Trigonometric Equations

Particular Solutions

Pg 36-42

Learning Objectives:

- MM12 [4.1.1] Evaluate Exact Values for Sine, Cosine, and Tangent
- MM12 [4.1.2] Applying Pythagorean Identity to Evaluate Trigonometric Functions
- MM12 [4.1.3] Apply Supplementary and Complementary Relationship to Evaluate Trigonometric Functions
- MM12 [4.1.4] Solve Particular Solution for Trigonometric Functions





Section A: Introduction to Circular Functions

Sub-Section: Radians and Degrees

Definition

Radians and Degrees

$$\mathbf{1}^c = \left(\frac{180}{\pi}\right)^0$$

$$1^o = \left(\frac{\pi}{180}\right)^c$$

$$180^{\circ} = \pi^{c}$$

Question 1

a. Find $\left(\frac{\pi}{4}\right)^c$ in degrees:

b. Find 12° in radians:

$$12^{\circ} \cdot \frac{\pi^{\circ}}{180^{\circ}} = \frac{1}{15} \pi$$

$$= \left(\frac{\pi}{15}\right)^{\circ} \pi$$



Sub-Section: Unit Circle



REMINDER: Don't forget!



$$sin = \frac{8}{H}$$

$$cos = \frac{4}{H}$$

Hypotenuse








Exploration: Unit Circle

- The unit circle is simply a circle of radius _________.
- lt can be divided into **four quadrants**:



- We can use the elementary definition of the trigonometric functions.
- Select the option below!

 $sin(\theta) = [X \ Value(Y \ Value), Gradient]$

 $cos(\theta) = X Value, Y Value, Gradient$


 $tan(\theta) = [X \ Value, Y \ Value, Gradient]$




Unit Circle



The unit circle is simply a circle of radius 1.








Sub-Section: Period




Discussion: What would happen if we rotate 360° to the value of sin and cos?





Discussion: What would happen if we rotate 180° to the value of tan?



REMINDER: Don't forget! Transformation



$$f(x) \to f(nx)$$

Dilation by factor $\frac{1}{n}$ from the y-axis.





Period of
$$sin(\underline{n}x)$$
 and $cos(\underline{n}x)$ functions $=\frac{2\pi}{n}$

Period of
$$tan(nx)$$
 functions = $\frac{\pi}{n}$

where
$$n = \text{coefficient of } x \text{ and } n > 0$$



Question 2

Find the period of each of the following trigonometric functions:

a.
$$p(x) = \tan\left(\frac{x}{2}\right)$$

$$P = \frac{\pi}{\left(\frac{1}{2}\right)} = 2\pi$$

b.
$$q(x) = \cos\left(\frac{3}{2}x + \frac{\pi}{3}\right)$$

$$P = \frac{2\pi}{\left(\frac{3}{2}\right)} = \frac{4\pi}{3} \eta$$

Ouestion 3 Extension.

Question 3 Extension.

Find the period of the following trigonometric function:

$$f(x) = \cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{3}\right)$$

$$f(x) = \cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{3}\right)$$

$$LCM(\pi, Gr) = 12\pi_{//}$$





Sub-Section: Pythagorean Identities




<u>Discussion:</u> What does $\sin^2(\theta) + \cos^2(\theta)$ equal to?





Pythagorean Identities





$$\sin^2(\theta) + \cos^2(\theta) = 1$$

Can be used for finding one trigonometry function by using the other.



Question 4

Find the value of cos (x) given that $sin(x) = \frac{2}{3}$ and x is first quadrant.

$$50^{2}(x) + cos^{2}(x) = 1$$

$$\frac{4}{9} + \cos^2(x) = 1$$

$$\omega s^2(x) = \frac{5}{9}$$

$$\frac{4}{9} + \cos^2(x) = 1$$

$$\cos^2(x) = \frac{5}{9}$$

$$\cos(x) = \frac{\pm \sqrt{5}}{3} \qquad \text{its in the 1st } 0.$$

$$\Rightarrow \cos(x) = \frac{\sqrt{5}}{3} = \sqrt{3}$$

$$\Rightarrow \cos(x) = \frac{\sqrt{5}}{3} / \sqrt{2}$$

NOTE: Consider the quadrant to determine signs.





Sub-Section: Exact Values



The Exact Values Table



x	0 (0°)	$\frac{\pi}{6}$ (30°)	$\frac{\pi}{4} (45^{\circ})$	$\frac{\pi}{3} \ (60^{\circ})$	$\frac{\pi}{2} \ (90^{\circ})$
sin(x)	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
$\cos(x)$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
tan(x)	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Undefined

TIP: Use the fact that sin is the y value, cos is the x value and the tangent is the gradient to remember the values well!







Question 5

Without looking at the exact value table, evaluate the following.

a. $\sin\left(\frac{\pi}{6}\right)$



b. $\cos\left(\frac{\pi}{3}\right)$

c. $\sin\left(\frac{\pi}{2}\right)$

d. $\tan\left(\frac{\pi}{6}\right)$








Exploration: Exact values

- Hold your left palm up facing yourself.
- Imagine the thumb to align with the positive y-axis, so the thumb is 9-axis.
- Imagine the pinky to align with the positive x-axis, so the pinky is _____.
- Can you guess what angles the index finger, middle finger and ring finger represent?
- Label the angles on a unit circle as well as the finger.



$$sin(\theta) = \frac{\sqrt{(the number of fingers below)}}{2}$$

$$cos(\theta) = \frac{\sqrt{(the\ number\ of\ fingers\ above)}}{2}$$





Active Recall: The Exact Values Table



x	0 (0°)	$\frac{\pi}{6}$ (30°)	$\frac{\pi}{4} (45^{\circ})$	$\frac{\pi}{3}$ (60°)	$\frac{\pi}{2} \ (90^{\circ})$
sin(x)	O	10	₹ <u>7</u>	42	1
$\cos(x)$	1	৸	1/2	专	O
tan(x)	0	162	1	13	undefind

Space for Personal Notes



Section B: Symmetry



Sub-Section: Supplementary Relationships


<u>Discussion:</u> Given that $\sin(30) = \frac{1}{2}$, how do we solve for $\sin(150)$?



What does Reflection in the y-axis look like?

Exploration: Reflection in y-axis

Consider the unit circle.



- Reflect the angle around the y-axis on the unit circle above.
- \blacktriangleright What is the angle in terms of θ ?



CONTOUREDUCATION

- How does that affect the sine / cosine / tangent functions?
- Scan the QR code below and have a look!



$$\cos(\pi - \theta) = +/\cos(\theta)$$

$$\sin(\pi - \theta) = -\sin(\theta)$$


$$\tan(\pi - \theta) = + /\cot(\theta)$$

CONTOUREDUCATION

Question 6


a. Find the angle at which $\frac{5\pi}{6}$ corresponds to in the first quadrant.

TG



b. What would $\sin\left(\frac{5\pi}{6}\right)$ equal to given that $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$?

こえり




c. What would $\cos\left(\frac{5\pi}{6}\right)$ equal to given that $\cos\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}$?

= -45/



d. What would $\tan\left(\frac{5\pi}{6}\right)$ equal to given that $\tan\left(\frac{\pi}{6}\right) = \frac{1}{\sqrt{3}}$?

· -! //



NOTE: Visualise on the unit circle.

ALSO NOTE: In the second quadrant, the y value (sin) is positive.






What does a Reflection in the x-axis look like?



Exploration: Reflection in x-axis

Consider the unit circle.



- Reflect the angle around the x-axis on the unit circle above.
- \blacktriangleright What is the angle in terms of θ ?
- How does that affect the sine / cosine / tangent functions?
- Scan the QR code below and have a look!



$$\cos(-\theta) = - \cos(\theta)$$


$$\sin(-\theta) = +\sin(\theta)$$

$$\tan(-\theta) = + \Box \tan(\theta)$$


CONTOUREDUCATION

Question 7


a. Find the angle at which $\frac{11\pi}{6}$ corresponds to in the first quadrant.



b. What would $\sin\left(\frac{11\pi}{6}\right)$ equal to given that $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$?




c. What would $\cos\left(\frac{11\pi}{6}\right)$ equal to given that $\cos\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}$?



d. What would $\tan\left(\frac{11\pi}{6}\right)$ equal to given that $\tan\left(\frac{\pi}{6}\right) = \frac{1}{\sqrt{3}}$?





NOTE: In the fourth quadrant, the x value (cos) is positive.






What does Reflection in both axes look like?



Exploration: Reflection in both axes

Consider the unit circle.



- Reflect the angle around both axes on the unit circle above.
- Mhat is the angle in terms of θ ? π + θ
- How does that affect the sine / cosine / tangent functions?
- Scan the QR code below and have a look!



$$\cos(\pi + \theta) = + \cos(\theta)$$


$$\sin(\pi + \theta) = + / \sin(\theta)$$

$$\tan(\pi + \theta) = + - \tan(\theta)$$

CONTOUREDUCATION


Question 8

a. Find the angle which $\frac{7\pi}{6}$ corresponds to in the first quadrant.




b. What would $\sin\left(\frac{7\pi}{6}\right)$ equal to given that $\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$?






c. What would $\cos\left(\frac{7\pi}{6}\right)$ equal to given that $\cos\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2}$?



d. What would $\tan\left(\frac{7\pi}{6}\right)$ equal to given that $\tan\left(\frac{\pi}{6}\right) = \frac{1}{\sqrt{3}}$?





NOTE: In the third quadrant, the gradient (tangent) is positive.






Let's summarise



Supplementary Relationships





- Simply look at the quadrant to find the correct sign.
 - **G** Second Quadrant $(\pi \theta)$

$$\cos(\pi - \theta) = -\cos(\theta)$$

$$\sin(\pi - \theta) = +\sin(\theta)$$

$$\tan(\pi - \theta) = -\tan(\theta)$$

• Third Quadrant $(\pi + \theta)$

$$\cos(\pi + \theta) = -\cos(\theta)$$

$$\sin(\pi + \theta) = -\sin(\theta)$$

$$\tan(\pi + \theta) = + \tan(\theta)$$



G Fourth Quadrant $(-\theta)$

$$\cos(-\theta) = +\cos(\theta)$$

$$\sin(-\theta) = -\sin(\theta)$$

$$\tan(-\theta) = -\tan(\theta)$$

- Steps
 - **1.** Equate to $\cos / \sin / \tan(\theta)$.
 - 2. Determine the sign (\pm) by considering the quadrant.

Try the following question!



TIP: Simply draw the unit circle.





Question 9 Walkthrough.

If $cos(\theta) = 0.8$ where θ is a first quadrant angle, evaluate the following.

C supp







b.
$$\sin(\pi - \theta)$$

$$\frac{\sin(\pi - \theta)}{y - \alpha^2} = \frac{0.6}{4}$$

$$\sin^2\theta = 1 - \cos\theta$$

$$\sin^2\theta = 1 - 0.64$$

$$= 0.36$$

c.
$$tan(\pi + \theta)$$

$$tan(\pi+\theta) = + tan\theta$$

$$m \to 03$$


$$= \frac{\sin \theta}{\cos \theta} = \frac{0.6}{0.8} = \frac{3}{4}$$



Question 10

If $sin(\theta) = 0.6$ where θ is a first quadrant angle, evaluate the following.

a. $sin(\pi + \theta)$




b. $cos(\pi + \theta)$

$$\cos^2\theta = -\sin^2\theta$$

= 0.64
 $\cos\theta = 0.8$ as to in Q1,
=) = $\cos(\pi + \theta) = -0.8$

c. $tan(\pi - \theta)$



Space for Personal Notes

邓立








Let's have a look at this right-angle triangle.





Exploration: Understanding Complementary Relationships

Take a look at this right-angle triangle:



 \blacktriangleright What does $\sin(\theta)$ equal to?

$$\sin(\theta) = \frac{b}{c}$$

On the triangle above, label $\frac{\pi}{2} - \theta$.

What does $\cos\left(\frac{\pi}{2} - \theta\right)$ equal to?

$$\cos\left(\frac{\pi}{2}-\theta\right)=$$

What do you notice?

$$\sin(\theta) = \cos(\frac{\pi}{2} - \theta)$$



Discussion: Given that sin and cos swaps, what would happen to the value of tan?



$$tou\left(\frac{\pi}{2}-\theta\right)=\frac{\sin\left(\frac{\pi}{2}-\theta\right)}{\omega\left(\frac{\pi}{2}-\theta\right)}=\frac{\cos\theta}{\sin\theta}=\frac{1}{\tan\theta}$$



Exploration: Understanding Complementary Relationships

 \blacktriangleright We swap x: cos and y: sin of the unit circle.

$$\sin\left(\frac{\pi}{2}-\theta\right)\to\cos(\theta)$$

$$\cos\left(\frac{\pi}{2} - \theta\right) \to \sin(\theta)$$

$$\tan\left(\frac{\pi}{2}-\theta\right)\to\frac{1}{\tan(\theta)}$$



Question 11

If $sin(\theta) = \frac{1}{3}$ where θ is a first quadrant angles, evaluate the following.

a.
$$\cos\left(\frac{\pi}{2} - \theta\right)$$

b.
$$\sin\left(\frac{\pi}{2} - \theta\right)$$

$$\cos^2\theta = /-\sin^2\theta$$

$$-\omega_1(\theta) = \frac{2\sqrt{2}}{5} \rightarrow \theta \text{ is in } Q_1$$

c.
$$\tan\left(\frac{\pi}{2} - \theta\right)$$

$$\tan\left(\frac{\pi}{2}-\theta\right)=\frac{1}{\tan\theta}$$


$$= \frac{\cos \theta}{\sin \theta} = \frac{\left(\frac{2\sqrt{2}}{3}\right)}{\left(\frac{1}{3}\right)} = 2\sqrt{2}$$





Exploration: Generalising Complementary Relationships

Take a look at the unit circle below.



- Mark the angle $\frac{\pi}{2} \theta$. Which quadrant is it in?
- Mark the angle $\frac{\pi}{2} + \theta$. Which quadrant is it in?
- Mark the angle $\frac{3\pi}{2} \theta$. Which quadrant is it in?
- Mark the angle $\frac{3\pi}{2} + \theta$. Which quadrant is it in? $\Box \Phi$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \Theta - \sin\left(\theta\right)$$

$$\underline{\sin\left(\frac{\pi}{2} - \theta\right)} = \underbrace{+} -\underline{\cos(\theta)}$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \theta - \frac{1}{\tan(\theta)}$$





$$\frac{\pi}{\cos\left(\frac{\pi}{2} + \theta\right)} = +/\sin\left(\theta\right)$$

$$\underline{\sin}\left(\frac{\pi}{2} + \theta\right) = \Theta - \cos(\theta)$$

$$\underline{\tan}\left(\frac{\pi}{2} + \theta\right) = + \underbrace{\cos(\theta)}$$

Third Quadrant



$$\cos\left(\frac{3\pi}{2} - \theta\right) = + \sin\left(\theta\right)$$

$$\frac{\mathbf{y}}{\sin}\left(\frac{3\pi}{2} - \theta\right) = +/\Theta \cos(\theta)$$

$$\tan\left(\frac{3\pi}{2} - \theta\right) = + - \frac{1}{\tan(\theta)}$$



G Fourth Quadrant ______ ← θ



$$\frac{\kappa}{\cos\left(\frac{3\pi}{2} + \theta\right)} = \frac{\kappa}{\cos\left(\frac{3\pi}{2} + \theta\right)} = \frac{\kappa}{\sin\left(\frac{3\pi}{2} + \theta\right)} = \frac{\kappa}{\sin\left(\frac{3\pi}{2}$$

$$\frac{\mathbf{x}}{\cos\left(\frac{3\pi}{2} + \theta\right)} = \mathbf{+} - \sin\left(\theta\right)$$

$$\frac{\mathbf{y}}{\sin\left(\frac{3\pi}{2} + \theta\right)} = \mathbf{+} \mathbf{-} \cos(\theta)$$

$$\tan\left(\frac{3\pi}{2} + \theta\right) = + \left(\frac{1}{\tan(\theta)}\right)$$



Definition

Complementary Relationships

Consider the quadrant for signs.

First Quadrant $\left(\frac{\pi}{2} - \theta\right)$

Step 1: Change function $\frac{\text{Step 2: Chech sign}}{\cos\left(\frac{\pi}{2} - \theta\right) = +\sin(\theta)}$ $\left(\frac{x}{y} + Q?\right)$

$$\sin\left(\frac{\pi}{2} - \theta\right) = +\cos(\theta)$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = +\frac{1}{\tan(\theta)}$$

Second Quadrant $\left(\frac{\pi}{2} + \theta\right)$

$$\sin\left(\frac{\pi}{2} + \theta\right) = +\cos(\theta)$$

$$\cos\left(\frac{\pi}{2} + \theta\right) = -\sin(\theta)$$

$$\tan\left(\frac{\pi}{2} + \theta\right) = -\frac{1}{\tan(\theta)}$$

Third Quadrant $\left(\frac{3\pi}{2} - \theta\right)$

$$\sin\left(\frac{3\pi}{2} - \theta\right) = -\cos(\theta)$$

$$\cos\left(\frac{3\pi}{2} - \theta\right) = -\sin(\theta)$$

$$\tan\left(\frac{3\pi}{2}-\theta\right)=\frac{1}{\tan(\theta)}$$

CONTOUREDUCATION

Fourth Quadrant $\left(\frac{3\pi}{2} + \theta\right)$

$$\sin\left(\frac{3\pi}{2} + \theta\right) = -\cos(\theta)$$

$$\cos\left(\frac{3\pi}{2} + \theta\right) = +\sin(\theta)$$

$$\tan\left(\frac{3\pi}{2} + \theta\right) = -\frac{1}{\tan(\theta)}$$

- Steps
 - 1. Note complementary relationship by identifying a vertical angle $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$.
 - **2.** Equate to the opposite trigonometric function $\cos / \sin / \frac{1}{\tan(\theta)}$.
 - **3.** Determine the sign (\pm) by considering the quadrant.



Question 12 Walkthrough.

If $\sin(\alpha) = \frac{4}{5}$ and $\cos(\beta) = \frac{1}{5}$ where α, β are first quadrant angles, evaluate the following.

a.
$$\cos\left(\frac{\pi}{2} - \alpha\right)$$

$$\cos(\frac{\pi}{3}-a) = \sin(a)$$

$$= \frac{4}{5} \pi$$

b.
$$\sin\left(\frac{\pi}{2} + \alpha\right)$$

$$\frac{\sin(\frac{\pi}{2} + \alpha)}{y, \alpha} = \bigoplus_{\infty} (\alpha)$$

$$= \cos(\alpha)$$

$$= \frac{3}{2}$$

$$=$$
 \pm ω s(a) \pm \sin (B)

=
$$\frac{3 \ln (\beta) - \cos(\alpha)}{5} - \frac{3}{5}$$

= $\frac{2\sqrt{6}-3}{5}$

$$\cos^2(a) = 1 - \sin^2(a)$$

$$= 1 - \frac{16}{25}$$

$$= \frac{9}{25}$$

$$= (05(a)) = \frac{3}{5}$$
(yas a is in G1

$$\sin^2(\beta) = 1 - \cos^2(\beta)$$

$$= 1 - \frac{1}{25}$$

$$= \frac{24}{25}$$
 $\sin(\beta) = \frac{154}{155} - \frac{26}{5}$

Gar Bisin (3



Question 13

If $sin(\alpha) = 0.6$ and $cos(\beta) = 0.4$ where α, β are first quadrant angles, evaluate the following.

a.
$$\cos\left(\frac{\pi}{2} + \alpha\right)$$

a.
$$\cos\left(\frac{1}{2} + a\right)$$

$$=-\sin(a)$$

b.
$$\sin\left(\frac{\pi}{2} - \alpha\right)$$

$$= cos(a)$$

$$= cos(a)$$

$$= 0.8$$

$$\cos^2(\alpha) = 1 - \sin^2(\alpha)$$
= 0.64

$$cos(a) = 0.8$$
 as a is in Q1

g
$$\rightarrow$$
 Q4 \rightarrow Q3
c. $\sin\left(\frac{3\pi}{2} + \alpha\right) - \cos\left(\frac{3\pi}{2} - \beta\right)$

=
$$-\omega s(a) - (-sin(\beta))$$

=
$$sin(\beta)-cos(\alpha)$$

$$Sin^2(\beta) = 1 - cos^2(\beta)$$



<u>Discussion:</u> How can we tell apart from supplementary relationship and complementary relationships?



Supplementary vs Complementary



Complementary: $trig(Vertical\ Angle \pm \theta)$



Question 14

If $sin(\alpha) = 0.6$ and $cos(\beta) = 0.4$ where α, β are first quadrant angles, evaluate the following.

a.
$$\sin(\pi + \alpha)$$

$$= -\sin(a)$$

b.
$$\sin\left(\frac{\pi}{2} + \beta\right)$$

$$= \omega_s(\beta)$$

$$=+\sin(\alpha)-\left(-\cos(\beta)\right)$$



Section C: Solving Trigonometric Equations

Sub-Section: Particular Solutions



Active Recall: Period of Trigonometric Function



Period of sin(nx) and cos(nx) functions = ______

Period of
$$tan(nx)$$
 functions = ______

where n = coefficient of x and n > 0



<u>Discussion</u>: How often would the solution to $\sin(x) = \frac{1}{2}$ repeat?



Particular Solutions



- Solving trigonometric equation for finite solutions.
- Steps
 - Make the trigonometric function the subject.
 - **2.** Find the necessary angle for one period.
 - **3.** Solve for x by equating the necessary angles to the inside of the trigonometric functions.
 - **4.** Add and subtract the period to find all other solutions in the domain.



Question 15 Walkthrough.

Solve the following equations for x over the domain specified.

$$2\sin(2x) + \sqrt{3} = 0$$
 for $x \in [0, 2\pi]$

$$Sin(2n) = \frac{-\sqrt{3}}{2}$$

$$\sin\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}}{2}$$


Step 2: Find Angle For 1 Period

$$2x = \pi + \frac{\pi}{3}, 2\pi - \frac{\pi}{3}$$

$$2x = \frac{4\pi}{3}, \frac{5\pi}{3}$$

Step 3: Remodt Domai

$$P = \frac{2\pi}{J} = \pi$$



Space for Personal Notes



Question 16

Solve the following equations for x over the domains specified.

a.
$$\sin(4x) = -1$$
 for $x \in [-\pi, \pi]$.

$$Sin(\frac{1}{2}) = 1$$

$$Sin\left(\frac{3t}{2}\right) = -1$$

$$4x = \frac{3\pi}{2}$$

Period =
$$\frac{2\pi}{7} = \frac{\pi}{2}$$

$$\therefore Z = \frac{-5\pi}{8}, \frac{-\pi}{8}, \frac{3\pi}{8}, \frac{7\pi}{8}$$

b.
$$2\cos\left(2x - \frac{\pi}{2}\right) + 1 = 0 \text{ for } x \in [0, 2\pi].$$

$$\cos(2x-\frac{\pi}{2})=\frac{-1}{2}$$

$$\omega_s(\frac{\pi}{3}) = \frac{1}{2} \implies \omega_s(\theta) = \bigcirc$$

$$\omega s(\theta) = \bigcirc$$

$$f = \frac{2\pi}{a} = \pi$$



Question 17 Extension.

Solve the following equation for x over the domain specified.

$$\sin^2(2x) - 9\sin(2x) - 5\cos^2(2x) + 8 = 0 \text{ for } x \in (0, \pi)$$

$$\sin^2(2x) - 9\sin(2x) - 5(1-\sin^2(2x)) + 8 = 0$$

$$\sin^2(2x) - 9\sin(2x) - 5 + 5\sin^2(2x) + f = 0$$

$$6\sin^2(2x) - 9\sin(2x) + 3 = 0$$

$$6a^2 - 9a + 3 = 0$$

$$(2a-1)(a-1)=0$$

$$Sin(2x) = \frac{1}{2}$$
 or $sin(2n) = 1$






Question 18 Walkthrough.

Solve the following equations for x over the domains specified.


$$2\tan (2x + \pi) + 2\sqrt{3} = 0 \text{ for } x \in [0, 2\pi]$$



$$\tan(2n+E) = -\sqrt{3}$$

$$\tan(\frac{\pi}{2}) = \sqrt{3} \implies \tan \Theta$$

$$0 = \sqrt{2}, 0.4$$



$$2x = \frac{\pi}{3}$$

$$2 = \frac{2\pi}{6}, \frac{2\pi}{6}, \frac{8\pi}{6}, \frac{11\pi}{6}$$



Discussion: Why do we need to find one angle only for tangents?



() Every soln is a period aport

Question 19

Solve the following equations for x over the domains specified.

$$\sqrt{3}\tan\left(x - \frac{\pi}{3}\right) - 1 = 0 \text{ for } x \in (0, 3\pi)$$

$$\begin{array}{c|c} 2: & \underline{\pi}_{2}, \underline{3\pi}_{2}, \underline{5E}_{2} \end{array}$$



Calculator Commands: Particular Solutions on Technology



Mathematica

Solve
[TrigEquation &&
Domain, x]

TI-Nspire

⊘ Solve (*trigequation*, *x*)|*domain*

Casio Classpad


Solve
 (trigequation, x)|domain

Question 20 Tech-Active.

Solve the following equations for x over the domains specified.

$$\sqrt{3} \tan \left(x - \frac{\pi}{3}\right) - 1 = 0 \text{ for } x \in (0, 3\pi).$$

b.
$$2\cos\left(2x - \frac{\pi}{2}\right) + 1 = 0 \text{ for } x \in [0, 2\pi].$$



Space for Personal Notes





Contour Check

□ <u>Learning Objective</u>: [4.1.1] – Evaluate exact values for sine, cosine, and tangent

Key Takeaways

Radians and Degrees

$$\mathbf{1}^c = \left(\frac{180}{\pi}\right)^{\mathbf{0}}$$

$$\mathbf{1}^{o} = \left(\frac{\pi}{180}\right)^{c}$$

$$180^{\circ} = \pi^{c}$$

■ The Exact Values Table:

x	0 (0°)	$\frac{\pi}{6} (30^{\circ})$	$\frac{\pi}{4} (45^{\circ})$	$\frac{\pi}{3} \left(60^{\circ}\right)$	$\frac{\pi}{2} (90^{\circ})$
$\sin(x)$					
$\cos(x)$					
tan(x)					



Learning Objective: [4.1.2] - Applying Pythagorean identity to evaluate
trigonometric functions

Key Takeaways

Pythagorean Identity:

$$\sin^2(\theta) + \cos^2(\theta) = \underline{\hspace{1cm}}$$

□ <u>Learning Objective</u>: [4.1.3] – Apply supplementary and complementary relationship to evaluate trigonometric functions

Key Takeaways

- Supplementary Relationships:
- ☐ Look at the quadrant to find the correct sign.
 - Second Quadrant $(\pi \theta)$
 - \square Reflection around y-axis.

$$\cos(\pi - \theta) =$$

$$\sin(\pi - \theta) = \underline{\hspace{1cm}}$$

$$\tan(\pi - \theta) = \underline{\hspace{1cm}}$$



- O Third Quadrant $(\pi + \theta)$
 - \square Reflection around x- and y-axis.

$$\cos(\pi + \theta) =$$

$$\sin(\pi + \theta) =$$

$$\tan(\pi + \theta) = \underline{\hspace{1cm}}$$

- Fourth Quadrant $(-\theta)$
 - \square Reflection around x-axis.

$$\cos(-\theta) =$$

$$sin(-\theta) = \underline{\hspace{1cm}}$$

$$tan(-\theta) = \underline{\hspace{1cm}}$$

- Steps
 - **1.** Equate to $\cos / \sin / \tan(\theta)$.
 - **2.** Determine the sign (\pm) by considering the quadrant.



- Complementary Relationships:
 - First Quadrant $\left(\frac{\pi}{2} \theta\right)$

$$\cos\left(\frac{\pi}{2}-\theta\right)=$$

$$\sin\left(\frac{\pi}{2}-\theta\right)=$$

$$\tan\left(\frac{\pi}{2}-\theta\right) = \underline{\hspace{1cm}}$$

O Second Quadrant $\left(\frac{\pi}{2} + \theta\right)$

$$\sin\left(\frac{\pi}{2}+\theta\right)=$$

$$\cos\left(\frac{\pi}{2}+\theta\right)=$$

$$\tan\left(\frac{\pi}{2}+\theta\right) = \underline{\hspace{1cm}}$$

O Third Quadrant $\left(\frac{3\pi}{2} - \theta\right)$

$$\sin\left(\frac{3\pi}{2}-\theta\right)=\underline{\hspace{1cm}}$$

$$\cos\left(\frac{3\pi}{2}-\theta\right)=$$

$$\tan\left(\frac{3\pi}{2}-\theta\right)=\underline{\hspace{1cm}}$$



O Fourth Quadrant $\left(\frac{3\pi}{2} + \theta\right)$

$$\sin\left(\frac{3\pi}{2}+\theta\right)=\underline{\hspace{1cm}}$$

$$\cos\left(\frac{3\pi}{2}+\theta\right)=$$

$$\tan\left(\frac{3\pi}{2}+\theta\right)=\underline{\hspace{1cm}}$$

O Steps:

- 1. Note complementary relationship by identifying a vertical angle $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$.
- **2.** Equate to the opposite trigonometric function $\cos / \sin / \frac{1}{\tan(\theta)}$.
- **3.** Determine the sign (\pm) by considering the quadrant.



Learning Objective: [4.1.4] - Solve particular solution for trigonometric functions						
Key Takeaways						
□ Period						
Period of $sin(nx)$ and $cos(nx)$ functions =						
Period of $tan(nx)$ functions =						
where $oldsymbol{n}=$ coefficient of $oldsymbol{x}$ and $oldsymbol{n}>0$						
□ Particular Solutions:						
 Solving trigonometric equations for finite solutions. 						
O Steps:						
1. Make the trigonometric function the						
2. Find the necessary for one period.						
3. Solve for x by equating the necessary angles to the of the trigonometric functions.						
4. Add and subtract the to find all other solutions in the domain.						



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

VCE Mathematical Methods ½

Free 1-on-1 Support

Be Sure to Make the Most of These (Free) Services!

- Experienced Contour tutors (45 + raw scores, 99 + ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

1-on-1 Video Consults	Text-Based Support
 Book via bit.ly/contour-methods-consult-2025 (or QR code below). One active booking at a time (must attend before booking the next). 	 Message <u>+61 440 138 726</u> with questions. Save the contact as "Contour Methods".

Booking Link for Consults
bit.ly/contour-methods-consult-2025



Number for Text-Based Support +61 440 138 726

