



Website: [contoureducation.com.au](http://contoureducation.com.au) | Phone: 1800 888 300

Email: [hello@contoureducation.com.au](mailto:hello@contoureducation.com.au)

## VCE Mathematical Methods $\frac{3}{4}$ Coordinate Geometry Exam Skills [1.6] Workbook

### Outline:



|   |          |                                       |          |
|---|----------|---------------------------------------|----------|
| <b><u>Recap</u></b>                                 | Pg 02-15 |                                       |          |
| <b><u>Warm-Up Test</u></b>                          | Pg 16-19 | <b><u>Exam 1 Questions</u></b>        | Pg 26-30 |
| <b><u>Coordinate Geometry Exam Skills</u></b>       | Pg 20-25 | <b><u>Tech-Active Exam Skills</u></b> | Pg 31-33 |
| ➤ Reflect a Point Around a Vertical/Horizontal Line |          | <b><u>Exam 2 Questions</u></b>        | Pg 34-39 |
| ➤ Reflect a Point Around a Line                     |          |                                       |          |
| ➤ Application of Angle Between Two Lines            |          |                                       |          |

### Learning Objectives:


- ❑ MM34 [1.6.1] - Apply midpoint to find a reflected point.
- ❑ MM34 [1.6.2] - Find the angle between a line and  $x$ -axis or two lines.



## Section A: Recap

*All the students who were here last week, skip to section B: Warm-Up Test!*

### Midpoint



- The midpoint,  $M$ , of two points  $A$  and  $B$  is simply the point halfway between  $A$  and  $B$ .

$$M(x_m, y_m) = \left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

- The midpoint can be found by taking the \_\_\_\_\_ of the  $x$ -coordinate and  $y$ -coordinate of the two points.

### Distance Between Two Points


- The distance between two points  $(x_1, y_1)$  and  $(x_2, y_2)$  can be found using Pythagoras' theorem:

$$\text{Distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

### Question 1

Find the points on the line  $y = 2x - 6$  which has a distance of  $\sqrt{5}$  from the point  $(2, 1)$ .

#### Horizontal Distance




Horizontal Distance = \_\_\_\_\_ where  $x_2 > x_1$

➤ Find the difference between their  $x$ -values.



### Vertical Distance




Vertical Distance= \_\_\_\_\_ where  $y_2 > y_1$

- Find the difference between their  $y$ -values.



### Parallel Lines



- Parallel lines have the same gradient.


$$m_1 = m_2$$

Space for Personal Notes

### Question 2

Find a line that is parallel to  $y = 3x - 1$  passing through the point  $(-2, 6)$ .


### Perpendicular Lines



### Question 3

Find a line that is perpendicular to  $y = 3x - 1$  passing through the point  $(1, 0)$ .

### Angle Between a Line and the $x$ -axis




- The angle between a line and the \_\_\_\_\_ direction of the  $x$ -axis (anticlockwise) is given by:

$$\tan(\theta) = m$$

#### Question 4

Find the angle made between the line  $y = -x + 2$  and the  $x$ -axis measured in the anticlockwise direction.

#### Acute Angle Between Two Lines



$$\theta = |\tan^{-1}(m_1) - \tan^{-1}(m_2)|$$

➤ Alternatively:

$$\tan(\theta) = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

For your understanding, note that this formula is derived from the  $\tan$  compound angle formula covered in SM34.

**NOTE:**  $|x|$  just takes the positive value of  $x$ .


**Question 5 Tech-Active.**

Find the acute angle between the lines  $x - 3y = 2$  and  $y = \frac{4}{5}x - 2$ . Give your answer in degrees correct to two decimal places.

**Exploration: Geometry of the Number of Solutions Between Linear Graphs**

➤ Unique Solution

$$m_1 \neq m_2$$




They just need to have \_\_\_\_\_.



➤ Infinite Solutions

$$m_1 = m_2 \text{ AND } c_1 = c_2$$




They just need to have the same \_\_\_\_\_ and the same \_\_\_\_\_.

In other words, they have to be the \_\_\_\_\_.

➤ No Solutions

$$m_1 = m_2 \text{ AND } c_1 \neq c_2$$






They need to have the \_\_\_\_\_ but \_\_\_\_\_ + c.

They have to be two different \_\_\_\_\_ lines.



### General Solutions of Simultaneous Linear Equations

➤ Two linear equations are either:

-  The same line is expressed in a different form. In this case, they have infinitely many solutions.
-  Unique lines that are parallel. In this case, they have no solutions.
-  Unique lines which are not parallel. In this case, they have exactly one solution.

Space for Personal Notes

Consider the following pair of simultaneous equations in terms of  $a \in \mathbb{R} \setminus \{0\}$ :

$$ax + 3y = 1$$

$$2x + (a + 1)y = 1$$

- a.** Find the value of  $a$  for which there are no solutions to the simultaneous equations.

- c. Find the value of  $a$  for which there are infinite solutions to the simultaneous equations.

### Solving Systems of Linear Equations with Parameters

- Occurs when solving for three variables with two equations. We simply,

$$\text{Let } x = k, \text{ or}$$

$$\text{Let } y = k, \text{ or}$$

$$\text{Let } z = k$$

- And solve simultaneously.

Space for Personal Notes

**Question 7**

Solve the following system of linear equations with the parameter of  $k$ .

$$x + 3z = 1$$

$$x + y = 2$$

Space for Personal Notes




## Addition of Ordinates

### ➤ Definition:

🔄 Technique used to graph the sum/difference of two functions.

$$e.g. y = \sin(x) + \sqrt{x}$$



➤ The addition of ordinates involves adding the \_\_\_\_\_ of two functions.

## Add two y-values

### ➤ Steps to sketching $f(x) + g(x)$ :

1. Sketch  $f(x)$  and  $g(x)$  on the same axes.
2. Plot points for  $f(x) + g(x)$  by adding the **y-values** of  $f(x)$  and  $g(x)$ .

➤ At  $x$ -intercepts, the sum equals to the \_\_\_\_\_. **Why?**


➤ At intersections, the sum equals to \_\_\_\_\_ the  $y$ -value. **Why?**

➤ When functions are equidistant from  $x$ -axis, the sum equals to \_\_\_\_\_. **Why?**

3. Join the plotted points.

**Question 8**

Plot the sum of the two functions given below, using the addition of ordinates.



Space for Personal Notes

## Section B: Warm-Up Test (15 Marks)

INSTRUCTION: 15 Marks. 15 Minutes Writing.



### Question 9 (3 marks)

Given that the distance between point  $A(3, 4)$  and point  $B(m, 2)$  is 3 units, find the possible values of  $m$ .

---

---

---

---

---

---

---

---

---

Space for Personal Notes



**Question 10** (3 marks)

Find the equation of the line that passes through  $(2, 1)$  and is perpendicular to a line that makes an angle of  $60^\circ$  with the positive direction of the  $x$ -axis.

---

---

---

---

---

---

---

---

---

---

---

Space for Personal Notes

**Question 11** (4 marks)

Sarah is standing at point  $Q(7, 3)$  and wants to walk to the road, which is described by  $y = 2x - 5$ . But Sarah wants to reach the road by covering the least amount of distance possible.

- a. Find the equation of the line that is perpendicular to  $y = 2x - 5$  and passes through the point  $Q(7, 3)$ . (2 marks)

---

---

---

---

---

---

- b. Hence, find the shortest distance that Sarah can travel to reach the road. (2 marks)

---

---

---

---

---

---

**Space for Personal Notes**

**Question 12** (5 marks)

Consider the simultaneous linear equations:

$$kx + 4y = 6$$

$$2x + (k - 2)y = 3$$

Where  $k$  is a real constant.

- a. Find the values of  $k$  for which there is a unique solution to the simultaneous equations. (2 marks)

---

---

---

---

---

---

- b. Find the values of  $k$  for which there are infinitely many solutions. (2 marks)

---

---

---

---

---

---

- c. Find the values of  $k$  for which there are no solutions. (1 mark)

---


---

## Section C: Coordinate Geometry Exam Skills

### Sub-Section: Reflect a Point Around a Vertical/Horizontal Line


#### Exploration: Reflection of a Point Around a Vertical/Horizontal Line

- Consider a point reflected around  $y = 3$ .



- What do you notice about their midpoint?
- What equation can we construct?

#### Reflection of a Point Around a Vertical/Horizontal Line




- Midpoint must be on the line of reflection.

**Question 13**

Find the reflection of  $(3, 1)$  around  $x = 1$ .

Space for Personal Notes

## Sub-Section: Reflect a Point Around a Line




*How about non-vertical/horizontal lines?*



### Exploration: Reflection of a Point in a Line



- Consider point  $A$  which is reflected across the line  $y = mx + c$  to point  $B$ , the line  $AB$  is \_\_\_\_\_ to the mirror.




- The distance between  $A$  and the line is \_\_\_\_\_ to the distance between the line and point  $B$ .
- Where would the midpoint of  $A$  and  $B$  lie?

Space for Personal Notes



Finding the Reflection of a Point in a Line:



► Steps:

1. Find the perpendicular line passing through the point.
2. Find the intersection between the original line and the perpendicular line.
3. Find the reflected point  $(x, y)$  by treating the intersection from 2. as the midpoint between the original and reflected point.

**Question 14 Walkthrough.**

Find the reflection of  $(1, 2)$  in the line  $y = x$ .

**Active Recall: Steps for Finding the Reflection of a Point in a Line**



1. Find the \_\_\_\_\_ line passing through the point.
2. Find the \_\_\_\_\_ between the original line and the perpendicular line.
3. Find the reflected point  $(x, y)$  by treating the intersection from 2. as the \_\_\_\_\_ between the original and reflected point.

**Question 15**

Find the reflection of  $(3, 1)$  in the line  $y = x + 2$ .

**Question 16 Extension.**

Find the equation of the line in the form  $y = mx + c$  that the point  $(1, 2)$  is reflected in the point  $(4, 6)$ .

Space for Personal Notes





## Sub-Section: Application of Angle Between Two Lines

### Question 17 Walkthrough.

It is known that the angle between  $y = x - 1$  and  $y = mx + 1$  is given by  $60^\circ$ . Find the value(s) of  $m$ .

### Question 18

It is known that the angle between  $y = 4x - 3$  and  $y = mx + 5$  is given by  $45^\circ$ . Find the value(s) of  $m$ .

**Section D: Exam 1 Questions (16 Marks)****Question 19** (5 marks)

Consider the simultaneous linear equations:

$$(k + 1)x + 3y = 6$$

$$4x + (k - 3)y = 4$$

Where  $k$  is a real constant.

- a.** Find the values of  $k$  for which there is a unique solution to the simultaneous equations. (2 marks)

---

---

---

---

---

---

---

---

- b.** Find the value of  $k$  for which there are infinitely many solutions. (2 marks)

---

---

---

---

c. Find the value of  $k$  for which there are no solutions. (1 mark)

---


---

---

Space for Personal Notes

**Question 20** (3 marks)

Consider the diagram below:



The rectangle  $OPQR$  has a vertex  $Q(4, 3)$  on the line that passes through  $U$  and  $V$ .

- a. Find an expression for  $v$  in terms of  $u$ . (1 mark)

---



---



---



---

- b. Hence, find an expression for the shaded area in terms of  $u$ . (2 marks)

---



---



---



---



---

**Question 21** (6 marks)

The point  $P(5, 2)$  is reflected in the line  $y = 2x - 3$  to become the point  $P'$ .

- a. Find the coordinates of  $P'$ . (3 marks)

---

---

---

---

---

---

---

---

- b. Find the minimum distance between the point  $P$  and the line  $y = x - 1$ . (3 marks)

---

---

---

---

---

---


---

---

**Space for Personal Notes**

**Question 22** (2 marks)

The graphs of  $f$  and  $g$  are sketched on the axes below. Sketch the graph of  $f + g$  on the same axes.



Space for Personal Notes

## Section E: Tech-Active Exam Skills



### Calculator Commands: Simultaneous Equations on CAS

#### System of Linear Equations

##### Example:

The simultaneous linear equations  $ax - 3y = 5$  and  $3x - ay = 8 - a$  have no solution for:

- A.  $a = 3$
- B.  $a = -3$
- C. Both  $a = 3$  and  $a = -3$ .
- D.  $a \in \mathbb{R} \setminus \{3\}$
- E.  $a \in \mathbb{R} \setminus [-3, 3]$

```
system_solve(a*x-3*y=5,3*x-a*y=8-a,a)
```

- ▶ Solving:  $\begin{bmatrix} a \cdot x - 3 \cdot y = 5 \\ 3 \cdot x - a \cdot y = 8 - a \end{bmatrix}$
- ▶ Unique Solution:  $a \neq -3$  and  $a \neq 3$
- ▶ No Solutions:  $a = -3$
- ▶ Infinite Solutions:  $a = 3$

#### Overview:

This program takes two linear equations and a parameter and finds the parameter values for the system to obtain a unique solution, no solution, or infinite solutions.

#### Input:

*system\_solve*(*< equation 1 >*,  
*< equation 2 >*,  
*< parameter >*)

#### Other Notes:

The program can only handle one parameter.

➤ Or menu – 3 – 7.

➤ UDF line functions:

### Normal Line

```
normal_line(x^3-x,x,2)

  ▶ Derivative: 3·x2-1
  ▶ Gradient: 11
  ▶ Perpendicular Gradient:  $-\frac{1}{11}$ 
  ▶ Passes Through: [2 6]
  ▶ x-Intercept: [68 0]
  ▶ Vertical Intercept:  $\left[0 \frac{68}{11}\right]$ 
  ▶ Normal Line:
    
$$\frac{68}{11} - \frac{x}{11}$$

```

#### Overview:

This program will find all the necessary information related to a normal line at a point on a function, which includes:

- The derivative.
- The gradient and perpendicular gradient.
- The point on the function the normal line passes through.
- The axis intercepts of the normal line.
- The equation of the normal line.

#### Input:

***normal\_line(< function >, < variable >, < x point >)***

### Tangent Line

```
tangent_line(x^3-x,x,2)

  ▶ Derivative: 3·x2-1
  ▶ Gradient: 11
  ▶ Passes Through: [2 6]
  ▶ x-Intercept:  $\left[\frac{16}{11} 0\right]$ 
  ▶ Vertical Intercept: [0 -16]
  ▶ Tangent Line:
    
$$11 \cdot x - 16$$

```

#### Overview:

This program will find all the necessary information related to a tangent line at a point on a function, which includes:

- The derivative.
- The gradient of the tangent line.
- The point on the function the tangent line passes through.
- The axis intercepts of the tangent line.
- The equation of the tangent line.

#### Input:

***tangent\_line(< function >, < variable >, < x point >)***





## Calculator Commands: Finding the Angle Between Two Lines

- The angle between two lines with gradients  $m_1$  and  $m_2$  respectively is

$$\theta = |\tan^{-1}(m_1) - \tan^{-1}(m_2)|$$

### ➤ Mathematica

- Use the Abs[] function.

```
In[126]:= Abs[ArcTan[2] - ArcTan[1]] / Degree // N
Out[126]:= 18.4349
```

### ➤ TI-Nspire

- Find the modulus sign.



Calculator interface showing the calculation of the angle between two lines with gradients 2 and 1. The result is 18.4349.

### ➤ Casio Classpad

- Modulus sign under Math1.

Calculator interface showing the calculation of the angle between two lines with gradients 2 and 1. The result is 18.4349882.

## Calculator Commands: Finding the Gradients of Lines Given the Angle They Make

- If we know the angle and one of the gradients  $m_1$  or  $m_2$  then we can find the other gradient by Solving,

$$\tan(\theta) = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

- e.g. Find the gradient of the line that makes an angle of  $60^\circ$  with  $y = -x$ .

### ➤ Mathematica

### ➤ TI-Nspire

- Find the modulus sign.



### ➤ Casio Classpad

- Modulus sign under Math1.



**Section F: Exam 2 Questions (13 Marks)****Question 23** (1 mark)

The perpendicular bisector of the points  $(3, 6)$  and  $(7, -4)$  is:

- A.  $y = -x + 2$
- B.  $y = \frac{2}{5}x - 1$
- C.  $y = -\frac{5}{2}x + 1$
- D.  $y = x + \frac{2}{5}$

**Question 24** (1 mark)

It is known that the lines  $y = mx + 4$  and  $y = 3x - 5$  make an angle of  $45^\circ$  when they intersect. The possible values of  $m$  are:

- A.  $m = -\frac{1}{2}$  only
- B.  $m = 2$  only
- C.  $m = -2, \frac{1}{2}$
- D.  $m = -2, -\frac{1}{2}$

Space for Personal Notes

**Question 25** (1 mark)

The simultaneous linear equations:

$$2x - (k + 3)y = 8$$

$$(2 - k)x - 2y = 3$$

Where  $k$  is a real constant that has no solutions for:

- A.  $k = 1$  only
- B.  $k = -2$  only
- C.  $k = -2, 1$
- D.  $k \in \mathbb{R} \setminus \{-2, 1\}$

**Question 26** (1 mark)

The acute angle made between the lines  $y = 2x - 3$  and  $y = -x + 1$  correct to the nearest degree is:

- A. 108
- B. 70
- C. 72
- D. 51

Space for Personal Notes

**Question 27** (1 mark)

The simultaneous linear equations:

$$(k + 1)x - 3y = 1, 0 \leq x \leq 1$$

$$4x - (k + 5)y = 2, 0 \leq x \leq 1$$


Where  $k$  is a real constant always has exactly one solution for:

- A.  $k > 1$
- B.  $k > -6$
- C.  $k \in [-6, \infty) \setminus \{1\}$
- D.  $k \geq -7$

Space for Personal Notes

**Question 28** (8 marks)

Emma is riding a bicycle on a hill, which starts from the origin as shown in the diagram. When she reaches the point  $(10, 6)$ , she fires a laser at an angle of  $\theta = 60^\circ$  above the horizontal. Meanwhile, David fires a laser from the origin that is parallel to Emma's laser such that both lasers travel together through space.



- a. Find the shortest distance between the two laser paths. Give your answer correct to two decimal places. (3 marks)

---

---

---

---

---

---


---

---

---

---

Emma now changes the direction in which she fires her laser.



- b. At what angle,  $\phi$ , should Emma have fired her laser such that the two lasers would intersect at  $x = 8$ ? Give your answer correct to two decimal places. (3 marks)

---

---

---

---

---

---

---

---

---

---

- c. Calculate the acute angle between these two laser paths when they intersect. Express your answer in degrees correct to two decimal places. (1 mark)

---

---

---

---

- d. Calculate the vertical distance between the two lasers when  $x = 10$ . Give your answer correct to two decimal places. (1 mark)

---

---

---

---

Space for Personal Notes



## Contour Check

### Learning Objective: [1.6.1] - Apply Midpoint to Find a Reflected Point.

#### Key Takeaways

- ☐ The line between a point and its reflection is \_\_\_\_\_ to the line it is reflected in.
- ☐ The \_\_\_\_\_ of a line and its reflection lies on the line it is reflected in.
- ☐ **Steps** for Finding the Reflection of a Point in a Line
  - ☐ Find the \_\_\_\_\_ line passing through the point.
  - ☐ Find the \_\_\_\_\_ between the original line and the perpendicular line.
  - ☐ Find the reflected point  $(x, y)$  by treating the intersection from 2. as the \_\_\_\_\_ between the original and reflected point.

### Learning Objective: [1.6.2] - Find the Angle Between a Line and $x$ -axis or Two Lines.

#### Key Takeaways

- ☐ To find the angle between a line and the  $x$ -axis, we can use equation  $m =$  \_\_\_\_\_.
- ☐ To find the angle between two lines, we can use  $\theta =$  \_\_\_\_\_ or  $\tan(\theta) =$  \_\_\_\_\_.



## VCE Mathematical Methods $\frac{3}{4}$ Free 1-on-1 Support



### Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

| <u>1-on-1 Video Consults</u>   | <u>Text-Based Support</u>   |
|--|---|
| <ul style="list-style-type: none"><li>➤ Book via <a href="https://bit.ly/contour-methods-consult-2025">bit.ly/contour-methods-consult-2025</a> (or QR code below).</li><li>➤ One active booking at a time (must attend before booking the next).</li></ul> | <ul style="list-style-type: none"><li>➤ Message <a href="tel:+61440138726">+61 440 138 726</a> with questions.</li><li>➤ Save the contact as "Contour Methods".</li></ul> |

[Booking Link for Consults](https://bit.ly/contour-methods-consult-2025)  
[bit.ly/contour-methods-consult-2025](https://bit.ly/contour-methods-consult-2025)



[Number for Text-Based Support](tel:+61440138726)  
[+61 440 138 726](tel:+61440138726)