



Website: contoureducation.com.au | Phone: 1800 888 300
Email: hello@contoureducation.com.au

VCE Mathematical Methods $\frac{3}{4}$
Polynomials [1.7]
Test

20 Marks. 1 Minute Reading. 22 Minutes Writing

Results:

Test Questions	_____ / 17
Extension Test Question	_____ / 3



Section A: Test Questions (17 Marks)

Question 1 (5 marks)

Tick whether the following statements are true or false.

	True	False
a. A root which corresponds to a polynomial with a factor of $x + 4$ is 4. (0.5 marks)		
b. If $f(2) = 4$, then $f(x) \div (2x - 4)$ has a remainder of 4. (0.5 marks)		
c. The remainder of $(x^3 + x^2 - x + 1) \div (x + 1)$ is 2. (0.5 marks)		
d. To factorise a quartic, we must first find two roots by trial and error. (0.5 marks)		
e. The rational root theorem suggests that $ax^3 + bx^2 + cx + d$ will have roots that are factors of d divided by the factors of a . (0.5 marks)		
f. All odd functions must have odd powers for power functions. (0.5 marks)		
g. Even functions are always symmetrical around the x -axis. (0.5 marks)		
h. The maximal domain of $f(x) = x^{\frac{3}{2}}$ is given by $[0, \infty)$. (0.5 marks)		
i. $f(x) = x^{\frac{2}{3}}$ is an even function. (0.5 marks)		
j. If $f(-x) = -f(x)$, then it is an even function. (0.5 marks)		

Space for Personal Notes

Question 2 (3 marks)

Consider the function $f(x) = x^3 + ax^2 + bx$. If $x - 2$ is a factor of $f(x)$ and the remainder of $f(x) \div (x - 3)$ is given by 12, find the value(s) of a and b .

Space for Personal Notes

Question 3 (3 marks)

Solve the following equation for x .


$$2x^3 - 5x^2 - 28x + 13 = -2$$

Space for Personal Notes

Question 4 (3 marks)

Sketch the graph of the following function on the axes below. Label all axes intercepts with their coordinates.

$$y = (x - 2)^2(x - 3)(x + 1)$$



Space for Personal Notes

Question 5 (3 marks)

It is known that $f(x)$ is an even function where $f(-2) = 4$ and $f'(-2) = 3$.

Let $g(x) = f(x) + 2$.

Find the values of $g(2)$ and $g'(2)$.

Space for Personal Notes

Section B: Extension Test Question (3 Marks)**Question 6** (3 marks)

Consider $f(x) = 2x^3 - 3x^2 - 12x + 2$.

Given that the turning point occurs at $x = -1$ and $x = 2$, find the value(s) of k such that $f(x) + k = 0$ has 3 solutions.

Space for Personal Notes



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

VCE Mathematical Methods $\frac{3}{4}$

Free 1-on-1 Support



Be Sure to Make The Most of These (Free) Services!

- Experienced Contour tutors (45+ raw scores, 99+ ATARs).
- For fully enrolled Contour students with up-to-date fees.
- After school weekdays and all-day weekends.

<u>1-on-1 Video Consults</u>	<u>Text-Based Support</u>
➤ Book via bit.ly/contour-methods-consult-2025 (or QR code below). ➤ One active booking at a time (must attend before booking the next).	➤ Message +61 440 138 726 with questions. ➤ Save the contact as "Contour Methods".

[Booking Link for Consults](https://bit.ly/contour-methods-consult-2025)
bit.ly/contour-methods-consult-2025



[Number for Text-Based Support](tel:+61440138726)
[+61 440 138 726](tel:+61440138726)