



Website: contoureducation.com.au | Phone: 1800 888 300
Email: hello@contoureducation.com.au

VCE Specialist Mathematics $\frac{1}{2}$ Sequences and Series [1.3] Workbook

Outline:



Introduction to Sequences and Series

Pg 2-8

- Sequences
- Recurrence Relations
- Introduction to Series

Arithmetic Sequence and Series

Pg 9-18

- Introduction to Arithmetic Sequence
- Arithmetic Recurrence Relation
- Arithmetic Mean
- Arithmetic Series

Geometric Sequence and Series

Pg 19-29

- Geometric Sequence
- Geometric Recurrence Relation
- Geometric Mean
- Geometric Series
- Infinite Geometric Series

Section A: Introduction to Sequences and Series

Sub-Section: Sequences



Sequences

- Consider Samuel, our beloved Contour Tutor.



- Lets say he eats a chocolate bar on the first day.
- The next day, he eats two chocolate bars.
- How many chocolate bars do you think he eats on the third day?
- Hands up if you think it's 3 chocolate bars!
 - 🇨 You guys have taken an **arithmetic** approach!
- Hands up if you think it's 4 chocolate bars!
 - 🇨 You guys have taken a **geometric** approach!
 - 🇨 **We will consider the two types of sequences today!**



Sequences

$$t_n = f(n)$$

- **Definition:** A sequence is an ordered list of numbers following a certain **pattern**.
- It is a _____ of the order _____.

Question 1 Walkthrough.

Construct the first 3 terms for the sequence given by $t_n = 2n + 1$.

NOTE: t_n stands for the n^{th} term. Eg: For t_3 , our value of n is equal to 3.

ALSO NOTE: This was a sequence of odd numbers!



Question 2

The sequence is defined by $t_n = 2^n + 1$. Identify the term number for which t_n equals 9.

Space for Personal Notes

Sub-Section: Recurrence Relations



What if we define the term t_n with respect to the previous term (t_{n-1})?



Recurrence Relations



➤ Definition:

- A recurrence relation is when we define a term (t_n), in terms of the previous one (t_{n-1}).
- Recurrence relations generate sequences of the form:

$$t_n = f(t_{n-1}) \text{ where } t_1 = a$$

Or


$$t_{n+1} = f(t_n) \text{ where } t_1 = a$$

- It must always include a _____.

Analogy: Recurrence Relations



- It's like knocking off a sequence of dominos!



- We focus on the _____ between the two terms (dominos).

Question 3


Consider the following recurrence relation.

$$t_n = 3t_{n-1} + 2, \quad t_1 = 2$$

State the value of t_3 .


Why do we always need a first term?

Analogy: Reason for why recurrence relations always need a first term.



- Can you knock off any of these dominos without knocking over the first one?
- Similarly, how can we solve for any term in recurrence relation without the first term?
It's impossible!

Sub-Section: Introduction to Series




What does the word "series" mean?



Series



► Definition:

 A series is the sum of the first n terms of a sequence.

$$S_n = \sum_{i=1}^n t_i$$

Question 4 Walkthrough.

Consider the sequence given by $t_n = 3n - 4$.

Evaluate S_2 .

Question 5

Consider the sequence given by $t_n = 2n + 2$.

Evaluate S_4 .

Key Takeaways

- ✓ Sequence follows a certain pattern.
- ✓ Recurrence relation is a relationship between the next term and the current term.
- ✓ Series is the sum of the first n terms.



Section B: Arithmetic Sequence and Series

Sub-Section: Introduction to Arithmetic Sequence



Let's go back to the previous context



Arithmetic Sequence



- Consider Samuel, our beloved Contour Tutor.



- He ate one chocolate bar on the first day.
- The next day, he ate two chocolate bars.
- Arithmetic sequence suggests Sam eats 3 chocolate bars on the third day !

Discussion: In an arithmetic sequence, how do we go from one term to the next?





Arithmetic Sequences

$$a, \quad a + d, \quad a + 2d, \quad a + 3d$$

► Definition

-  An **arithmetic sequence** is one where the **common difference** is added or subtracted to get the next term.

$$t_n = a + (n - 1)d$$

-  Where d is the common difference, and a is the first term.

Discussion: Why do we add $(n - 1)d$ instead of nd ? How many differences do we add from t_1 to t_n ?



Question 6 Walkthrough.

Consider the arithmetic sequence defined by $t_n = 12 + 6(n - 1)$.

Identify the common difference, first term and the 5th term.

Question 7

Consider the arithmetic sequence defined by $t_n = -1 + 6n$.

Identify the common difference, first term and the 8th term.

NOTE: Read the question carefully. Sometimes, they expand the $n - 1$ factor to confuse you.



Space for Personal Notes

Sub-Section: Arithmetic Recurrence Relation



What about recurrence relations for arithmetic sequence?



Discussion: What must be the relationship between the current term (t_n) and the previous term (t_{n-1}) for an arithmetic sequence?



Formula: Recurrence Relation for Arithmetic Sequence



$$t_n = t_{n-1} + d \text{ where } t_1 = a$$


Question 8

Consider the following n^{th} term rule for the arithmetic sequence:

$$t_n = 10 - 3(n - 1)$$

Find the recurrence relation which corresponds to it.

Sub-Section: Arithmetic Mean



What do we mean by the Arithmetic Mean?



Exploration: Finding arithmetic mean.



- Consider three terms of an arithmetic sequence with the common difference of d .

$$t_1 = a, t_2 = a + d \text{ and } t_3 = a + 2d.$$

- Say that t_2 is an arithmetic mean (average) of t_1 and t_3 .
- Try finding the sum of t_1 and t_3 .

$$t_1 + t_3 = \underline{\hspace{2cm}}$$

- What should we do to $t_1 + t_3$ to find t_2 ?

$$t_2 = \frac{t_1 + t_3}{2}$$

The Arithmetic Mean



- **Definition:**


$$\text{Arithmetic Mean of } a \text{ and } b = \frac{a + b}{2}$$

NOTE: Arithmetic mean is same as the so called "normal average".




Space for Personal Notes

Sub-Section: Arithmetic Series



Discussion: What would be the most efficient way of adding all the whole numbers 1 – 100?



$$1 + 2 + 3 + \dots + 98 + 99 + 100$$

Question 9

Find the sum of all the odd numbers from 1 to 99.

$$1 + 3 + 5 + \dots + 95 + 97 + 99.$$

Now let's generalise it for all arithmetic sequences!



Arithmetic Series (Form 1)

- Use the following formula, if we know the first term, last term and number of terms.

$$S_n = \frac{n}{2}(a + l)$$

► Where n = number of terms, a = first term and l = last term.

► $\frac{n}{2}$ can be thought as the _____.

► $a + l$ can be thought as the _____.

Question 10 Walkthrough.

Consider the arithmetic sequence with $t_1 = -5$ and $t_4 = 10$.

Find S_4 .

Question 11

Consider the arithmetic sequence with $t_1 = 3$ and $t_9 = 19$.

Find S_9 .

Now let's generalise it for all arithmetic sequences!

Exploration: Arithmetic series for when we don't know the last term (l).

➤ Recall the series formula-

$$S_n = \frac{n}{2}(a + l)$$

➤ Which term would l be?

➤ How can we define t_n for an arithmetic sequence?

➤ On the space below, substitute $l = a + d(n - 1)$ to the series formula!



Arithmetic Series (Form 2)

► Use the following formula, if we know the first term, common difference and number of terms.

$$S_n = \frac{n}{2} (2a + d(n - 1))$$

◀ Where n = number of terms, a = first term and d = common difference.

Question 12 Walkthrough.

Consider the arithmetic sequence with $t_n = 2 + 3(n - 1)$.

Find S_{10} .

Space for Personal Notes

Question 13

Consider the arithmetic sequence with $t_n = -2 + \frac{1}{3}(n - 1)$.

Find S_{13} .


Key Takeaways



- ✓ Arithmetic sequence has a common difference between the next term and the current one.
- ✓ Arithmetic sequence is given by $t_n = a + (n - 1)d$.
- ✓ Arithmetic mean of a and b is $\frac{a+b}{2}$.
- ✓ Arithmetic sum is given by $\frac{n}{2}(a + l)$ or $\frac{n}{2}(2a + (n - 1)d)$.

Section C: Geometric Sequence and Series

Sub-Section: Geometric Sequence



Now let's consider another type of sequence, "Geometric" sequences.




Arithmetic Sequence



- Remember again
- He ate one chocolate bar on the first day.
- The next day, he ate two chocolate bars.
- Geometric sequence will suggest that Sam eats 4 chocolate bars on the 3rd day!

Discussion: In geometric sequence, how do we go from one term to the next?





Geometric Sequences

$$a, \quad ar, \quad ar^2, \quad ar^3$$

► Definition:

- A Geometric sequence is one where we keep multiplying or dividing by **the common ratio** to get the next term.

$$t_n = ar^{n-1}$$

- Where r is the common ratio, and a is the first term.

Discussion: Why do we have a power of $n - 1$ instead of n ? How many ratios do we multiply from t_1 to t_n ?



Question 14 Walkthrough.

Consider the geometric sequence defined by $t_n = 2 \cdot \left(\frac{1}{3}\right)^{n-1}$.

Identify the common ratio, first term and the 4th term.

NOTE: Geometric sequence is an exponential!



Question 15

Consider the geometric sequence defined by $t_n = 6 \cdot (2)^n$.


Identify the common ratio, first term and the 2nd term.

NOTE: Read the question carefully. Sometimes, they expand the $n - 1$ power to confuse you!



Space for Personal Notes


Sub-Section: Geometric Recurrence Relation



What about recurrence relations for geometric sequence?



Discussion: What must be the relationship between the current term (t_n) and the previous term (t_{n-1}) for a geometric sequence?



Recurrence Relation for Geometric Sequence



$$t_n = t_{n-1} \times r \text{ where } t_1 = a$$

Question 16

Consider the following n^{th} term rule for the geometric sequence

$$t_n = 2 \cdot 4^{n-1}$$

Find the recurrence relation which corresponds to it.

Sub-Section: Geometric Mean



How do we find a geometric term between two other geometric terms?



Exploration: Finding geometric mean.



- Consider three terms of a geometric sequence with the common ratio of r .

$$t_1 = a, t_2 = ar \text{ and } t_3 = ar^2$$

- Geometric mean simply means a _____.

Here we can say, t_2 is a geometric mean (average) of _____.

- Find the product of t_1 and t_3 .

$$t_1 \cdot t_3 = \underline{\hspace{2cm}}$$

- What should we do to $t_1 \cdot t_3 = a^2 r^2$ to find $t_2 = ar$?

$$t_2 = \underline{\hspace{2cm}}$$

The Geometric Mean



- **Definition:** The geometric mean of two numbers a and b is the geometric term in between a and b .

$$a \qquad \sqrt{ab} \qquad b$$

$$\text{Geometric Mean of } a \text{ and } b = \sqrt{ab}$$

Space for Personal Notes



TIP: Remember the similarity!


- Arithmetic Mean: We add the two and divide by 2.
- Geometric Mean: We multiply the two and square root.

Question 17

- a.** Find the geometric mean of 5 and 20.
- b.** Explain in words why 10 being a geometric mean makes sense.

Space for Personal Notes

Sub-Section: Geometric Series



Geometric Series

➤ **Definition:** Sum of first n geometric terms is given by:

$$S_n = \frac{a(r^n - 1)}{r - 1}$$

➤ Where n = number of terms, a = first term and r = common ratio.

Question 18 Walkthrough.

Consider the geometric sequence $t_n = 2 \cdot (3)^{n-1}$.

Find S_4 .

Question 19

Consider the geometric sequence with $t_n = 4 \cdot \left(\frac{1}{2}\right)^{n-1}$.

Find S_5 .

Space for Personal Notes

Sub-Section: Infinite Geometric Series



What does infinite geometric series even mean?



Context




► Imagine a frog jumping from one end of the pond to the other.

Here's the catch!

The frog always jumps half of remaining distance.

The frog always jumps half of remaining distance.



Question 20

Construct a geometric sequence for which its terms represent the % of the distance between the two ends of the pond that the frog covers in his n^{th} jump.

NOTE: Notice how, even if the frog jumps infinitely, its distance covered is still a finite number.

- Even if we add infinitely many geometric terms, the series(sum) can still be finite.
- This is called "Zeno's Paradox".

NOTE: If the common ratio is higher than 1 like the above discussion, the infinite series will not be a finite number.

The Infinite Geometric Series



- **Definition:** The sum of infinitely many geometric terms is given by

$$S_{\infty} = \frac{a}{1 - r}$$

 **IMPORTANT:** Only works when $-1 < r < 1$.

Question 21 Walkthrough.

Identify the first term, common ratio and hence find the infinite series.

$$1 + \frac{1}{10} + \frac{1}{100} + \frac{1}{1000} + \dots$$

Question 22

Identify the first term, common ratio and hence find the infinite series.

$$1 - \frac{2}{3} + \frac{4}{9} - \frac{8}{27} + \dots$$

NOTE: The common ratio must be between -1 and 1 for an infinite series to be a finite number.



Key Takeaways



- ✓ Geometric sequence has a common ratio between the next term and the current one.
- ✓ Geometric sequence is given by $t_n = ar^{n-1}$.
- ✓ Geometric mean of a and b is \sqrt{ab} .
- ✓ Geometric sum is given by $\frac{a(r^n - 1)}{r - 1}$.
- ✓ Infinite geometric sum is given by $\frac{a}{1 - r}$, where $-1 < r < 1$.



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

VCE Specialist Mathematics ½

Free 1-on-1 Consults

What Are 1-on-1 Consults?

- **Who Runs Them?** Experienced Contour tutors (45+ raw scores and 99+ ATARs).
- **Who Can Join?** Fully enrolled Contour students.
- **When Are They?** 30-minute 1-on-1 help sessions, after school weekdays, and all day weekends.
- **What To Do?** Join on time, ask questions, re-learn concepts, or extend yourself!
- **Price?** Completely free!
- **One Active Booking Per Subject:** Must attend your current consultation before scheduling the next :)

SAVE THE LINK, AND MAKE THE MOST OF THIS (FREE) SERVICE!

Booking Link

bit.ly/contour-specialist-consult-2025

