



Website: contoureducation.com.au | Phone: 1800 888 300
Email: hello@contoureducation.com.au

VCE Specialist Mathematics ½

Proofs II [2.2]

Workbook

Outline:

Proving Conditional Statements Pg 2-18 <ul style="list-style-type: none"> ➤ Conditional Statements ➤ Methods of Proving Conditional Statements ➤ Direct Proof ➤ Proof by Contrapositive ➤ Proof by Contradiction 	Universal and Existential Quantifiers Pg 24-34 <ul style="list-style-type: none"> ➤ Universal Quantifier ➤ Existence Quantifier ➤ Negation of Universal and Existence Statement ➤ Disproving a Universal Statement ➤ Disproving an Existence Statement
Converse Statements and Equivalent Statements Pg 19-23 <ul style="list-style-type: none"> ➤ Converse Statements 	Other Proof Techniques Pg 35-42 <ul style="list-style-type: none"> ➤ Telescopic Cancelling ➤ Proof by Induction

Learning Objectives:

- SM12 [2.2.1] - Direct proofs, proofs by contrapositive and contradiction.
- SM12 [2.2.2] - Converse and the equivalent of a conditional statement.
- SM12 [2.2.3] - Proofs involving the universal and existence quantifiers.
- SM12 [2.2.4] - Proofs by induction and telescoping series.



Section A: Proving Conditional Statements

Sub-Section: Conditional Statements


Context

- After this workbook, you will have the skills to tackle proof questions in Unit 3 and 4 level!

Exploration: Visualisation of Conditional Statements

$$A \Rightarrow B$$

Visualisation of $A \Rightarrow B$



If A , then B .

Conditional Statements

- Conditional Statement: Hypothesis implies conclusion
- Note: Notation for "implies": \Rightarrow

"Hypothesis \Rightarrow Conclusion"

NOTE: Order Matters!

Let's look at some questions together!

Question 1 Walkthrough.

Write a conditional statement for the following:

A customer will receive a coupon if they spend \$500.

If the customer spends \$500, then they will receive a coupon

Recall!

Active Recall: Conditional statements

► The relationship between the hypothesis and the conclusion is that the hypothesis implies the conclusion.



Your Turn!

Question 2

Write a conditional statement for the following:

Doing strength training grows your muscles.

If you do strength training, then you will grow your muscles

Question 3 Additional Question.

Write a conditional statement for the following:

All books in the library's collection are about history.

Space for Personal Notes

Sub-Section: Methods of Proving Conditional Statements



Okay, how do we prove them?



Proving Conditional Statements



1. Direct Proof

2. Proof by Contrapositive

3. Proof by Contradiction

Space for Personal Notes

Sub-Section: Direct Proof

Context

- The proof question for the most recent VCAA Exam 1 was a direct proof!

REMINDER

- From last week, we found that an even number can be written in the form:

$$2k, k \in \mathbb{Z} \rightarrow \text{integers}$$

- An odd number can be written in the form:

$$2k+1, k \in \mathbb{Z}$$

- If a number is divisible by 3, then it can be written in the form:

$$3k, k \in \mathbb{Z}$$

- If a number is not divisible by 3, then it can be written in the form:

$$3k+1, 3k+2, k \in \mathbb{Z}$$

- For proofs involving divisibility, we often need to split into cases.

Method 1: Direct Proof

- The simplest method of proof. These are the proofs we have been working on in [2.1].
- Here we are not altering the statement we need to prove.

Question 4 Walkthrough.

Prove that if n is even, then n^3 is divisible by 8.

$$\text{Let } n = 2k, k \in \mathbb{Z}$$

$$n^3 = (2k)^3$$

$$n^3 = 8k^3$$

$$\text{Let } k^3 = m, m \in \mathbb{Z}$$

$$n^3 = 8m$$

$\therefore n^3$ is divisible by 8.

NOTE: We did not change the statement itself! Hence, a direct proof.

Recall!

Active Recall: Direct proofs

► In a direct proof, we do not change the original statement.

Your Turn!

Question 5

Prove that for all integers m and n , if m is divisible by 5 and n is divisible by 2, then $10m + 3n$ is even.

$$\text{Let } m = 5k, n = 2\ell, k, \ell \in \mathbb{Z}$$

$$10m + 3n = 10(5k) + 3(2\ell)$$

$$= 50k + 6\ell$$

$$= 2(25k + 3\ell)$$

$$\text{Let } x = 25k + 3\ell, x \in \mathbb{Z}$$

$$= 2x \leftarrow \therefore 10m + 3n \text{ is even.}$$

Question 6 Additional Question.

Prove that if the last two digits of n are divisible by 4, then n is divisible by 4.

Space for Personal Notes

Sub-Section: Proof by Contrapositive



Context

- Check out this question from the Sample Exam for Specialist Mathematics Units 3 and 4 Exam 2, which asks about contrapositive statements.

Question 1

Consider the following statement.

'For all integers n , if n^2 is even, then n is even.'


Which one of the following is the contrapositive of this statement?

- A. For all integers n , if n^2 is odd, then n is odd.
- B. There exists an integer n such that n^2 is even and n is odd.
- C. There exists an integer n such that n is even and n^2 is odd.
- D. For all integers n , if n is odd, then n^2 is odd.
- E. For all integers n , if n is even, then n^2 is even.

What is a contrapositive statement?



Contrapositive Statement



(Contrapositive Statement of $A \Rightarrow B$) is $\neg B \Rightarrow \neg A$

NOTE: Swap the order and negate the statements.

① Flip
② negate.



Question 7 Walkthrough.

Write down a contrapositive of the following statement:

If you starve for a day, then you are hungry.

$$A \Rightarrow B$$

$$\neg B \Rightarrow \neg A$$

If you are not hungry, then you are not starving for a day

NOTE: We negate and flip the order.



Recall!



Active Recall: Contrapositive statements



► The contrapositive of the statement $A \Rightarrow B$ is:



Your Turn!



Question 8

Write down the contrapositive of the following statement:

If all the sides of a triangle have equal length, then the triangle is equilateral.

If a triangle is not equilateral, then the sides have different lengths



Question 9 Additional Question.

Write down the contrapositive of the following statement:

If x is an irrational number, then \sqrt{x} is irrational as well.

Discussion: What happens when you prove the contrapositive?

proving the original



Method 2: Proof by Contrapositive (Indirect Proof)



- ▶ Instead of proving $A \Rightarrow B$, we can prove its contrapositive $\neg B \Rightarrow \neg A$.

Prove a contrapositive statement instead.

- ▶ Considered to be an indirect proof, as the original statement is altered.

Steps:

1. Set up the contrapositive statement.
2. Prove the contrapositive statement to be true.
3. Conclude by saying "As contrapositive is true, the original statement is true."

Let's look at some questions together!

Question 10 Walkthrough.

Prove the following statement by contrapositive:

If $x^2 - 6x + 5$ is even, then x is odd.

If x is even, then $x^2 - 6x + 5$ is odd

Let $x = 2k, k \in \mathbb{Z}$

$$\begin{aligned} x^2 - 6x + 5 &= (2k)^2 - 6(2k) + 5 \\ &= 4k^2 - 12k + 4 + 1 \\ &= 2(k^2 - 6k + 2) + 1 \end{aligned}$$

Let $m = k^2 - 6k + 2, m \in \mathbb{Z}$

$$\text{—————} = 2m + 1 \leftarrow \text{odd}$$

Since the contrapositive is true, the original statement is true.

Discussion: Imagine if we tried to do a direct proof for the previous question. Would it be easy?

No!

Recall!

Active Recall: Proof by Contrapositive

Steps:

1. Set up the contrapositive statement.
2. Prove the contrapositive statement to be true.
3. Conclude by saying "As contrapositive is true, the original statement is true."

Your Turn!

Question 11

Prove the following conditional statement using contrapositive:

Let $x \in \mathbb{R}$. If x is irrational, then $\sqrt{x + \frac{1}{5}}$ is irrational.

If $\sqrt{x + \frac{1}{5}}$ is rational, then x is rational

$$\text{Let } \sqrt{x + \frac{1}{5}} = \frac{a}{b}, \quad a, b \in \mathbb{Z} \\ b \neq 0$$

$$x + \frac{1}{5} = \frac{a^2}{b^2}$$

$$x = \frac{a^2}{b^2} - \frac{1}{5}$$

$$x = \frac{5a^2 - b^2}{5b^2} = \frac{m}{n}$$

$$\text{Let } m = 5a^2 - b^2 \\ m \in \mathbb{Z} \\ n = 5b^2 \\ n \in \mathbb{Z} \setminus \{0\}$$

Question 12 Additional Question.

Prove the following statement below by first writing down its contrapositive.:

$$\text{Let } a, b \in \mathbb{R}^+. \text{ If } \sqrt{ab} \neq \frac{a+b}{2}, \text{ then } a \neq b.$$

Note: This means that if the geometric mean and arithmetic mean of two numbers a and b are not equal, then a and b are themselves not equal.

Space for Personal Notes

Sub-Section: Proof by Contradiction



Context

- Check out this proof-by-contradiction question from the Specialist Mathematics Units 3 and 4 Sample Exam 2. Give it a try after completing this sub-section!

Question 4 (3 marks)

Use proof by contradiction to prove that if n is odd, where $n \in \mathbb{N}$, then $n^3 + 1$ is even.

What is a contradicting statement?



Contradicting Statement



(Contradicting Statement of $A \Rightarrow B$) is $A \Rightarrow \neg B$

- Negate the conclusion.

Question 13

State the contradicting statement of the following:

If x is rational, then x^2 is rational.

If x is rational, then x^2 is irrational

Discussion: If we prove that the contradicting statement is false, then what are we also proving?



Original is true



Method 3: Proof by Contradiction (Indirect Proof)

To Prove $A \Rightarrow B$

Assume $A \Rightarrow \neg B$ is true \rightarrow math/logic

And show that the assumption is FALSE.

$B \vee B' =$

► Steps:

1. First, assume that the contradicting statement is true.
2. Show that the assumption has a contradiction, and is hence false.
3. Conclude by saying "Since the contradicting statement is false, the original statement is true."

► Considered to be an indirect proof, as the original statement is altered.

A B

Let's look at some questions together!

Question 14 Walkthrough.

Prove the simple statement below using contradiction:

$\log_5(9)$ is irrational

Assume: $\log_5(9)$ is rational

$$\log_5(9) = \frac{a}{b} \quad , \quad a, b \in \mathbb{Z} \quad b \neq 0$$

$$(5^{\frac{a}{b}})^b = (9)^b$$

$$5^a = 9^b$$

This is a contradiction

Since 5^a can't be equal to 9^b where a and b are integers, and $b \neq 0$

Since our assumption is FALSE, the original statement is TRUE



Active Recall: Steps for Proof by Contradictions

► Steps:

1. First, assume that the contradicting statement is _____,
2. Show that the assumption _____, and is hence false.
3. Conclude by saying "_____"



Active Recall: De Morgan's Law

$$\neg(A \wedge B) = \neg A \vee \neg B$$

$$\neg(A \vee B) = \neg A \wedge \neg B$$



Your Turn!

Question 15

Let $a, b \in \mathbb{R}$. Prove that if $a + b > 150$, then $a > 75$ or $b > 75$.

$$\text{If } a + b > 150, \text{ then } a \leq 75 \text{ and } b \leq 75$$

$$\text{①} + \text{②} \Rightarrow a + b \leq 150$$

← This

Directly contradicts our assumption

Therefore the original statement must be true

Question 16 Additional Question.

Use proof by contradiction to prove the following statement:

Let $x, y > 0$ and $x \neq y$. Show that $\frac{x}{y} + \frac{y}{x} > 2$.

Key Takeaways



- ✓ Direct proof involves proving without changing the conditional statement.
- ✓ The contrapositive of a statement $A \rightarrow B$ is given by $\neg B \rightarrow \neg A$.
- ✓ Proof by contrapositive involves proving the contrapositive of the conditional statement instead.
- ✓ Proof by contradiction requires first assuming that the contradicting statement is true, then showing contradiction.

Space for Personal Notes

Section B: Converse Statements and Equivalent Statements

Sub-Section: Converse Statements

Context: Sam is being Converse-

- ▶ Sam is being annoying and talks in the opposite direction of what Jacob says:

Jacob: "If you like chocolate, then you are Sam"

- ▶ What would Sam say?

Sam:

If you are Sam, then you like chocolate

- ▶ Here we call Sam the "converse" of Jacob.

$$\begin{array}{l} A \Rightarrow B \\ B \Rightarrow A \end{array}$$

Converse Statements

- ▶ **Definition:** Conditional statement that flows in the opposite direction.

(Converse Statement of $A \Rightarrow B$) is $B \Rightarrow A$

Question 17 Walkthrough.

For the following statement, write down the converse statement, and conclude whether the converse is true:

If a shape is a square, then it is a rectangle. \leftarrow True.

If the shape is a rectangle, then it is a square

\uparrow
False

NOTE: This is NOT the same as negation! It's simply an opposite flow.

Recall!



Active Recall: Converse statements



- ▶ The converse of the statement $A \Rightarrow B$ is

Your Turn!



Question 18

For the following statement, write down the converse statement, and conclude whether the converse is true:

If x is divisible by 2 and 5, then it is divisible by 10.

If x is divisible by 10, then it is divisible by 2 and 5

true

true

Discussion: If your friend told you that x is divisible by 2 and 5, is that equivalent to x is divisible by 10?



Yes

What happens if the converse is also true?

Equivalent Statements (Biconditional)

- It is a conditional statement where if the original is proven to be true, its converse is ALWAYS true.

$$A \Rightarrow B \text{ and } B \Rightarrow A$$

$$A \Leftrightarrow B$$

A is true, if and only if B

- In description, B is true if and only if A is true.
- To prove equivalent statements, we prove each direction separately.

Question 19 Walkthrough.

$$A \Leftrightarrow B$$

Let n be an integer. Prove that n is even, if and only if $3n + 3$ is odd.

$$A \Rightarrow B$$

$$\text{Let } n = 2k, k \in \mathbb{Z}$$

$$3n + 3 = 3(2k) + 3$$

$$= 6k + 2 + 1$$

$$= 2(3k + 1) + 1$$

$$\text{Let } m = 3k + 1, m \in \mathbb{Z} \Rightarrow 2m + 1$$

$A \Rightarrow B$ is true

$$B \Rightarrow A$$

If $3n + 3$ is odd, then n is even

Using the contrapositive,
If n is odd, then $3n + 3$ is even

$$\text{Let } n = 2k + 1$$

$$3n + 3 = 3(2k + 1) + 3$$

$$= 6k + 6$$

$$= 2(3k + 3)$$

$$\text{Let } l = 3k + 3, l \in \mathbb{Z}$$

$$= 2l$$

$B \Rightarrow A$ is also true

Since the contrapositive is true, the original statement is also true

NOTE: For if and only if (equivalent statement), we must prove both converse statements.

$$\therefore A \Leftrightarrow B$$

Recall!

Active Recall: Proving biconditional statements

► To prove an "if and only if" statement $A \Leftrightarrow B$, we need to show both $A \Rightarrow B$ and $B \Rightarrow A$.

Your Turn!

Question 20

$$A \Leftrightarrow B$$

Let n be an integer. Prove that n is odd, if and only if n^2 is odd.

$$A \Rightarrow B$$

If n is odd, n^2 is odd

$$\text{Let } n = 2k+1, k \in \mathbb{Z}$$

$$n^2 = (2k+1)^2$$

$$= 4k^2 + 4k + 1$$

$$= 2(2k^2 + 2k) + 1$$

$$\text{Let } m = 2k^2 + 2k$$

$$= 2m + 1$$



$A \Rightarrow B$ is true.

$$B \Rightarrow A$$

If n^2 is odd, n is odd

Using the contrapositive,
If n is even, the n^2 is even

$$\text{Let } n = 2k, k \in \mathbb{Z}$$

$$n^2 = 4k^2$$

$$= 2(2k^2)$$

$$\text{Let } m = 2k^2$$

$$= 2m \quad \checkmark$$

Since the contrapositive is true, the original is also true

$B \Rightarrow A$ is true.

◦◦ $A \Leftrightarrow B$.

Question 21 Additional Question.

Let x be a real number. Show that $x^2 + y^2 = 0$, if and only if $x = 0$ and $y = 0$.

Key Takeaways



- ☒ The converse statement of $A \rightarrow B$ is given by $B \rightarrow A$.
- ☒ Equivalent statement is when a statement and its converse both are proved to be true.
- ☒ "If and only if" stands for equivalent statement.

Space for Personal Notes

Section C: Universal and Existential Quantifiers

Sub-Section: Universal Quantifier

How do we represent all members of a given set?

Universal Quantifiers

➤ Universal quantifier is a way to represent all members of a given set.

For all real numbers x , x^2 is never negative.

➤ Notation: \forall (Universal Quantifier) "For all"

$$\forall x \in \mathbb{R}, x^2 \geq 0$$

Discussion: Where does the symbol \forall come from?

Upside down A
↑
All

Question 22 Walkthrough.

Convert the following conditional statement using the universal quantifier.

For all real numbers x , e^x is strictly positive.

$$\forall x \in \mathbb{R}, e^x > 0$$

True

Recall!



Your Turn!



Question 23

Rewrite the following statement using the universal quantifier:

For all integers n , $n^2 - 4n$ is an integer.

$$\forall n \in \mathbb{Z}, n^2 - 4n \in \mathbb{Z}$$

Question 24 Additional Question.

Rewrite the following statement using the universal quantifier:

For all natural numbers n , n is at most equal to n^2 .

$$\forall n \in \mathbb{N}, n \leq n^2$$

Sub-Section: Existence Quantifier

How do we represent certain members of a given set?

Existence Quantifiers

► existence quantifier is a way to represent certain members of a given set.

There exists an integer such that $x^2 - x - 12 = 0$.

► Notation: \exists (Existential Quantifier) 'there exists'

$\exists x \in \mathbb{Z}, x^2 - x - 12 = 0$

Discussion: How can we prove the existence statement?

Find an example

Question 25 Walkthrough.

Convert the following conditional statement using the existence quantifier.

There exist a real number x such that $x^2 + 4 = 16$.

$\exists x \in \mathbb{R}, \text{ s.t. } x^2 + 4 = 16$

Recall!

Active Recall: Existence quantifier

► The symbol for the existence quantifier \exists represents some members of the set.

Your Turn!

Question 26

Rewrite the following statement using the existence quantifier.

There exists a positive whole number n such that $2^n = 4$.

$$\exists n \in \mathbb{N} \text{ s.t. } 2^n = 4$$

\mathbb{Z}^+

Question 27 Additional Question.

Rewrite the following statement using the existence quantifier.

There exists a positive real number x such that $x^2 < x$.

$$\exists x \in \mathbb{R}^+ \text{ s.t. } x^2 < x$$

Sub-Section: Negation of Universal and Existence Statement

Discussion: What would be the opposite of saying that all living humans breathe?



Discussion: What would be the opposite of saying that some humans are taller than 190 cm?



Negation of Universal and Existence Statements



- ▶ They are opposites of each other (with opposite conclusions).

\neg Universal Statement = Existence Statement with Opposite Conclusion

- ▶ And vice versa.

Question 28 Walkthrough.

Write down the negation of the following statements.

- a. For all natural numbers, $3n \geq 2n - 1$.

$$\neg (\forall n \in \mathbb{N}, 3n \geq 2n - 1)$$

$$\exists n \in \mathbb{N}, 3n < 2n - 1$$

There exists a natural number n , such that $3n < 2n - 1$

- b. There exists a real number x such that $x^2 = 4$.

$$\neg (\exists x \in \mathbb{R}, \text{ s.t. } x^2 = 4)$$

$$\forall x \in \mathbb{R}, \text{ s.t. } x^2 \neq 4$$

For all real numbers x , $x^2 \neq 4$

Recall!



Active Recall: Negation involving the universal and existence quantifiers.



\neg Universal statement = \exists , negate conclusion
 \neg Existence statement = \forall , negate conclusion

Your Turn!



Question 29

Write down the negation for the following statements below.

- a. If n is a natural number, then $n + 1 > n$.

There exists a natural number n , such that $n + 1 < n$

- b. There exists an integer k such that $k^2 = k + 4$.

For all integers k , $k^2 \neq k + 4$

Question 30 Additional Question.

Write the negation for the following statements below.

- a. If n is a natural number, then $6n^2 + 4n + 1$ is divisible by n .

- b. There exist integers p and q such that $q \neq 0$ such that $\pi = \frac{p}{q}$.

Sub-Section: Disproving a Universal Statement.



Context

- Consider the following questionable statement:

"All of the vehicles in the carpark are black."

- How can you disprove this statement?

→ Find a vehicle that isn't black

- Basically, you have just provided a counter example.

Disproving Universal Statement



- We prove the *opposite* (negation) existence statement.

- We call this proof by counter example.

- 🇬🇧 Giving a counter example will be proving an opposite existence statement.

Question 31 Walkthrough.

∇

Disprove the following statement: For all positive integers m , if m is prime then $m^2 + 4$ is also prime.

Provide counter example

$$m = 2,$$

$$m^2 + 4 = 8$$

Since $m=2$ is a counter example, we have disproven the statement

NOTE: We simply give ourselves a counter example to disprove a universal statement.



Recall!



Active Recall: Disproving a universal statement.



► To show that a “for all ...” statement is not true, it suffices to provide a _____.

Your Turn!



Question 32

Disprove the following statement:

$\forall a, b \in \mathbb{R}$, if $a^2 = b^2$ then $a = b$.

$$a = 2$$

$$(1)^2 = (-1)^2$$

$$b = -2$$

↑
counter example.
 $a \neq b$

Question 33 Additional Question.

Disprove the statement given below:

For all positive integers n , $2^n > n^2$.

Sub-Section: Disproving an Existence Statement

Disproving Existence Statements

► We prove the opposite (negation) universal statement

Question 34 Walkthrough.

Disprove the following statement: There exists $n \in \mathbb{N}$ such that $n^2 + 9n + 20$ is a prime number.

$\exists n \in \mathbb{N}$ s.t. $n^2 + 9n + 20$ is prime

$\forall n \in \mathbb{N}$ s.t. $n^2 + 9n + 20$ is NOT prime

$$n^2 + 9n + 20 = (n+5)(n+4)$$

(1) Since $n \in \mathbb{N}$, $n+5 \neq 1$
 $n+5 \neq n^2 + 9n + 20$
 $n+4 \neq 1$
 $n+4 \neq n^2 + 9n + 20$
 $\therefore \forall n \in \mathbb{N}$, $n^2 + 9n + 20$ is not prime

NOTE: To disprove their exist statement, you must show the opposite universal statement as true.

Recall!!

Active Recall: Disproving an existence statement.

► To disprove a "there exists ..." statement, we need to show _____

Your Turn!



Question 35

Disprove the following statements:

There exists a real number x , such that $10 + 3x^2 = 3 + x^2$.

$$\forall x \in \mathbb{R}, 10 + 3x^2 \neq 3 + x^2$$

$$2x^2 \neq -7$$

↑
true mathematical
statement

We have disproven the OG statement

Question 36 Additional Question.

Disprove the following statement below:

There exists $n \in \mathbb{N}$ so that $n^3 - 2n^2 + 5n + 4$ is divisible by $n + 2$.



Key Takeaways

- ✓ There exists a quantifier \exists .
- ✓ For all quantifiers \forall .
- ✓ To prove a there exists statement, simply give an example.
- ✓ To prove for all statements, you must prove for all values.
- ✓ To disprove a there exist statement, prove the opposite for all statements.
- ✓ To disprove a for-all statement, prove the opposite there exist statement.

Space for Personal Notes


Section D: Other Proof Techniques

Sub-Section: Telescopic Cancelling



Telescopic Cancelling

- Telescoping series is a series (sum of terms) where all terms cancel out except for the first and last.
- The name comes from the visualisation as shown below.



- Proofs of telescoping series can be done via telescopic cancelling.

Space for Personal Notes

Question 37 Walkthrough.

a. Using partial fractions, find the values of a and b such that:

$$\frac{3}{k(k+1)} = \frac{a}{k} + \frac{b}{k+1}$$

$$= \frac{a(k+1) + bk}{k(k+1)}$$

$$3 = a(k+1) + bk$$

$$\text{sub } k=0$$

$$3 = a(1)$$

$$a=3$$

$$\text{sub } k=-1$$

$$-b=3$$

$$b=-3$$

$$\frac{3}{k(k+1)} = \frac{3}{k} - \frac{3}{k+1}$$

b. Hence, prove that:

$$k=1$$

$$k=2$$

$$k=3$$

$$\frac{3}{1 \times 2} + \frac{3}{2 \times 3} + \dots + \frac{3}{n \times (n+1)} = \frac{3n}{n+1}$$

$$\left(3 - \frac{3}{2}\right) + \left(\frac{3}{2} - \frac{3}{3}\right) + \left(\frac{3}{3} - \frac{3}{4}\right) \dots \dots \left(\frac{3}{n} - \frac{3}{n+1}\right)$$

$$= 3 - \frac{3}{n+1}$$

$$= \frac{3(n+1) - 3}{n+1} = \frac{3n + 3 - 3}{n+1} = \frac{3n}{n+1}$$

Recall!

Active Recall: Telescoping sums

► The key idea for telescoping sums is to _____.

Your Turn!

Question 38

- a. Using partial fraction decomposition, find values of A and B such that:

$$\frac{1}{n^2 + 3n + 2} \equiv \frac{A}{n+1} + \frac{B}{n+2} \quad \rightarrow = \frac{1}{n+1} - \frac{1}{n+2}$$

$$\frac{1}{(n+2)(n+1)} = \frac{A(n+2) + B(n+1)}{(n+2)(n+1)}$$

$$1 = A(n+2) + B(n+1)$$

$n = -1$
 $1 = A$
 $n = -2$
 $1 = -B$
 $B = -1$

- b. Hence, prove that:

$$\sum_{n=1}^k \frac{1}{n^2 + 3n + 2} = \frac{k}{2(k+2)}$$

$n=1$
 $\left(\frac{1}{2} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \dots + \left(\frac{1}{k+1} - \frac{1}{k+2}\right)$

$n=2$
 $= \frac{1}{2} - \frac{1}{k+2}$
 $= \frac{k+2-2}{2(k+2)}$
 $= \frac{k}{2(k+2)}$

Question 39 Additional Question.

By using the fact that $\frac{2(2k+1)}{k^2(k+1)^2} = \frac{2}{k^2} - \frac{2}{(k+1)^2}$, show that:

$$\frac{2 \times 5}{2^2 \times 3^2} + \dots + \frac{2(2n+1)}{n^2(n+1)^2} = \frac{(n-1)(n+3)}{2(n+1)^2}$$

Sub-Section: Proof by Induction



Context

- ▶ Check out this question from the Specialist Mathematics Units 3 and 4 Sample Exam.

Question 3 (4 marks)

Prove by mathematical induction that the number $9^n - 5^n$ is divisible by 4 for all $n \in \mathbb{N}$.



Proof by Induction

- ▶ We will be following the given steps: " TAPE ".

🔗 **Step T:** Test the statement for its first possible value.

🔗 **Step A:** Assume that the statement is true for $n = k$.

🔗 **Step P:** Prove that if the statement holds true for $n = k$, It also holds true for $n = k + 1$.

- ▶ Add "By Assumption".

🔗 **Step E:** "By the principle of mathematical induction, the statement is true for a set of values."

Space for Personal Notes

Question 40 Walkthrough.

$$2(1) - 1 = 1$$

Prove that for $n \in \mathbb{N}$, $1 + 3 + 5 \dots (2n - 1) = n^2$.

Test $n = 1$

Assume true $n = k$

$$1 = 1^2 \checkmark$$

$$1 + 3 + 5 \dots + (2k - 1) = k^2$$

Prove $n = k + 1$

$$1 + 3 + 5 \dots + (2k - 1) + (2(k + 1) - 1) = (k + 1)^2$$

$$k^2 + 2k + 1 = k^2 + 2k + 1 \checkmark$$

If the statement is true for $n = k$

Then it is also true for $n = k + 1$


However, the statement is also true for $n = 1$

Therefore $n \in \mathbb{N}$ $1 + 3 + 5 + \dots + 2n - 1 = n^2$ by induction

Exploration: How do we visualise the idea of proof by induction?

- Imagine we have dominos of statements we need to prove.
- Each domino is a statement with a specific value of n .

Dominoes of Statement



- ▶ In Step 1: We test/prove the first statement.
- ▶ In Steps A and P: We prove the relationship between a “previous” domino block, and the “next” one.
- ▶ If one domino falls, the next one also falls.
- ▶ Since the first one falls, all of them fall!

Recall!



Active Recall: Proofs by induction



- ▶ When doing a proof-by-induction question, we have four key steps:
1. _____ the _____ holds.
 2. _____ the statement holds for an $n = \underline{\hspace{1cm}}$ in the required set of values for k .
 3. _____ that the statement holds for _____.
 4. _____ using the principle of _____ that the statement holds for all k .

Space for Personal Notes

Your Turn!

Question 41

Prove that for all $n \in \mathbb{N}$, we have:

$$1 + 2 + \dots + n = \frac{n(n+1)}{2}$$

NOTE: $1 + 2 + \dots + n$ can be written as:

$$\sum_{i=1}^n i$$

Test $n=1$

$$1 = \frac{1(1+1)}{2}$$

$$= \frac{2}{2}$$

$$= 1 \quad \checkmark$$

Assume $n=k$ is true

$$1 + 2 + 3 + \dots + k = \frac{k(k+1)}{2}$$

Prove $n=k+1$

$$1 + 2 + 3 + \dots + k + (k+1) = \frac{(k+1)(k+2)}{2}$$

$$\frac{k(k+1)}{2} + k+1 = \frac{(k+1)(k+2)}{2}$$

$$\frac{k(k+1) + 2k+2}{2} = \frac{(k+1)(k+2)}{2}$$

$$LHS = \frac{k^2 + 3k + 2}{2}$$

$$RHS = \frac{k^2 + 2k + k + 2}{2}$$

$$= \frac{k^2 + 3k + 2}{2}$$

$$LHS = RHS \quad \checkmark$$

Therefore $1+2+3+\dots+n = n(n+1)/2$ by induction

Question 42 Additional Question.

Prove by induction that for any natural number $n \geq 1$, $n < 2^n$.

DM for discord
 alex.watson 715 douglas.mann 27
 alex.w@contour-education.com.au
 douglas
 Class code:
 C O 3

Key Takeaways



- ✓ For telescoping series proof, always look for terms to cancel each other.
- ✓ For induction proof always use TAPE: Test for the first case, assume it's true for $n = k$, prove for $n = k + 1$ and explain at the end.

Space for Personal Notes



Contour Check

Learning Objective: [2.2.1] – Direct proofs, proofs by contrapositive and contradiction.

Key Takeaways

- ☐ Direct proof is _____ the conditional statement.
- ☐ The contrapositive of a statement $A \rightarrow B$ is given by _____.
- ☐ Proof by contrapositive is proving the _____.
- ☐ Proof by contradiction requires first _____ and showing contradiction.

Learning Objective: [2.2.2] – Converse and the equivalent of a conditional statement.

Key Takeaways

- ☐ Converse statement of $A \rightarrow B$ is given by _____.
- ☐ Equivalent statement is when _____ both have to be true.
- ☐ "If and only if" stands for _____.

Learning Objective: [2.2.3] – Proofs involving the universal and existence quantifiers.

Key Takeaways

- ☐ There exists a quantifier \exists .
- ☐ For all quantifiers \forall .
- ☐ To prove a there exists statement, simply _____.
- ☐ To prove for all statements, you must prove for _____.
- ☐ To disprove a there exist statement, _____.
- ☐ To disprove a for-all statement, _____.

Learning Objective: [2.2.4] – Proofs by induction and telescoping series.

Key Takeaways

- ☐ For telescoping series proof, always look for terms to _____.
- ☐ For induction proof always use TAPE: _____
_____.



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

VCE Specialist Mathematics ½

1-on-1 Maths Consults

What Are 1-on-1 Maths Consults?

- Individual 30-minute consultations via Zoom with a Contour tutor, where you can ask questions, clarify doubts, and get tips, advice, and support in a one-on-one format.
- Complimentary (yes, entirely free) with your Contour enrolment (as long as you're enrolled in a Maths subject at Contour). You can book up to a week in advance.

SAVE THE LINK. AND MAKE THE MOST OF THIS (FREE) SERVICE!

Booking Link

bit.ly/maths-consults

