



Website: contoureducation.com.au | Phone: 1800 888 300

Email: hello@contoureducation.com.au

VCE Specialist Mathematics ½

Trigonometry II [3.2]

Workbook

Outline:



| | | | |
|--|----------|--|----------|
| <u>Introduction to Circular Functions</u> | Pg 2-9 | <u>Particular and General Solutions</u> | Pg 15-22 |
| ➤ Radians and Degrees | | ➤ Particular Solutions | |
| ➤ Unit Circle | | ➤ General Solutions | |
| ➤ Period | | <u>Graphs of Sine and Cosine</u> | Pg 23-29 |
| ➤ Pythagorean Identities | | ➤ Basics of Sine and Cosine Functions | |
| ➤ Exact Values | | ➤ Graphing Sine and Cosine Functions | |
| <u>Symmetry</u> | Pg 10-14 | <u>Graphs of Tangent</u> | Pg 30-33 |
| ➤ Supplementary Relationships | | ➤ Basics of Tangent Graphs | |
| | | ➤ Graphing Tangent Functions | |

Learning Objectives:

- SM12 [3.2.1] - Find Trig Ratios of Supplementary Relationships
- SM12 [3.2.2] - Find Particular and General Solutions
- SM12 [3.2.3] - Graph Sine, Cosine and Tangent functions



Section A: Introduction to Circular Functions

Sub-Section: Radians and Degrees

Radians and Degrees

$$1^c = \left(\frac{180}{\pi}\right)^0$$

$$1^0 = \left(\frac{\pi}{180}\right)^c$$

$$180^0 = \pi^c$$



$$360^0 = 2\pi$$



Question 1


- a. Find $\left(\frac{\pi}{4}\right)^c$ in degrees.

$$\frac{\pi}{4} = 45^0$$


- b. Find 12^0 in radians.

$$12 \times \frac{\pi}{180} = \frac{12\pi}{180} = \frac{\pi}{15}$$

Sub-Section: Unit Circle



Active Recall



$\sin = \frac{O}{H}$

$\cos = \frac{A}{H}$


$\tan = \frac{O}{A}$

Space for Personal Notes

What is a unit circle, and how do we use it?

Exploration: Unit Circle

- The unit circle is simply a circle of radius 1.
- Angles are measured from the positive x-axis anticlockwise.
- It can be divided into four quadrants:



- We can use the elementary definition of the trigonometric functions.
- Select the option below!

$$\sin(\theta) = [X \text{ Value}, Y \text{ Value}, \text{Gradient}]$$


$$\cos(\theta) = [X \text{ Value}, Y \text{ Value}, \text{Gradient}]$$

$$\tan(\theta) = [X \text{ Value}, Y \text{ Value}, \text{Gradient}]$$



Unit Circle

➤ The unit circle is simply a circle of radius 1.




$$\sin(\theta) = y$$

$$\cos(\theta) = x$$

$$\tan(\theta) = \text{gradient}$$

Discussion: For which quadrant is cos, sin and tangent positive?



Sub-Section: Period

Discussion: For what angle does cos, sin and tangent repeats itself?

2π π (180°)



Period of a Trigonometric Function



period of sin(nx) and cos(nx) functions = $\frac{2\pi}{|n|}$

period of tan(nx) functions = $\frac{\pi}{|n|}$

where n = coefficient of x .

Question 2

Find the period of each of the following trigonometric functions:

a. $p(x) = \tan(2x)$


$$\text{period} = \frac{\pi}{n} = \frac{\pi}{2}$$

b. $q(x) = \cos\left(\frac{5}{2}x + \frac{\pi}{3}\right)$


$$\text{period} = \frac{2\pi}{5/2} = 2\pi \times \frac{2}{5} = \frac{4\pi}{5}$$

Sub-Section: Pythagorean Identities

Discussion: What does $\sin^2(\theta) + \cos^2(\theta)$ equal to?

$$\begin{array}{c} \uparrow \quad \uparrow \quad = 1 \\ y^2 + x^2 = 1 \end{array}$$


Pythagorean Identities



$$\sin^2(\theta) + \cos^2(\theta) = 1$$

➤ Can be used for finding one trigonometry function by using the other.

Space for Personal Notes

How can we use it?



Question 3 Walkthrough.

Find the value of $\sin(x)$ given that $\cos(x) = \frac{1}{4}$ and x is the first quadrant.

+ve

$$\sin^2(x) + \cos^2(x) = 1$$

$$\sin^2(x) + \left(\frac{1}{4}\right)^2 = 1$$

$$\sin^2(x) = \frac{15}{16}$$

Since $\sin(x) \geq 0$ $\sin(x) = \pm \sqrt{\frac{15}{16}}$

$$\sin(x) = \sqrt{\frac{15}{16}}$$

NOTE: Always show the rejection by the quadrant.



Question 4

Find the value of $\cos(x)$ given that $\sin(x) = \frac{1}{3}$ and x is the second quadrant.

cos Ev

$-\frac{2\sqrt{2}}{3}$


$$\cos^2(x) = 1 - \frac{1}{9}$$

$$\cos^2(x) = \frac{8}{9}$$

$$\cos(x) = \pm \frac{2\sqrt{2}}{3}$$

$$\cos(x) = -\frac{2\sqrt{2}}{3}$$

Sub-Section: Exact Values



Exact values are super important to remember!



The Exact Values Table



| x | $0 (0^\circ)$ | $\frac{\pi}{6} (30^\circ)$ | $\frac{\pi}{4} (45^\circ)$ | $\frac{\pi}{3} (60^\circ)$ | $\frac{\pi}{2} (90^\circ)$ |
|-----------|---------------|----------------------------|----------------------------|----------------------------|----------------------------|
| $\sin(x)$ | 0 | $\frac{1}{2}$ | $\frac{\sqrt{2}}{2}$ | $\frac{\sqrt{3}}{2}$ | 1 |
| $\cos(x)$ | 1 | $\frac{\sqrt{3}}{2}$ | $\frac{\sqrt{2}}{2}$ | $\frac{1}{2}$ | 0 |
| $\tan(x)$ | 0 | $\frac{1}{\sqrt{3}}$ | 1 | $\sqrt{3}$ | Undefined |


TIP: Use the fact that sin is the y value, cos is the x value, and the tangent is the gradient to remember the values well!



Discussion: With your classmate next to you, test each other on the exact value table!



Discussion: The exact value table only has first-quadrant angles! How do we evaluate other quadrants?



Symmetry

Section B: Symmetry


Sub-Section: Supplementary Relationships

180°

What does reflection in the y-axis look like?

Exploration: Reflection in y-axis

- Consider the unit circle.




- Reflect the angle around the y-axis on the unit circle above.
- What is the angle in terms of θ ?

$$\pi - \theta$$

Question 5

Consider the angle $\frac{\pi}{6}$.

Find the angle after the reflection in the y-axis.




$$\pi - \frac{\pi}{6} = \frac{5\pi}{6}$$

What does a reflection in the x -axis look like?

Exploration: Reflection in x -axis

- Consider the unit circle.




- Reflect the angle around the x -axis on the unit circle above.
- What is the angle in terms of θ ?

Question 6

Consider the angle $\frac{\pi}{6}$

Find the angle after the reflection in the x -axis.



$$360^\circ - \theta$$

$$2\pi - \frac{\pi}{6} = \frac{11\pi}{6}$$

$$\boxed{-\frac{\pi}{6}} \quad \text{or} \quad \boxed{\frac{11\pi}{6}}$$


NOTE: Simply make the angle negative!

What does reflection in both axes look like?



Exploration: Reflection on Both Axes

- Consider the unit circle.




- Reflect the angle around both axes on the unit circle above.
- What is the angle in terms of θ ?

$$\pi + \theta$$

Question 7

Consider the angle $\frac{\pi}{6}$.

Find the angle after the reflection in both axes.




$$\pi + \frac{\pi}{6} = \frac{7\pi}{6}$$

Let's summarise!



Supplementary Relationships



➤ Simply look at the quadrant to find the correct sign.

🌀 Second Quadrant ($\pi - \theta$):

$$\cos(\pi - \theta) = -\cos(\theta)$$

$$\sin(\pi - \theta) = +\sin(\theta)$$

$$\tan(\pi - \theta) = -\tan(\theta)$$

🌀 Third Quadrant ($\pi + \theta$):

$$\cos(\pi + \theta) = -\cos(\theta)$$

$$\sin(\pi + \theta) = -\sin(\theta)$$

$$\tan(\pi + \theta) = +\tan(\theta)$$


Fourth Quadrant $(-\theta)$:

$$\cos(-\theta) = +\cos(\theta)$$

$$\sin(-\theta) = -\sin(\theta)$$

$$\tan(-\theta) = -\tan(\theta)$$

Try the following question!



Question 8

If $\sin(\theta) = -0.6$ where θ is a third quadrant angle, evaluate the following.

a. $\sin(\pi + \theta)$ \rightarrow quad 1 \Rightarrow sin (+ve)
 $= 0.6$

b. $\cos(\pi + \theta)$ \Rightarrow
 $\cos^2(\pi + \theta) + \sin^2(\pi + \theta) = 1$
 $\cos^2(\pi + \theta) = 1 - (0.6)^2$
 $= 1 - 0.36$
 $= 0.64$
 $\Rightarrow \cos(\pi + \theta) = \pm 0.8$
 (quad)
 $\cos(\pi + \theta) = -0.8$

c. $\tan(\pi - \theta)$
 $\tan(-\theta + \pi)$
 2nd quad + 180°
 $\tan(4th \text{ quad}) = \frac{\sin(-\theta + \pi)}{\cos(-\theta + \pi)} = \frac{-0.6}{0.8} = -\frac{3}{4}$

NOTE: The aim of the question is to convert the angle to theta!

Section C: Particular and General Solutions

Sub-Section: Particular Solutions


Active Recall: Period of Trigonometric Function

period of $\sin(nx)$ and $\cos(nx)$ functions = $\frac{2\pi}{n}$

period of $\tan(nx)$ functions = $\frac{\pi}{n}$

where n = coefficient of x .

Discussion: How often would the solution to $\sin(x) = \frac{1}{2}$ repeat?



every 2π

↳ every period

Particular Solutions

- Solving trigonometric equations for finite solutions.
- Steps:
 - 🔍 Make the trigonometric function the subject.
 - 🔍 Find the necessary angle for one period.
 - 🔍 Solve for x by equating the necessary angles to the inside of the trigonometric functions.
 - 🔍 Add and subtract the period to find all other solutions in the domain.

Question 9 Walkthrough.

Solve the following equations for x over the domains specified.



$$2 \sin(2x + \pi) - \sqrt{3} = 0 \text{ for } x \in [0, 2\pi]$$

$$\sin(2x + \pi) = \frac{\sqrt{3}}{2}$$

$$\frac{2\pi}{2} = \pi$$

$$2x + \pi = \frac{\pi}{3}$$

$$2x + \pi = \frac{2\pi}{3}$$

$$2x = -\frac{2\pi}{3}, -\frac{\pi}{3}$$

$$x = -\frac{\pi}{3}, -\frac{\pi}{6} + \text{period}$$

$$x = \frac{2\pi}{3}, \frac{5\pi}{6}, \frac{5\pi}{3}, \frac{11\pi}{6}$$

Question 10

Solve the following equations for x over the domains specified.

a. $\sin\left(x - \frac{\pi}{2}\right) = -\frac{1}{2}$ for $x \in [-\pi, \pi]$

$$x - \frac{\pi}{2} = \pi + \frac{\pi}{6}, -\frac{\pi}{6}$$

$$x = \frac{7\pi}{6} + \frac{\pi}{2}, -\frac{\pi}{6} + \frac{\pi}{2}$$

$$x = \frac{10\pi}{6}, \frac{2\pi}{6}$$

$$x = \frac{5\pi}{3}, \frac{\pi}{3}, -\frac{\pi}{3}, -\frac{5\pi}{3}$$

$$\text{period} = \frac{2\pi}{1} = 2\pi$$

$$x = -\frac{\pi}{3}, \frac{\pi}{3}$$

b. $2 \cos\left(2x + \frac{\pi}{6}\right) + 1 = 0$ for $x \in [0, 2\pi]$

$$\cos\left(2x + \frac{\pi}{6}\right) = -\frac{1}{2}$$

$$2x + \frac{\pi}{6} = \frac{2\pi}{3}, \frac{4\pi}{3}$$

$$2x = \frac{2\pi}{6}, \frac{7\pi}{6}$$

$$x = \frac{\pi}{4}, \frac{7\pi}{12}, \frac{5\pi}{4}, \frac{13\pi}{12}$$

Question 11 Walkthrough.

Solve the following equations for x over the domains specified.

$$\tan\left(x + \frac{\pi}{3}\right) - \sqrt{3} = 0 \text{ for } x \in [0, 2\pi]$$


$$\tan\left(x + \frac{\pi}{3}\right) = \sqrt{3}$$

$$x + \frac{\pi}{3} = \frac{\pi}{3}$$

$$x = 0$$

$$\text{period} = \frac{\pi}{n} = \frac{\pi}{1} = \pi$$

$$x = 0, \pi, 2\pi$$



Discussion: Why do we need to find one angle only for tangents?



$$\textcircled{1} \tan\left(x - \frac{\pi}{3}\right) = \frac{1}{\sqrt{3}}$$



$$\textcircled{3} \quad x - \frac{\pi}{3} = \frac{\pi}{6}$$

$$x = \frac{\pi}{2}$$

$$\textcircled{4} \quad \text{Period} = \frac{\pi}{n} = \frac{\pi}{1} = \pi$$

$$x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}.$$

Because the next one is a period away for tan (tan's period is half of the other ones)

Question 12

Solve the following equations for x over the domains specified.

$$\sqrt{3} \tan\left(x + \frac{\pi}{4}\right) + 1 = 0 \text{ for } x \in (0, 3\pi)$$

$$\tan\left(x + \frac{\pi}{4}\right) = -\frac{1}{\sqrt{3}}$$

$$x + \frac{\pi}{4} = -\frac{\pi}{6}$$

$$x = -\frac{\pi}{6} - \frac{\pi}{4}$$


$$x = \frac{-2\pi - 3\pi}{12}$$

$$x = -\frac{5\pi}{12}$$


$$\text{period} = \frac{\pi}{1} = \pi$$

$$x = -\frac{5\pi}{12}, \frac{7\pi}{12}, \frac{19\pi}{12}, \frac{31\pi}{12}$$

Sub-Section: General Solutions



Discussion: How many solutions would there be for $x \in \mathbb{R}$?



Infinite

General Solutions



➤ Finding infinite solutions to a trigonometric equation.

➤ Steps:

1. Make the trigonometric function the subject.

2. Find the necessary angle for one period.

3. Solve for x by equating the necessary angles to the inside of the trigonometric functions.

4. Add Period $\cdot n$ where $n \in \mathbb{Z}$.

period
 $+2\pi n$ $n \in \mathbb{Z}$

Space for Personal Notes


Question 13 Walkthrough.

Find the general solutions to the following equations:

$$2 \sin\left(2x + \frac{\pi}{2}\right) - 1 = 0$$

$$\sin\left(2x + \frac{\pi}{2}\right) = \frac{1}{2}$$

① 1 period



$$2x + \frac{\pi}{2} = \frac{\pi}{6}, \pi - \frac{\pi}{6}$$

$$2x + \frac{\pi}{2} = \frac{\pi}{6}, \frac{5\pi}{6}$$

$$2x = -\frac{\pi}{3}, \frac{\pi}{3}$$

$$x = -\frac{\pi}{6}, \frac{\pi}{6}$$

② + period $\cdot n$
period = $\frac{2\pi}{2} = \pi$

$$x = -\frac{\pi}{6} + \pi \cdot n, \frac{\pi}{6} + \pi n, \quad n \in \mathbb{Z}$$

NOTE: The steps are exactly the same as a particular solution except for adding the period. We simply add period $\times n$ instead.

ALSO NOTE: We must state that $n \in \mathbb{Z}$.





Discussion: What does the n have to be a whole number?

If it isn't a whole number, we aren't doing a whole number of rotations which doesn't guarantee the same value for any trig functions.

Question 14

Find the general solutions to the following equation:

$$2 \sin\left(-2x + \frac{\pi}{4}\right) = \sqrt{2}$$

period = $\frac{2\pi}{2} = \pi$

$$\sin\left(-2x + \frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}$$

$$-2x + \frac{\pi}{4} = \frac{\pi}{4}, \frac{3\pi}{4}$$

$$-2x = 0, \frac{\pi}{2}$$

$$x = 0, -\frac{\pi}{4}$$

$x = \pi n, -\frac{\pi}{4} + \pi n$
 $n \in \mathbb{Z}$

Question 15 Walkthrough.

Find the general solutions to the following equation:

$$\tan\left(3x - \frac{\pi}{4}\right) + \frac{1}{\sqrt{3}} = 0$$

$$\tan\left(3x - \frac{\pi}{4}\right) = -\frac{1}{\sqrt{3}}$$

period = $\frac{\pi}{3}$

$$3x - \frac{\pi}{4} = -\frac{\pi}{6}$$

$$3x = -\frac{\pi}{6} + \frac{\pi}{4}$$

$$3x = \frac{-2\pi + 3\pi}{12}$$

$$3x = \frac{\pi}{12}$$

$$x = \frac{\pi}{36}$$

$\rightarrow x = \frac{\pi}{36} + \frac{\pi}{3} \cdot n, n \in \mathbb{Z}$

NOTE: For tangents, we always get one general solution!



Question 16

Find the general solutions to the following equation:

$$2\sqrt{3} + 2 \tan\left(2\left(x + \frac{\pi}{6}\right)\right) = 0$$

`:= Solve[2 $\sqrt{3}$ + 2 Tan[2 (x + π / 6)] == 0, x] // Expand`
 [풀이 함수] [탄젠트] [확장]


$$s] = \left\{ \left\{ x \rightarrow -\frac{\pi}{3} + \frac{\pi c_1}{2} \text{ if } c_1 \in \mathbb{Z} \right\} \right\}$$

Section D: Graphs of Sine and Cosine

Sub-Section: Basics of Sine and Cosine Functions


What does a Sine and Cosine graph look like?

Exploration: Graph of Sine and Cosine




➤ Label below $Q1, Q2, Q3, Q4$ for the section of the graph that corresponds to respective quadrants.

➤ Sin




➤ Cos



Amplitude, Period and Average Value



For $y = A \sin/\cos (nx + b) + k$



Consider the sign of our graph

$$\text{Amplitude} = |A|$$

$$\text{Period} = \frac{2\pi}{n}$$

$$\text{Average Value} = k$$

Sub-Section: Graphing Sine and Cosine Functions



Steps for Sketching Transformations of sin and cos Functions

➤ Identify:


Amplitude

Period

Mean Value

Positive/Negative Shape

➤ And create a "mini version" of the graph you are about to draw.




Consider the sign of our graph


➤ Start plotting the function from when the angle = 0.

🔄 For instance, for $\sin\left(2x - \frac{\pi}{3}\right)$, start from $x = \frac{\pi}{6}$.

🔄 Why?



➤ Draw the start and end of the periods, and plot the halves (turning points).




➤ Find any x -intercepts.

➤ Join all the points!

Question 17 Walkthrough.

Sketch the graph of $f(x) = -\sin(2x) + 1$ for $x \in [0, 2\pi]$ on the axes below, labelling all intercepts and endpoints with their coordinates.




Space for Personal Notes

Question 18

Sketch the following on the axes below, labelling all intercepts, endpoints, and turning points with their coordinates.

$$y = 2 \sin \left(2 \left(x - \frac{\pi}{3} \right) \right) - \sqrt{3} \text{ for } x \in [0, 2\pi]$$




Question 19

Sketch the following on the axes below, labelling all intercepts, endpoints, and turning points with their coordinates.

$$y = 2 \cos\left(2x + \frac{\pi}{3}\right) - 1 \text{ for } x \in [0, 2\pi]$$

$$= 2 \cos\left(2\left(x + \frac{\pi}{6}\right)\right) - 1$$

$\frac{\pi}{6}$ left




Section E: Graphs of Tangent

Sub-Section: Basics of Tangent Graphs


What does the tangent graph look like?

Exploration: Graph of Tangents



➤ Label below $Q1, Q2, Q3, Q4$ for the section of the graph which corresponds to respective quadrants.


➤ $\tan(x)$



Discussion: Why do we have a vertical asymptote for a tangent?

Because $\tan(90)$ is undefined

Sub-Section: Graphing Tangent Functions



Steps for Sketching tan Functions

➤ Identify

⚙ The period = $\frac{\pi}{n}$.

➤ Find the vertical asymptotes by solving for the angle = $\frac{\pi}{2}$.

➤ Find other vertical asymptotes within the domain by adding the period to answer from the previous step.


⚙ For instance, for $\tan\left(2x - \frac{\pi}{3}\right)$, solve $2x - \frac{\pi}{3} = \frac{\pi}{2}$ for x .

➤ Plot the inflection point (h, k) (Midpoint of the two vertical asymptotes).

⚙ x value of inflection point = x value, which makes an angle = 0.

⚙ y value of inflection point = vertical translation of the function.

eg: $\tan(x-h)+k$




➤ Find any x -intercepts.

➤ Sketch a "cubic-like" shape.

Question 20 Walkthrough.

Sketch the graph of $y = 3 \tan(2x)$ for $x \in [-\pi, \pi]$.




Space for Personal Notes

Your turn!

Question 21


Sketch the following on the axes below, labelling all intercepts, points of inflection, and endpoints with their coordinates, and all asymptotes with their equations.



Question 22

Sketch the following on the axes below, labelling all intercepts, points of inflection, and endpoints with their coordinates and all asymptotes with their equations.

$$f: [0, \pi] \rightarrow \mathbb{R}, f(x) = -3 \tan(\pi + 4x) + \sqrt{3}$$






Contour Checklist

- Learning Objective: [3.2.1] - Find Trig Ratios of Supplementary Relationships


Key Takeaways



$$\begin{aligned} \sin &= \frac{O}{H} \\ \cos &= \frac{A}{H} \\ \tan &= \frac{O}{A} \end{aligned}$$

Unit Circle

- The unit circle is simply a circle of radius 1.



$$\sin(\theta) = \underline{y}$$

$$\cos(\theta) = \underline{x}$$

$$\tan(\theta) = \underline{m}$$

- Period of a Trigonometric Function

period of $\sin(nx)$ and $\cos(nx)$ functions = $\frac{2\pi}{n}$

period of $\tan(nx)$ functions = $\frac{\pi}{n}$

where n = coefficient of x .

- Pythagorean identity:

$$\sin^2(\theta) + \cos^2(\theta) = 1$$

- Supplementary relationships:

\rightarrow ASTC

- Second Quadrant ($\pi - \theta$)

$$\cos(\pi - \theta) = -\cos(\theta)$$

$$\sin(\pi - \theta) = \sin(\theta)$$

$$\tan(\pi - \theta) = -\tan(\theta)$$

- Third Quadrant ($\pi + \theta$)

$$\cos(\pi + \theta) = -\cos(\theta)$$

$$\sin(\pi + \theta) = -\sin(\theta)$$

$$\tan(\pi + \theta) = \tan(\theta)$$

- Fourth Quadrant ($-\theta$)

$$\cos(-\theta) = \cos(\theta)$$

$$\sin(-\theta) = -\sin(\theta)$$

$$\tan(-\theta) = -\tan(\theta)$$

□ Learning Objective: [3.2.2] - Find Particular and General Solutions

Key Takeaways

○ Particular Solutions

- Solving trigonometric equations **for finite solutions.**

○ Steps:

- Make the trigonometric function the subject.
- Find the necessary (call) angle for one period.
- Solve for x by equating the necessary angles to the inside of the trigonometric functions.
- Add and subtract the period to find all other solutions in the domain.

□ General Solutions

- Finding infinite solutions to a trigonometric equation.

○ Steps:


- Make the trigonometric function the subject.
- Find the necessary angle for one period.
- Solve for x by equating the necessary angles to the inside of the trigonometric functions.
- Add period $\cdot n$ where $n \in \mathbb{Z}$.

Learning Objective: [3.2.3] - Graph Sine, Cosine and Tangent functions

Key Takeaways

Amplitude, Period and Average Value

For $y = A \sin / \cos (nx + b) + k$



Amplitude = $|A|$

Period = $\frac{2\pi}{n}$

Average Value = k

Tan function:

Period = $\frac{\pi}{n}$

- Find the asymptotes by solving for angle $(\) = \frac{\pi}{2}$.
- Find the other asymptotes by adding the period to the previous answer.

For the point of inflection:

x value of inflection point = x value, which makes an angle = $\frac{\pi}{2}$.

y value of inflection point = vertical translation of the function.



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

VCE Specialist Mathematics ½

Free 1-on-1 Consults



What Are 1-on-1 Consults?

- **Who Runs Them?** Experienced Contour tutors (45 + raw scores and 99 + ATARs).
- **Who Can Join?** Fully enrolled Contour students.
- **When Are They?** 30-minute 1-on-1 help sessions, after-school weekdays, and all-day weekends.
- **What To Do?** Join on time, ask questions, re-learn concepts, or extend yourself!
- **Price?** Completely free!
- **One Active Booking Per Subject:** Must attend your current consultation before scheduling the next. :)

SAVE THE LINK, AND MAKE THE MOST OF THIS (FREE) SERVICE!



Booking Link

bit.ly/contour-specialist-consult-2025

