

Website: contoureducation.com.au | Phone: 1800 888 300

Email: hello@contoureducation.com.au

# VCE Specialist Mathematics ½ Complex Numbers I [8.1]

Homework

#### **Admin Info & Homework Outline:**

| Student Name                |             |
|-----------------------------|-------------|
| Questions You Need Help For |             |
| Compulsory Questions        | Pg 2-Pg 23  |
| Supplementary Questions     | Pg 24-Pg 38 |




### **Section A:** Compulsory Questions



### Sub-Section [8.1.1]: Find Polar and Rectangular Forms of Complex Numbers

| Question | 1 |
|----------|---|
| Question | J |



|      | <br> |  |
|------|------|--|
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
| <br> | <br> |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |
|      |      |  |



| c. | Convert $-\sqrt{3} + i$ into polar form. |
|----|--|
|    |  |
|    |  |
|    |  |
| ٠  |  |
|    |  |
|    |  |
|    |  |

| Space for Personal Notes |  |
|--------------------------|--|
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |
|                          |  |




**Question 2** 



a.

i. Convert  $\operatorname{cis}\left(\frac{7\pi}{3}\right)$  into a rectangular form.

ii. Plot your answer on the Argand diagram below:






b.

i. Convert  $-2\sqrt{3} + 2i$  into polar form.

ii. Plot your answer on the Argand diagram below:






c.

i. Convert  $-\frac{\sqrt{6}}{2} - \frac{\sqrt{6}}{2}i$  into polar form.

\_\_\_\_\_

ii. Plot your answer on the Argand diagram below:






#### **Question 3**




a. Below is a complex number plotted on an Argand diagram. State the number in polar form.






**b.** The complex number z is on the Argand diagram below:




i. State z in polar form.

| - <u></u> | <br> |  |
|-----------|------|--|
|           |      |  |
|           |      |  |
|           | <br> |  |
|           |      |  |

ii. Plot  $z^2$  on the Argand diagram below:



**c.** 3 complex numbers, a, b and c are plotted on the Argand diagram below:




i. State which complex number corresponds to  $cis\left(\frac{23\pi}{8}\right)$ .

| <br> |  |
|------|--|
|      |  |
|      |  |
|      |  |
|      |  |
|      |  |

\_\_\_\_\_

ii. Plot  $b^2$  on the Argand diagram below:





| Qu        | Question 4 Tech-Active.  |  |  |  |  |
|-----------|--|--|--|--|--|
| a.        | Convert $\operatorname{cis}\left(\frac{15\pi}{8}\right)$ into a rectangular form.  |  |  |  |  |
|           |  |  |  |  |  |
|           |  |  |  |  |  |
|           |  |  |  |  |  |
|           | ·  |  |  |  |  |
|           |  |  |  |  |  |
|           | , (G, (G, (G_+, (G, (G_+, (G_+ |  |  |  |  |
| <b>b.</b> | Convert $\frac{\sqrt{6}-\sqrt{2}}{2} + \frac{\sqrt{6}+\sqrt{2}}{2}i$ into polar form.  |  |  |  |  |
|           |  |  |  |  |  |
|           |  |  |  |  |  |
|           |  |  |  |  |  |
|           |  |  |  |  |  |
|           |  |  |  |  |  |
|           |  |  |  |  |  |





### <u>Sub-Section [8.1.2]</u>: Evaluate Operations of Complex Numbers

| Qu | Question 5   |  |  |  |
|----|--|--|--|--|
| a. | Let $u = 5\operatorname{cis}\left(\frac{3\pi}{8}\right)$ and $v = 2\operatorname{cis}\left(\frac{2\pi}{8}\right)$ . Evaluate $uv$ , leaving your answer in polar form.           |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
| b. | Let $u = 3\operatorname{cis}\left(\frac{5\pi}{6}\right)$ and $v = 2\operatorname{cis}\left(\frac{5\pi}{12}\right)$ . Evaluate $\frac{u}{v}$ , leaving your answer in polar form. |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |
|    |  |  |  |  |

| c. | Let $u = 4 + \frac{7}{2}i$ and $v = \frac{3}{5} + i$ . Evaluate $u + v$ . |
|----|---|
|    |   |
|    |   |
|    |   |
|    |   |
|    |   |



**Question 6** 



**a.** Let u=1+3i and  $v=\sqrt{3}+\sqrt{3}i$ . Evaluate uv, leaving your answer in rectangular form.

**b.** Simplify  $\frac{1+2i}{1-i}$ .

c. Let  $u = \operatorname{cis}\left(\frac{3\pi}{4}\right)$  and  $v = 2\operatorname{cis}\left(\frac{2\pi}{3}\right)$ . Evaluate u + v, leaving your answer in rectangular form.

**Question 7** 



**a.** Let  $u = \frac{\sqrt{\sqrt{2}+2}}{2} + \frac{\sqrt{2-\sqrt{2}}}{2}i$ . Evaluate  $u^2$ , leaving your answer in polar form.

**b.** Simplify  $\frac{-1+\sqrt{3}i}{2-\sqrt{3}i}$ .

c. Let  $u = \sqrt{2}\operatorname{cis}\left(\frac{\pi}{4}\right)$  and  $v = 1 + (1 - 2\sqrt{3})i$ . Evaluate  $u + \bar{v}$ , leaving your answer in polar form.



Question 8 Tech-Active.

**a.** Evaluate  $\operatorname{cis}\left(\frac{5\pi}{7}\right) + \operatorname{cis}\left(\frac{2\pi}{7}\right)$ , leaving your answer in rectangular form to 2 decimal places.

**b.** Let  $u = 5\sqrt{3} - \sqrt{2}i$  and  $v = \sqrt{7} - i$ . Simplify  $\frac{u}{v}$ .





### Sub-Section [8.1.3]: Apply De Moivre's Theorem

| Question 9 |   |  |  |
|------------|---|--|--|
| a.         | Let $a = \sqrt{2}\operatorname{cis}\left(\frac{\pi}{6}\right)$ . Evaluate $a^4$ , leaving your answer in rectangular form.                        |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |
| b.         | Let $v = \sqrt{3}\operatorname{cis}\left(\frac{5\pi}{7}\right)$ . Evaluate $v^3$ , leaving your answer in polar form with the principal argument. |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |
|            |   |  |  |



| C. | Evaluate $(1+i)^5$ , leaving your answer in rectangular form. |  |
|----|---|--|
|    | Evaluate (1 + t) , leaving your answer in rectangular form.   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    | 6 B 1N .  |  |
| Sp | ace for Personal Notes  |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
|    |   |  |
| Ī  |   |  |

**Question 10** 



**a.** Let  $u = -\frac{\sqrt{3}}{2} - \frac{1}{2}i$ . Evaluate  $u^3$ , leaving your answer in rectangular form.

**b.** Let  $v = 2^{\frac{1}{11}} \operatorname{cis}\left(\frac{\pi}{6}\right)$ . Evaluate  $v^{22}$ , leaving your answer in rectangular form.

**c.** Evaluate  $\frac{\left(1+\sqrt{3}i\right)^3}{\left(-\sqrt{2}+\sqrt{2}i\right)^4}$ , leaving your answer in rectangular form.

#### **Question 11**



**a.** Find the value(s) of n such that  $\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right)^n = \operatorname{cis}\left(\frac{\pi}{3}\right)$ .

**b.** Suppose  $u = \operatorname{cis}\left(\frac{2\pi}{7}\right)$ . Find the value(s) of n such that  $u^n$  is a real number.

**c.** Find the value(s) of n such that  $(3+3i)^n = ai$ , where  $a \in R$  and a < 0.



| Question 12 Tech-Active.   |  |
|--|--|
| Evaluate $\left(\sqrt{2}\operatorname{cis}\left(\frac{\pi}{12}\right)\right)^5$ , leaving your answer in rectangular form. |  |
|  |  |
|  |  |
|  |  |
|  |  |
|  |  |
|  |  |
|  |  |

| Space for Personal Notes |  |  |
|--------------------------|--|--|
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |
|                          |  |  |





### **Sub-Section**: Final Boss

| Question 13   |  |  |
|---|--|--|
| Consider the equation $z^6 - 1 = 0$ , $z \in C$ .   |  |  |
| <b>a.</b> Use De Moivre's theorem to verify that $z = \operatorname{cis}\left(\frac{\pi}{3}\right)$ is a solution to this equation. |  |  |
|   |  |  |
|   |  |  |
|   |  |  |
|   |  |  |
|   |  |  |
|   |  |  |
| <b>b.</b> Find the other five solutions to the equation.  |  |  |
|   |  |  |
|   |  |  |
|   |  |  |
|   |  |  |
|   |  |  |
|   |  |  |



| c. | Show that $cis(a) + cis(-a) = 2 cos(a)$ .                |
|----|--|
|    |  |
|    |  |
|    |  |
|    |  |
| d. | Hence, show that the sum of the roots of $z^6 - 1$ is 0. |
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |



| Qı   | Question 14   |  |  |
|------|---|--|--|
| De   | De Moivre's theorem states that $(\cos(\theta) + i\sin(\theta))^n = \cos(n\theta) + i\sin(n\theta)$ . |  |  |
| a.   | Show that De Moivre's theorem is true for $n = 2$ .   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
| b.   | Using induction, prove De Moivre's theorem for $n \in \mathbb{N}$ .                                   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
|      |   |  |  |
| ۲    | pace for Personal Notes   |  |  |
| <br> | Jace IVI I Elsvilai NULES   |  |  |
|      |   |  |  |



### Section B: Supplementary Questions



### Sub-Section [8.1.1]: Find Polar and Rectangular Forms of Complex Numbers

| Qu | uestion 15                                 |
|----|--|
| a. | Write $14i$ in polar form.                 |
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |
| b. | Convert $-2 - 2\sqrt{3}i$ into polar form. |
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |



| c. | Convert $\sqrt{2}\operatorname{cis}\left(\frac{-\pi}{4}\right)$ into a rectangular form. |
|----|--|
|    |  |
|    |  |
|    |  |
|    |  |

**Question 16** 



**a.** Convert  $-\frac{3\sqrt{3}}{2} + \frac{3}{2}i$  into polar form.

**b.** Convert  $\operatorname{cis}\left(\frac{420\pi}{4}\right)$  into Cartesian form.


**c.** Convert  $-\frac{5\sqrt{2}}{2} - \frac{5\sqrt{2}}{2}i$  into polar form.



#### **Question 17**



The complex number  $z = \sqrt{2}\operatorname{cis}\left(\frac{\pi}{6}\right)$  is shown in the Argand diagram below:



On the same set of axes, plot:

- **a.**  $z^2$
- **b.**  $z^4$
- c.  $\bar{z}$
- **d.**  $z\bar{z}$



#### **Question 18**



**a.** Use a compound formula to show that  $\sin\left(\frac{7\pi}{12}\right) = \frac{\sqrt{6}+\sqrt{2}}{4}$ .

**b.** Hence, show that  $\cos\left(\frac{7\pi}{12}\right) = \frac{\sqrt{2}-\sqrt{6}}{4}$ , using the fact that  $\sqrt{\frac{2-\sqrt{3}}{4}} = \frac{\sqrt{6}-\sqrt{2}}{4}$ .

c. Hence, state  $\operatorname{cis}\left(\frac{17\pi}{12}\right)$ .






### <u>Sub-Section [8.1.2]</u>: Evaluate Operations of Complex Numbers

**Question 19** 



**a.** Let  $u = 3\operatorname{cis}\left(\frac{\pi}{5}\right)$  and  $v = 2\operatorname{cis}\left(\frac{-4\pi}{5}\right)$ . Evaluate uv, leaving your answer in polar form.

**b.** Let  $u = \operatorname{cis}\left(\frac{-3\pi}{4}\right)$  and  $v = \operatorname{cis}\left(\frac{\pi}{4}\right)$ . Evaluate  $\frac{u}{v}$ .



| c. | Let $u = \sqrt{3} + 2i$ and $v = \sqrt{48} - i$ . Evaluate $u + v$ . |
|----|--|
|    |  |
|    |  |
|    |  |
|    |  |
|    |  |

| Space for Personal Notes |
|--------------------------|
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |
|                          |

| Question | 20 |
|----------|----|
| Question | _  |



**a.** Find  $\frac{1-i}{2+3i}$ .

**b.** Let  $a = (3 - \sqrt{2}i)$  and  $b = (2 + \sqrt{5}i)$ . Evaluate ab, leaving your answer in rectangular form.

c. Evaluate  $2\operatorname{cis}\left(\frac{3\pi}{4}\right) + \operatorname{cis}\left(\frac{\pi}{3}\right)$ , leaving your answer in an appropriate form.



| uestion 21  |  |
|---|--|
| Show that $z\bar{z} =  z ^2$ for any complex number z.          |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
| Evaluate $\frac{3+10i}{2-5i}$ .                                 |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
| It is known that $z = 1 - 2i$ . Find $z^4$ in rectangular form. |  |
| It is known that $z = 1 - 2t$ . Find $z = 1$ rectangular form.  |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |
|   |  |



| Question | 22 |
|----------|----|
|----------|----|



| u | RESTIVIT 22  |   |
|---|--|---|
|   | Show that for any complex number $z = a + bi$ where $a, b \in \mathbb{R}$ , $\text{Im}(z^3) = b(3a^2 - b^2)$ . |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   | Hence, prove or disprove the statement $z^3 \in \mathbb{R}$ if and only if $z \in \mathbb{R}$ .                |   |
|   | Hence, prove or disprove the statement $z^3 \in \mathbb{R}$ if and only if $z \in \mathbb{R}$ .                | _ |
|   | Hence, prove or disprove the statement $z^3 \in \mathbb{R}$ if and only if $z \in \mathbb{R}$ .                |   |
|   | Hence, prove or disprove the statement $z^3 \in \mathbb{R}$ if and only if $z \in \mathbb{R}$ .                |   |
|   |  |   |
|   |  |   |
|   |  |   |
|   | Hence, prove or disprove the statement $z^3 \in \mathbb{R}$ if and only if $z \in \mathbb{R}$ .                |   |





### Sub-Section [8.1.3]: Apply De Moivre's Theorem

| Qu  | Question 23   |  |  |  |  |
|-----|---|--|--|--|--|
| Eva | aluate the following, leaving your answer in an appropriate form. |  |  |  |  |
| a.  | $\left(3\operatorname{cis}\left(\frac{\pi}{6}\right)\right)^3$    |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
| b.  | $(-2+2i)^3$   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |
|     |   |  |  |  |  |

 $\mathbf{c.} \quad \left(2\operatorname{cis}\left(\frac{-\pi}{10}\right)\right)^2$ 

**Question 24** 



**a.** Evaluate  $(1 + \sqrt{3}i)^{48}$ , leaving your answer in the form  $a^b$  where a and b are integers.

| b.          |            |  |
|-------------|------------|--|
| j           | i <b>.</b> | Convert $2\sqrt{2} - 2\sqrt{2}i$ to polar form.  |
|             |            |  |
|             |            |  |
|             |            |  |
|             |            |  |
|             |            |  |
| i           | ii.        | Hence, evaluate $\frac{(2\sqrt{2}-2\sqrt{2}i)^7}{(-1+\sqrt{3}i)^{12}}$ . State your answer in the form $a+bi$ , where $a,b\in\mathbb{R}$ . |
|             |            |  |
|             |            |  |
|             |            |  |
|             |            |  |
|             |            |  |
| <b>c.</b> ] | Eva        | aluate $\left(\frac{\sqrt{2}+\sqrt{2}i}{\sqrt{3}+1}\right)^5$ , leaving your answer in polar form.   |
| -           |            |  |
| -           |            | ·  |
| -           |            |  |
| -           |            |  |
| -           |            |  |



| $\mathbf{O}$ | uestion | 25 |
|--------------|---------|----|
| ~            | acoulon |    |



| a. | Find the value of $n$ such that $(\sqrt{2} + \sqrt{2}i)^n = 1024 \operatorname{cis}(\theta)$ where $\theta \in (-\pi, \pi]$ , and state $\theta$ . |
|----|--|
|    |  |

**b.** Solve for the value(s) of n such that z is purely imaginary.

$$z = 9\operatorname{cis}\left(\frac{\pi}{10}n\right)$$

| _ |      |      |
|---|------|------|
|   |      |      |
| _ | <br> | <br> |
|   |      |      |
|   |      |      |
|   |      |      |
|   |      |      |
|   |      |      |
|   |      |      |
|   |      |      |
|   |      |      |
|   |      |      |

| c. | Let $z = \left(6\operatorname{cis}\left(\frac{\pi}{7}\right)\right)^n$ . Solve for the values of $n$ such that $z$ is a real number. |
|----|--|
|    |  |
|    |  |
|    |  |
|    |  |

#### **Question 26**



Solve for the value(s) of n such that z is a real number.

$$z = 4\operatorname{cis}\left(\frac{3(\pi+2)}{8}n\right)$$

| <br> |  |
|------|--|
|      |  |
|      |  |
| <br> |  |
|      |  |
|      |  |



Website: contoureducation.com.au | Phone: 1800 888 300 | Email: hello@contoureducation.com.au

### VCE Specialist Mathematics ½

## Free 1-on-1 Consults

#### What Are 1-on-1 Consults?

- **Who Runs Them?** Experienced Contour tutors (45 + raw scores and 99 + ATARs).
- Who Can Join? Fully enrolled Contour students.
- When Are They? 30-minute 1-on-1 help sessions, after-school weekdays, and all-day weekends.
- What To Do? Join on time, ask questions, re-learn concepts, or extend yourself!
- Price? Completely free!
- One Active Booking Per Subject: Must attend your current consultation before scheduling the next. :)

SAVE THE LINK, AND MAKE THE MOST OF THIS (FREE) SERVICE!

# G

### **Booking Link**

bit.ly/contour-specialist-consult-2025

