

2007

VCE Further Mathematics Trial Examination 1

Suggested Solutions

© Kilbaha Multimedia Publishing 2007



Kilbaha Multimedia Publishing ABN 47 065 111 373
PO Box 2227
Kew Vic 3101
Australia
Tel: 03 9817 5374
Fax: 03 9817 4334
chemas@chemas.com
www.chemas.com

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Multimedia Publishing.
- The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
- For authorised copying within Australia please check that your institution has a licence from **Copyright Agency Limited**. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.

Reproduction and communication for educational purposes The Australian Copyright Act 1968 (the Act) allows a maximum of one chapter or 10% of the pages of this work, to be reproduced and/or communicated by any educational institution for its educational purposes provided that educational institution (or the body that administers it) has given a remuneration notice to Copyright Agency Limited (CAL) under the Act.

For details of the CAL licence for educational institutions contact
CAL, Level 15, 233 Castlereagh Street, Sydney, NSW, 2000
Tel: (02) 9394 7600
Fax: (02) 9394 7601
Email: info@copyright.com.au


- Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.
- Schools which purchase a licence to use this material may distribute this electronic file to the students at the school for their exclusive use. This distribution can be done either on an Intranet Server or on media for the use on stand-alone computers.
- Schools which purchase a licence to use this material may distribute this printed file to the students at the school for their exclusive use.
- **The Word file (if supplied) is for use ONLY within the school**
- **It may be modified to suit the school syllabus and for teaching purposes.**
- **All modified versions of the file must carry this copyright notice**
- **Commercial use of this material is expressly prohibited**

Question 1 A $\frac{61}{73+61+146} \times \frac{100}{1} = \frac{61}{280} \times \frac{100}{1} = 21.786\%$	Question 2 D Having or not having a water tank or having one on order is categorical data. Scatter plots and histograms are used when both variables are numerical. For a stem and leaf you need two categories, but there are three categories here. You cannot find the median, lower quartile, etc. so cannot draw a box plot. Bar graphs are used for categorical data.
Question 3 C A long whisker to the left indicates the box plot is negatively skewed. The interquartile range = $80 - 60 = 20$	Question 4 C 99.7% lie within 3 standard deviations of the mean, so 0.3% lie outside this range. Half of $0.3\% = 0.15\%$. this means 0.15% lie above 3 standard deviations above the mean. $43 + 3 \times 15 = 88$. So the lowest mark for an A is 88
Question 5 A The standard deviation remains at 5 g., because the spread is unaltered. The mean weight of each bag is increased by 2g to 452g.	Question 6 B $r^2 = 0.1296$ $r = \pm\sqrt{0.1296}$ $r = \pm 0.36$ But the gradient of the line is negative, (-0.73) so the correlation coefficient must be negative $\therefore r = -0.36$
Question 7 B Draw 2 vertical lines so that there are 3 points to the left of the first vertical line, 4 points in the middle, and 3 points to the right of the last vertical line. $x_l = 2, y_l = 2$ $x_u = 10, y_u = 18$ gradient = $\frac{18-2}{10-2} = \frac{16}{8} = 2$	Question 8 E The gradient is approximately -28. This means that for every dollar increase in fares, the number of passengers decreased by 28.
Question 9 D For y versus x the graph would look like A or C, not B. However, the graph would contain the point $(1,2)$. Hence, not A, B, or C. For y versus $\log x$ the graph would look like D or E but would contain the point $(1,5)$. Hence, D	


Question 10 B Enter data in graphics calculator in L_1 and L_2 using Stat Edit Press Stat Calc Linear Regression Line L_1, L_2 This gives $r^2 = 0.305$	Question 11 D To use seasonal indices, there must be a secular trend and the variation must be seasonal.
Question 12 D Quarterly average = $\frac{240 + 226 + 210 + 120}{4} = 199$ Seasonal Index for 3rd quarter = $\frac{210}{199} = 1.055$	Question 13 A The median of the first three points is 500 so the point (1985,500) should lie on the graph, hence C, D or E. The median of points 3,4, and 5 is 500, so the third point on the smoothed graph should be (1995,500). Hence E

Question 1 D Each term in the sequence is found by adding 6 to the term before it. Hence it is an arithmetic sequence with a common difference of 6	Question 2 B This is a geometric sequence with a common ratio of $-\frac{1}{2}$ $a = 8, n = 12$ $t_n = ar^{n-1}$ $t_n = 8\left(-\frac{1}{2}\right)^{11} = -\frac{1}{256}$
Question 3 C $t_7 = 3 \times t_6 + 4$ $8017 = 3 \times t_6 + 4$ $8013 = 3 \times t_6$ $2671 = t_6$ $t_6 = 3 \times t_5 + 4$ $2671 = 3 \times t_5 + 4$ $2667 = 3 \times t_5$ $889 = t_5$	Question 4 C $t_4 = t_{1+3} = t_{1+2} + t_{1+1} + t_1 = t_3 + t_2 + t_1 = 1 + 1 + 1 = 3$ $t_5 = t_{2+3} = t_{2+2} + t_{2+1} + t_2 = t_4 + t_3 + t_2 = 3 + 1 + 1 = 5$
Question 5 C $4 + 6 + 8 + \dots$ Sum of Arithmetic Sequence $S_n = \frac{n}{2}[2a + (n-1)d]$ $S_{10} = \frac{10}{2}[8 + 9 \times 2] = 5 \times 26 = 130\text{m}$	Question 6 E Value at beginning of first year = 32,000 Value at beginning of second year = $32,000(0.85)$ Value at beginning of third year = $32,000(0.85)^2$ Following the pattern, Value at beginning of fifth year = $32,000(0.85)^4$ $= \\$16704.20$ This is closest to \$16704

Question 7 D Arithmetic Sequence. $a = 85$ $d = -7$ $t_n = a + (n - 1)d = 85 + (n - 1)(-7)$ Use sequence mode on graphics calculator and press $y =$ $n(min) = 1$ $\mu(n) = 85 + (n - 1)(-7)$ $\mu(n \text{ min}) = 85$ Press second table and scroll down to get -48 This corresponds to $n = 20$	Question 8 A $\frac{72}{x} = \frac{x}{8}$ $x^2 = 576$ $x = \pm\sqrt{576} = \pm 24$ Common ratio = $\frac{\pm 24}{8} = \pm 3$
Question 9 D $S_\infty = \frac{a}{1-r} = \frac{2}{1-\frac{2}{5}} = 3.3333$ $S_n = \frac{a(1-r^n)}{1-r} = \frac{2(1-0.4^n)}{0.6}$ Enter this equation in sequence mode in the graphics calculator $n(min) = 1$ $\mu(n \text{ min}) = 2$ Using the table scroll down $n = 5 \quad \mu(n) = 3.2992$ $n = 6 \quad \mu(n) = 3.3197$ $n = 7 \quad \mu(n) = 3.3279$ $3.3333 - 3.2992 = 0.0341$ which is too big $3.3333 - 3.3197 = 0.0136$ which is too big $3.3333 - 3.3279 = 0.0054$ which is less than 0.01	

Question 1 E $b^2 = a^2 + c^2 - 2ac \cos 50^\circ$ $b^2 = 27^2 + 41^2 - 2 \times 27 \times 41 \times \cos 50^\circ$ $b = \sqrt{27^2 + 41^2 - 2 \times 27 \times 41 \times \cos 50^\circ}$ $b = 31.4$ AC is closest to 31cm	Question 2 D $\text{Area} = \frac{1}{2} ac \sin b$ $\text{Area} = \frac{1}{2} \times 4 \times 7 \times \sin 64^\circ$ $\text{Area} = 12.58 \text{ cm}^2$
Question 3 E In this right angled isosceles triangle, AB = AC, so $\angle BAC = \theta$. The sum of the angles of a triangle = 180° $\therefore \theta = 45^\circ$ $\sin 45^\circ = \cos 45^\circ = \frac{1}{\sqrt{2}} = 0.707$	Question 4 A  $\text{gradient} = \frac{100}{3000} = 0.033$ which is closest to 0.03
Question 5 C In $\triangle ADB$, $AD^2 = 20^2 - 16^2$ $AD = \sqrt{20^2 - 16^2} = 12$ In $\triangle BDC$, $DC^2 = 34^2 - 16^2$ $DC = \sqrt{34^2 - 16^2} = 30$ $AC = AD + DC = 12 + 30 = 42 \text{ cm}$	Question 6 E Let the length of the original cube = x Then the original volume = x^3 Length of larger cube = $4x$ Volume of larger cube = $(4x)^3 = 64x^3$ $\therefore$ volume has increased by a factor of 64

Question 7 A



$$\angle NXZ = \angle XZQ \text{ (alternate angles)} = 142^\circ$$

$$\angle NZQ = 180^\circ$$

$$\angle XZY = 340 - (180 + 142) = 18^\circ$$


Question 8 C

$\Delta s ABO$ and CDO are similar (AAA)

$$\therefore \frac{x}{5.2} = \frac{2.5}{10}$$

$$x = \frac{2.5 \times 5.2}{10} = 1.3 \text{ m} = 130 \text{ cm}$$

Question 9 A



$$\tan 30^\circ = \frac{30}{DC}$$

$$DC = \frac{30}{\tan 30^\circ} = 51.9615 = AB$$

$$AC^2 = AB^2 + 18^2$$

$$AC^2 = 51.9615^2 + 18^2$$

$$AC = \sqrt{51.9615^2 + 18^2}$$

$$AC = 54.99$$

$$\tan \theta = \frac{30}{AC} = \frac{30}{54.99}$$

$$\theta = \tan^{-1}\left(\frac{30}{54.99}\right) = 28.6^\circ \text{ which is closest to } 29^\circ$$

Question 1 C They both meet where the lines meet. Read the distance from the distance axis. This is 15 km.	Question 2 E Jake travels $25 + 25 = 50$ km Jenny travels $5 + 15 = 20$ km. So Jake travels 30 km further than Jenny
Question 3 E $28 = a \times 2^3$ $28 = a \times 8$ $a = \frac{28}{8} = 3.5$	Question 4 D $2 \times 3 - 5 \times 2 + b = 0$ $6 - 10 + b = 0$ $b = 10 - 6 = 4$
Question 5 B The charge per hour is the gradient of the line $= \frac{90 - 30}{3 - 0} = \frac{60}{3} = 20$ The fixed cost is the y intercept = 30	Question 6 D The shaded region is above the line or equal to the line $y = x - 3$ $\therefore y \geq x - 3$ $\therefore$ not C or E The vertical line is $x = 5$ and the required region is greater than or equal to this line $\therefore x \geq 5$, so not B The horizontal line is $y = 4$ and the required region is less than or equal to this line so $y \leq 4$ $\therefore$ D

Question 7 B $3y = -6x + 2$ $y = -2x + \frac{2}{3}$ gradient of this line is -2 $\therefore$ gradient of parallel line is -2 $y = -2x + c$ when $x = 4, y = 10$ $10 = -2 \times 4 + c$ $10 = -8 + c$ $c = 18$ $\therefore y = -2x + 18$ $y + 2x - 18 = 0$	Question 8 E If you multiplied equation 1 by 2 and equation 2 by 3 you would have $-6b$ in each equation. The b could be eliminated by subtraction. Question 9 A Eqn of revenue $R = mx + c$ Points on revenue line are (0,0) and (200,600) $m = \frac{600 - 0}{200 - 0} = 3$ $y = 3x + c$ c is the y intercept = 0 $R = 3x$ Eqn of cost $C = mx + c$ Points on cost line are (0,400) and (200,600) $m = \frac{600 - 400}{200 - 0} = \frac{200}{200} = 1$ $c = 400$ $C = x + 400$ When $x = 150, C = 150 + 400 = 550$ When $x = 150, R = 3 \times 150 = 450$ Therefore, loss = 100
--	---

Question 1 C $R = \frac{100I}{PT} = \frac{100 \times 945}{4800 \times 3} = 6.5625$ This is closest to 6.6%	Question 2 D Amount paid = $7000 + 24 \times 1330 = \\$38,920$ Interest = Amount paid – Cost Interest = $38920 - 36000 = \\$2,920$
Question 3 B Sale price = 85% of 1300 = 1105 Price paid by customer = 88% of 1105 = \$972.40	Question 4 C Use TVM solver under finance on a graphics calculator. $N = 48$ $I = 10$ $Pv = -3000$ $PMT = 0$ $FV =$ $P / Y = 4$ $C / Y = 4$ This gives $FV = 9814.47$ Interest = Final value – Amount invested Interest = $9814.47 - 3000 = 6814.47$ This is closest to \$6814
Question 5 C On 9 Jan. have $7648 + 2004 = 9652$ On 29 Jan have 7632 Amt. taken out of account = $9652 - 7632$ = \$2020	
Question 6 B Interest will be paid for the months of Dec., Jan. and Feb. Dec. min monthly balance = 9243 Jan. min monthly balance = 7632 Feb. min monthly balance = 7532 Interest for Dec. = $9243 \times \frac{1}{100} \times 1 = 92.43$ Interest for Jan. = $7632 \times \frac{1}{100} \times 1 = 76.32$ Interest for Feb. = $7532 \times \frac{1}{100} \times 1 = 75.32$ Total Interest = $92.43 + 76.32 + 75.32 = \\244.07	Question 7 E $(0.92)^n \times 2000 = 800$ Enter $y = (0.92)^n \times 2000$ in graphics calculator Then 2nd Table When $x = 11$ $y = 799.27$ So 11 years.

Question 8 E

Compound interest rate increases more rapidly with time. This means the gradient of the graph increases with time. The only graph that does this is E

Question 9 C

Use TVM solver under finance on a graphics calculator.

$$N = 48$$

$$I = 9.2$$

$$PV = 20000$$

$$PMT = -300$$

$$FV =$$

$$P / Y = 12$$

$$C / Y = 12$$

This gives $FV = 11,395.90$

Paid back lump sum of 2000 leaves

9395.90 owing

$$N =$$

$$I = 9.2$$

$$PV = 9395.90$$

$$PMT = -400$$

$$FV = 0$$

$$P / Y = 12$$

$$C / Y = 12$$

This gives $N = 25.8$ months which is closest to 2 years.

Question 1 A A complete graph has each vertex linked to every other vertex	Question 2 E For an Euler circuit to exist, each vertex must be of even degree.
Question 3 E A to A is 0 B to B is 0 C to C is 1 D to D is 0 Hence, the diagonal is 0,0,1,0	Question 4 C A tree is a connected graph with no loops, circuits or multiple edges. A, B, D, and E have loops, multiple edges or circuits.
Question 5 E Degree of A = 3 Degree of B = 4 Degree of C = 5 Degree of D = 3 Degree of E = 5 Degree of F = 4 Need a line joining any two of A, C, D or E, so that two of the vertices only remain odd. From given choices it has to be C to D	Question 6 D Cameron is the parent of Donald and Anthea, not of Chris

Question 7 B

F is a prerequisite for D so not C
C is a prerequisite for D so not A or D
D is a prerequisite for E so not E

Question 8 B

Ben has dominance over Carrie who in turn has dominance over David.

Question 9 B

B,C,D,G lie on the critical path because their float time is zero. Hence, A is not true.
If all the crash times are implemented, only those on the critical path will reduce the overall time of the project. Hence, C is not true. E does not lie on the critical path, so D is not true. The total time for the project is the length of the critical path so E is not true. E does not lie on the critical path, so it does not affect the overall time of the project.

Question 1 A $\det A = (4 \times -3) - (-2 \times -1) = -12 - 2 = -14$	Question 2 D $2A = \begin{bmatrix} 2a & 2b \\ 2c & 2d \end{bmatrix}$ $2A - B = \begin{bmatrix} 2a - e & 2b - f \\ 2c - g & 2d + h \end{bmatrix}$
Question 3 B AB does not exist because the number of columns of A does not equal the number of rows of B. BA is a 4×3 matrix. Order of BA is number of rows of B $\times$ number of columns of A	Question 4 C A has no solutions because the lines are parallel. B has one, unique, solution. (2,4) D has a unique solution because $\det D = 3 - 2 = 1$ E has a unique solution because $\det E = -12 - 12 = -24$ C has many solutions because the two lines are the same.
Question 5 A This can be done on the graphics calculator or $A - B = \begin{bmatrix} 4 & 1 \\ 2 & 3 \end{bmatrix} - \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 2 & 2 \end{bmatrix}$ $C^2 = \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix} \times \begin{bmatrix} 1 & 2 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} 7 & 4 \\ 6 & 7 \end{bmatrix}$ $(A - B)C^2 = \begin{bmatrix} 2 & 1 \\ 2 & 2 \end{bmatrix} \times \begin{bmatrix} 7 & 4 \\ 6 & 7 \end{bmatrix} = \begin{bmatrix} 20 & 15 \\ 26 & 22 \end{bmatrix}$	Question 6 E From Y to X is 0.25 so the first row second column is 0.25 so not B or C or D From X to Z is 0.05 so not A Hence, answer E

Question 7 A S_0 is the initial number of people choosing milk and dark chocolate. $S_0 = \begin{bmatrix} 2180 \\ 1260 \end{bmatrix}$	Question 8 B $T = \begin{bmatrix} 0.72 & 0.35 \\ 0.28 & 0.65 \end{bmatrix}$ $T^5 S_0 = \begin{bmatrix} 0.72 & 0.35 \\ 0.28 & 0.65 \end{bmatrix}^{5 \times} \begin{bmatrix} 2180 \\ 1260 \end{bmatrix} = \begin{bmatrix} 1913 \\ 1527 \end{bmatrix},$ using graphics calculator Number choosing dark chocolate = 1527																				
Question 9 D <table><tr><td></td><td>D</td><td>N</td><td>C</td><td></td></tr><tr><td>X</td><td>50</td><td>60</td><td>200</td><td>$\begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$</td></tr><tr><td>Y</td><td>20</td><td>100</td><td>30</td><td></td></tr><tr><td>Z</td><td>80</td><td>40</td><td>70</td><td></td></tr></table> $\begin{bmatrix} 100 + 60 + 600 \\ 40 + 100 + 90 \\ 160 + 40 + 210 \end{bmatrix} = \begin{bmatrix} 760 \\ 230 \\ 410 \end{bmatrix}$			D	N	C		X	50	60	200	$\begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$	Y	20	100	30		Z	80	40	70	
	D	N	C																		
X	50	60	200	$\begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$																	
Y	20	100	30																		
Z	80	40	70																		

End of suggested solutions
2007 Further Mathematics Trial Examination 1

KILBAHA MULTIMEDIA PUBLISHING PO BOX 2227 KEW VIC 3101 AUSTRALIA	TEL: (03) 9817 5374 FAX: (03) 9817 4334 chemas@chemas.com www.chemas.com
--	---