



THE SCHOOL FOR EXCELLENCE
UNIT 4 MATHEMATICAL METHODS 2006
COMPLIMENTARY WRITTEN EXAMINATION 1 - SOLUTIONS

QUESTION 1

- a. (i) Divide $3x+1$ into $3x^4 + 4x^3 - 14x^2 - 20x - 5$ using either synthetic or polynomial long division and get a remainder of zero:

$$\begin{array}{r}
 \overline{x^3 + x^2 - 5x - 5} \\
 3x+1 \overline{) 3x^4 + 4x^3 - 14x^2 - 20x - 5} \\
 \underline{3x^4 + x^3} \\
 3x^3 - 14x^2 - 20x - 5 \\
 \underline{3x^3 + x^2} \\
 -15x^2 - 20x - 5 \\
 \underline{-15x^2 - 5x} \\
 -15x - 5 \\
 \underline{-15x - 5} \\
 0
 \end{array}$$

- (ii) It follows from (i) that:

$$p(x) = 3x^4 + 4x^3 - 14x^2 - 20x - 5 = (3x+1)(x^3 + x^2 - 5x - 5).$$

- b. (i) Using trial and error it is readily found that $q(-1) = 0$.
It follows that $x - (-1) = x+1$ is a factor of $q(x)$.
- (ii) Divide $x+1$ into $q(x) = x^3 + x^2 - 5x - 5$ using either synthetic or polynomial long division to get:

$$q(x) = x^3 + x^2 - 5x - 5 = (x+1)(x^2 - 5)$$

Use difference of two squares formula:

$$= (x+1)(x - \sqrt{5})(x + \sqrt{5}).$$

$$\text{Therefore: } p(x) = (3x+1)q(x) = (3x+1)(x+1)(x - \sqrt{5})(x + \sqrt{5}).$$

QUESTION 2

a. By definition: $\int_0^b ax \, dx = 1.$

Therefore: $\frac{a}{2} [x^2]_0^b = 1$

$$\therefore \frac{a}{2} b^2 = 1 \Rightarrow ab^2 = 2.$$

b. By definition: $\overline{X} = \int_0^b (x)ax \, dx = a \int_0^b x^2 \, dx.$

Therefore: $\overline{X} = \frac{a}{3} [x^3]_0^b = \frac{a}{3} b^3.$

c. Using the result from part (b): $\frac{a}{3} b^3 = 1 \therefore ab^3 = 3. \quad (1)$

From part (a): $ab^2 = 2. \quad (2)$

Equation (1)/Equation (2): $b = \frac{3}{2}.$

Substitute $b = \frac{3}{2}$ into equation (2): $a \left(\frac{3}{2} \right)^2 = 2 \Rightarrow a = \frac{8}{9}.$

QUESTION 3

Substitute $w = 2^x$. Then: $2^{3x} + 2^{2x} - 2^{-x} = 1 \Leftrightarrow (2^x)^3 + (2^x)^2 - (2^x)^{-1} = 1$
 $\therefore w^3 + w^2 - w^{-1} = 1$

Multiply both sides by w : $\therefore w^4 + w^3 - 1 = w \Leftrightarrow w^4 + w^3 - w - 1 = 0$
 $(w^4 - 1) + (w^3 - w) = 0$
 $(w^2 - 1)(w^2 + 1) + w(w^2 - 1) = 0$
 $(w^2 - 1)(w^2 + w + 1) = 0.$

Use the null factor theorem: $w^2 + w + 1 = 0$ or $w^2 - 1 = 0.$
 $w^2 + w + 1 = 0$ has no real solution.
 $w^2 - 1 = 0 \Rightarrow w = \pm 1.$

Back-substitute $w = 2^x$: $2^x = \pm 1.$
 $2^x = -1$ has no real solution.
 $2^x = 1 \Rightarrow 2^x = 2^0 \Rightarrow x = 0.$

QUESTION 4

Let $y = f^{-1}(t)$. Then: $t = \frac{3y}{y^2 + 1} \therefore t(y^2 + 1) = 3y \therefore ty^2 - 3y + t = 0$.

Use the quadratic formula to solve for y : $y = \frac{3 \pm \sqrt{9 - 4t^2}}{2t}$.

To choose between the two potential solutions for y , use the fact that

$$f(a) = b \Rightarrow f^{-1}(b) = a.$$

For example, since $f(0) = 0$ it is required that $f^{-1}(0) = 0$.

Since $\frac{3 + \sqrt{9 - 4(0)^2}}{2(0)}$ is undefined, it follows by elimination that $y = f^{-1}(t) = \frac{3 - \sqrt{9 - 4t^2}}{2t}$.

Note: $\lim_{t \rightarrow 0} \frac{3 - \sqrt{9 - 4t^2}}{2t} = 0$.

QUESTION 5

a. Let $y = \frac{u}{v}$ where $u = \cos^2 x$ and $v = x$.

Then $\frac{dv}{dx} = 1$.

To find $\frac{du}{dx}$ use the **Chain Rule**:

Let $w = \cos x$ so that $u = w^2$.

Then $\frac{du}{dw} = 2w$ and $\frac{dw}{dx} = -\sin x$.

Then $\frac{du}{dx} = \frac{du}{dw} \times \frac{dw}{dx} = (2w)(-\sin x) = -2 \cos x \sin x$.

From the **Quotient Rule**:

$$\frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2} = \frac{(x)(-2 \cos x \sin x) - (\cos^2 x)(1)}{x^2} = \frac{-2x \sin x \cos x - \cos^2 x}{x^2}.$$

- b.** To find x -coordinate of stationary points solve $\frac{dy}{dx} = 0$:

$$\frac{dy}{dx} = \frac{-2x \sin x \cos x - \cos^2 x}{x^2} = \frac{-\cos x(2x \sin x + \cos x)}{x^2} = 0$$

Use the null factor theorem:

$$\therefore \cos x = 0 \quad \text{or} \quad 2x \sin x + \cos x = 0.$$

Solve for x : $\cos x = 0$

$$x = \frac{(2n+1)\pi}{2} \text{ where } n \in J.$$

Solve for y : $\cos x = 0$

$$y = \frac{2(0)^2}{(2n+1)\pi} = 0$$

c. $2x \sin x + \cos x = 0$

$$\therefore 2x \sin x = -\cos x$$

$$\therefore 2x \frac{\sin x}{\cos x} = -1$$

$$\therefore 2x \tan x = -1.$$

QUESTION 6

Let the point where $y = \frac{x}{2} - 1$ is normal to $y = bx^4 + 1$ have coordinates (x_1, y_1) .

Gradient of normal to curve at (x_1, y_1) :

$$m_{\text{tangent}} = \frac{dy}{dx} = 4bx^3.$$

Therefore $m_{\text{tangent}} = 4bx_1^3$ at (x_1, y_1) .

$$m_{\text{normal}} m_{\text{tangent}} = -1 \therefore m_{\text{normal}} = -\frac{1}{m_{\text{tangent}}} \therefore m_{\text{normal}} = -\frac{1}{4bx_1^3}. \quad (1)$$

$$\text{Gradient of line: } m = \frac{1}{2}. \quad (2)$$

$$\begin{aligned} \text{Equate equations (1) and (2): } -\frac{1}{4bx_1^3} &= \frac{1}{2} \Rightarrow 2bx_1^3 = -1 \\ \therefore 2bx_1^3 &= -1. \end{aligned} \quad (3)$$

Since the point (x_1, y_1) lies on both the line and the curve, it follows that

$$y_1 = \frac{x_1}{2} - 1. \quad (4)$$

$$y_1 = bx_1^4 + 1. \quad (5)$$

Equate equations (4) and (5):

$$\frac{x_1}{2} - 1 = bx_1^4 + 1 \therefore 2bx_1^4 - x_1 + 4 = 0. \quad (6)$$

From equation (3) it follows that $2bx_1^4 = -x_1$.


$$\text{Substitute } 2bx_1^4 = -x_1 \text{ into equation (6): } -x_1 - x_1 + 4 = 0 \therefore x_1 = 2.$$

$$\text{Substitute } x_1 = 2 \text{ into equation (4): } y_1 = 0.$$

$$\text{Substitute } x_1 = 2 \text{ and } y_1 = 0 \text{ into equation (5): } 0 = b(2)^4 + 1 \therefore b = -\frac{1}{16}.$$

QUESTION 7

a. and b.



c. Note that $y = |2f(x)| = \left| 2\cos\left(\frac{x}{2}\right) \right| = \begin{cases} 2\cos\left(\frac{x}{2}\right) & \text{for } -\pi \leq x \leq \pi \\ -2\cos\left(\frac{x}{2}\right) & \text{for } \pi \leq x \leq 3\pi \end{cases}$

Therefore:

$$\begin{aligned} \text{Area} &= \int_{-\pi}^{\pi} 2\cos\left(\frac{x}{2}\right) - \cos\left(\frac{x}{2}\right) dx + \int_{\pi}^{3\pi} -2\cos\left(\frac{x}{2}\right) - \cos\left(\frac{x}{2}\right) dx \\ &= \int_{-\pi}^{\pi} \cos\left(\frac{x}{2}\right) dx - 3 \int_{\pi}^{3\pi} \cos\left(\frac{x}{2}\right) dx \\ &= 2 \left[\sin\left(\frac{x}{2}\right) \right]_{-\pi}^{\pi} - 6 \left[\sin\left(\frac{x}{2}\right) \right]_{\pi}^{3\pi} \\ &= 2 \left\{ \sin\left(\frac{\pi}{2}\right) - \sin\left(-\frac{\pi}{2}\right) \right\} - 6 \left\{ \sin\left(\frac{3\pi}{2}\right) - \sin\left(\frac{\pi}{2}\right) \right\} = 2\{1 - (-1)\} - 6\{-1 - 1\} = 16. \end{aligned}$$

QUESTION 8

$$g(x) = f(-2x) + 1.$$

Transformations: Reflection in the vertical axis.
 Dilation by a scale factor of $\frac{1}{2}$ from the vertical axis.
 Translation by +1 units from the x-axis.

The order in which these transformations are applied does not matter.

QUESTION 9

a. Volume of a right circular cone: $V = \frac{1}{3}\pi r^2 h$.

$$\text{Semi-vertical angle of } 45^\circ \Rightarrow \tan 45^\circ = \frac{r}{h} \Rightarrow r = h.$$

$$\text{Therefore: } V = \frac{1}{3}\pi h^3.$$

$$\text{After 30 minutes } V = (30)(0.3) = 9 \text{ m}^3.$$

$$\text{Therefore: } 9 = \frac{1}{3}\pi h^3$$

$$h^3 = \frac{27}{\pi}$$

$$h = \frac{3}{\pi^{1/3}} \text{ meters.}$$

b. From the chain rule: $\frac{dh}{dt} = \frac{dh}{dV} \times \frac{dV}{dt}$.

$$\text{Given: } \frac{dV}{dt} = 0.3 \text{ m}^3 / \text{min}.$$

$$\text{From part (a): } V = \frac{1}{3}\pi h^3 \therefore \frac{dV}{dh} = \pi h^2 \therefore \frac{dh}{dV} = \frac{1}{\pi h^2}.$$


$$\text{Therefore: } \frac{dh}{dt} = \frac{1}{\pi h^2} \times 0.3.$$

$$\text{Substitute } t = 30 \text{ minutes} \Rightarrow h = \frac{3}{\pi^{1/3}} \text{ meters:}$$

$$\frac{dh}{dt} = \frac{1}{\pi \left(\frac{3}{\pi^{1/3}} \right)^2} \times 0.3 = \frac{0.3\pi^{1/3}}{9\pi} = \frac{1}{30\pi^{1/3}} \text{ m/min.}$$

QUESTION 10

Draw a tree diagram:



$\Pr(\text{Living in the city in 2 years time} \mid \text{currently living in the country})$

$$= \left(\frac{1}{7}\right)\left(\frac{9}{10}\right) + \left(\frac{6}{7}\right)\left(\frac{1}{7}\right)$$

$$= \frac{9}{70} + \frac{6}{49} = \frac{63}{490} + \frac{60}{490} = \frac{123}{490}.$$

QUESTION 11

a. By definition: $p + \frac{1}{7} + 3p + 3p + 4p + p = 1$

$$\therefore 12p = \frac{6}{7}$$

$$\therefore p = \frac{1}{14}.$$

b. Conditional probability: Require $\Pr(X < 5 \mid X > 2)$.

x	1	2	3	4	5	6
$\Pr(X = x)$	p	$\frac{1}{7}$	$3p$	$3p$	$4p$	p

$$\Pr(X < 4 \mid X > 2) = \frac{\Pr(X = 3) + \Pr(X = 4)}{\Pr(X = 3) + \Pr(X = 4) + \Pr(X = 5) + \Pr(X = 6)} = \frac{6p}{11p} = \frac{6}{11}.$$

BONUS QUESTION

QUESTION 12

- a. Let $y = uv$ where $u = x$ and $v = e^{-x}$.

Then $\frac{du}{dx} = 1$ and $\frac{dv}{dx} = -e^{-x}$.

From the Product Rule: $\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$

$$\therefore \frac{dy}{dx} = (x)(-e^{-x}) + (e^{-x})(1) = e^{-x} - xe^{-x}.$$

- b. Anti-differentiate both sides of $\frac{dy}{dx} = e^{-x} - xe^{-x}$ with respect to x :

$$y = \int (e^{-x} - xe^{-x}) dx$$

Substitute $y = xe^{-x}$:

$$\therefore xe^{-x} = \int e^{-x} dx - \int xe^{-x} dx = -e^{-x} - \int xe^{-x} dx$$

$$\therefore \int xe^{-x} dx = -xe^{-x} - e^{-x} = -(x+1)e^{-x},$$

Note: The arbitrary constant of anti-differentiation is omitted since only *an* anti-derivative is required.

- c. (i) By definition: $\overline{X} = \int_0^{\log_e 3} (x) \frac{3}{2} e^{-x} dx = \frac{3}{2} \int_0^{\log_e 3} xe^{-x} dx.$

Substitute the result from part (b):

$$\overline{X} = -\frac{3}{2} \left[(x+1)e^{-x} \right]_0^{\log_e 3} = -\frac{3}{2} \left\{ (1 + \log_e 3)e^{-\log_e 3} - 1 \right\}$$

Use the log rule $\log_a \frac{1}{b} = -\log_a b$:

$$= \frac{3}{2} \left\{ 1 - (1 + \log_e 3)e^{\log_e \frac{1}{3}} \right\}$$

Use the log rule $a^{\log_a b} = b$:

$$= \frac{3}{2} \left\{ 1 - \frac{1}{3} (1 + \log_e 3) \right\} = 1 - \frac{1}{2} \log_e 3$$

Use the log rule $c \log_a b = \log_a b^c$ and the power rule $b^{1/2} = \sqrt{b}$:

$$= 1 - \log_e \sqrt{3}.$$

(ii) Let the median value of X be equal to k .

By definition: $\frac{1}{2} = \int_0^k \frac{3}{2} e^{-x} dx$.

Therefore:

$$\frac{1}{3} = \int_0^k e^{-x} dx = -[e^{-x}]_0^k = 1 - e^{-k}$$

$$\therefore \frac{2}{3} = e^{-k}$$

$$-k = \log_e \frac{2}{3} \quad .$$

$$k = -\log_e \frac{2}{3} = \log_e \frac{3}{2}$$