



Online & home tutors Registered business name: itute ABN: 96 297 924 083

Mathematical Methods

2008

Trial Examination 2

SECTION 1 Multiple-choice questions

Instructions for Section 1

Answer **all** questions.

Choose the response that is **correct** for the question.

A correct answer scores 1, an incorrect answer scores 0.

Marks will **not** be deducted for incorrect answers.

No marks will be given if more than one answer is completed for any question.

Question 1

Consider the function $f : D \rightarrow R$, $f(x) = \tan\left(\frac{x}{2}\right)$ and $D = \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$. Which one of the following statements is true?

A. f has asymptotes at $x = -\frac{\pi}{2}$ and $x = \frac{\pi}{2}$.

B. f has asymptotes at $x = -\pi$ and $x = \pi$.

C. f has no asymptotes.

D. f has asymptotes at $x = -2\pi$ and $x = 2\pi$.

E. f has asymptotes at $x = -\frac{\pi}{4}$ and $x = \frac{\pi}{4}$.

Question 2

Given $b = 2 \log_2\left(\frac{a}{2}\right)$, $a =$

A. $e^{\frac{1}{2}(b+2)\log_e 2}$

B. $2e^{\frac{1}{2}(b+2)}$

C. $2^{\frac{1}{2}(b+1)}$

D. $2^{\frac{1}{2}(b-1)}$

E. $e^{\frac{b}{2}\log_e 2+1}$

Question 3

The equation $e^{2x} - 2e^x + k = 0$ has two solutions when

A. $k \in R$

B. $k \in R^+$

C. $k \in R^-$

D. $k \in (-\infty, 1)$

E. $k \in (0.3, 0.7)$

Question 4

If $f(x) = 2 \cos(3x)$, then $\frac{1}{4} f\left(\frac{\pi}{6} - \frac{1}{3}x\right) =$

A. $\frac{1}{2} \cos(x)$


B. $\frac{1}{2} \sin(x)$

C. $\frac{1}{2} \cos\left(\frac{\pi}{6} - \frac{1}{3}x\right)$

D. $\frac{1}{2} \sin\left(\frac{\pi}{6} - \frac{1}{3}x\right)$

E. $\frac{1}{2} \sin\left(\frac{1}{3}x\right)$

Question 5



The function shown in the graph is in the form $f(x) = a\sqrt{1-bx^2}$. The values of a and b are respectively

- A. 1 and 4 B. 4 and 1 C. 1 and 2 D. 1 and 1 E. 2 and $\frac{1}{4}$

Question 6

For $a, b, c \in \mathbb{R}^+$, the range of the function $|a - be^{-x}| - c$ is

- A. $[a - c, \infty)$ B. $[0, \infty)$ C. $(-\infty, b + c]$ D. $[-c, \infty)$ E. $(-\infty, c]$

Question 7

Which one of the following expressions **cannot** be expressed as a polynomial function of x ?

- A. $(3\sqrt{x} + x)(3\sqrt{x} - x)$
 B. $(1 - x\sqrt{2})(2 + x\sqrt{2})$
 C. $\sqrt[3]{x^3 - 3x^2 + 3x - 1}$
 D. $\frac{x^{\frac{3}{2}} - (2x)^{\frac{5}{2}}}{x^{-\frac{3}{2}}}$
 E. $|x^4 - x^2 - 1|$

Question 8

Which one of the following **cannot** be the possible number of intersections of a function and its inverse?

- A. 0 B. 1 C. 2 D. No more than 2 E. Any number

Question 9

If $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = \cos\left(\frac{\pi x}{2}\right)$ and $g : [1, \infty) \rightarrow \mathbb{R}$, $g(x) = \frac{1}{x} - 1$, the range of $f \circ g$ is

- A. $[0, 1)$ B. $(0, 1]$ C. $[-2, 0]$ D. $[0, -2]$ E. $[0, -2)$


Question 10

Given $f(x) = 1 - \sqrt{x}$, which one of the following statements is true?


- A. $f(xy) + f\left(\frac{x}{y}\right) = 2f(x)$
- B. $f\left(\frac{x}{y}\right) - f\left(\frac{y}{x}\right) = 2[f(x) - f(y)]$
- C. $f(xy) + f\left(\frac{x}{y}\right) = f(x)f(y)$
- D. $f\left(\frac{x}{y}\right) = \frac{f(x) - f(y)}{1 - f(y)}$
- E. $f(2x) - f(2y) = [f(x) - f(y)][f(x) + f(y)]$

Question 11

For graph of $y = f(x)$ is shown below.



Which one of the following is the graph of $y = f'(x)$?



Question 12

Given $P(x) = \frac{f(\log_e x)}{g(\sqrt{x})}$, $P'(x) =$

- A. $\frac{g(\sqrt{x})f'(\log_e x) - f(\log_e x)g'(\sqrt{x})}{2x\sqrt{x}g(x)}$
- B. $\frac{2\sqrt{x}g(\sqrt{x})f'(\log_e x) - xf(\log_e x)g'(\sqrt{x})}{2x\sqrt{x}[g(\sqrt{x})]^2}$
- C. $\frac{g(\sqrt{x})f'(\log_e x) - f(\log_e x)g'(\sqrt{x})}{2x\sqrt{x}[g(\sqrt{x})]^2}$
- D. $\frac{g(x)f'(x) - f(x)g'(x)}{2x\sqrt{x}[g(x)]^2}$
- E. $\frac{2\sqrt{x}g(x)f'(x) - xf(x)g'(x)}{2x\sqrt{x}[g(x)]^2}$

Question 13

The equation of the normal to the curve $y = 2x^3 - 3ax^2 + 5$ at $x = a$ is

- A. $y = x$ B. $y = 0$ C. $x = 0$ D. $y = a$ E. $x = a$

Question 14

The gradient of the tangent to the curve $y = \frac{x \tan(\sqrt{xe^x})}{2\pi}$ at $x = \frac{3}{4}$ equals


- A. 2.373 B. 2.372 C. 23.41 D. 23.42 E. 3.668

Question 15

Using the approximation $f(x+h) \approx f(x) + hf'(x)$ for $f(x) = \sqrt{1-x}$, the approximate value of $f(-3.1)$ is

- A. $\frac{79}{40}$ B. $\frac{81}{40}$ C. $\frac{197}{100}$ D. $\frac{2531}{1250}$ E. $\frac{4937}{2500}$

Question 16



Using ‘right rectangles’ of unit width, the approximate area bounded by $x = -1$, $x = 3$ and the two curves $y = f(x)$ and $y = \frac{1}{2}f(x)$ shown in the figure above is closest to

- A. 5.4 B. 6.0 C. 6.6 D. 10.8 E. 13.2

Question 17

Given the continuous and **decreasing** function $f(x) = (x + 2)g(x)$, and $F(x) = \int f(x)dx$, the area bounded by the x -axis, $x = a$, $x = b$ and $y = f(x)$ for $a < -2 < b$ is

- A. $2F(-2) - F(a) - F(b)$
B. $F(b) - F(a)$
C. $F(a) - F(b)$
D. $F(b) + F(a)$
E. $2F(-2) + F(a) - F(b)$


Question 18

Roll a fair die twice.

The probability of getting the difference of the two uppermost numbers greater than 1 is

- A. $\frac{2}{3}$ B. $\frac{5}{6}$ C. $\frac{5}{9}$ D. $\frac{11}{18}$ E. $\frac{13}{18}$

Question 19



The graph above shows a probability distribution of random variable X . The value of $\bar{X}$ equals

- A. $\frac{51}{15}$ B. $\frac{53}{15}$ C. $\frac{51}{30}$ D. $\frac{53}{30}$ E. 3

Question 20

	$X > 0.36$	$X < 0.36$	
$X > 0.64$	0.19		
$X < 0.64$		0.48	
			1

The above is a probability table for continuous random variable X with some missing probability values. When $X < 0.64$, the probability that $X > 0.36$ equals

- A. $\frac{11}{27}$ B. $\frac{13}{25}$ C. $\frac{33}{100}$ D. $\frac{33}{52}$ E. $\frac{19}{48}$

Question 21

The heights of the children in a queue for an amusement park ride are normally distributed. 95.4% of the children in the queue have heights between 1.25 m and 1.35 m, i.e. within $\mu \pm 2\sigma$. The percentage of the children in the queue with heights between 1.28 m and 1.38 m is closest to

- A. 95% B. 90% C. 85% D. 80% E. 75%

Question 22

The waiting time t (hours) to see a doctor has a probability distribution given by the probability density function $f(t) = \frac{1}{\sqrt{3} \cos^2 t}$, $0 \leq t \leq \frac{\pi}{3}$. The median waiting time t is closest to

- A. 0.9 B. 0.8 C. 0.7 D. 0.6 E. 0.5

SECTION 2 Extended-answer questions

Instructions for Section 2

Answer **all** questions.

A decimal approximation will not be accepted if an **exact** answer is required to a question.

In questions where more than one mark is available, appropriate working **must** be shown.

Where an instruction to **use calculus** is stated for a question, you must show an appropriate derivative or antiderivative.


Unless otherwise indicated, the diagrams in this exam are **not** drawn to scale.

Question 1

Consider the function $f(x) = \frac{16}{x^4} - 1$. Length is measured in metres and time in seconds.

- a. Show that the equations of the inverse of $f(x)$ is $y = \pm \frac{2}{(x+1)^{\frac{1}{4}}}$. 2 marks

- b. Sketch the graph of the inverse of $f(x)$. Label the asymptote(s) and intercept(s).



2 marks

- c. Let L be the length of a vertical line touching the inverse of $f(x)$. Express L in terms of x .

1 mark

The vertical line is at $x = 0$ when $t = 0$, and it moves in the positive x direction at a speed of 2 metres per second.

- d. i. Show that the rate of decrease of L is $\frac{2}{(x+1)^{\frac{5}{4}}}$.

- d. ii. Find the rate of decrease of L when $t = 7.5$.


2 + 1 = 3 marks

- e. Find the average rate of increase of the area swept by the vertical line in the time interval $[0, 7.5]$.

3 marks

Total 11 marks

Question 2



The diagram above shows a pond with water filled to a depth of h m. The shaded part is the water surface and has an area $A = 2\pi[1 - (1 - h)^2]$ m². The volume of water is $V = 2\pi\left[\frac{2}{3} - (1 - h) + \frac{(1 - h)^3}{3}\right]$ m³.

- a. Factorise $2\pi[1 - (1 - h)^2]$ completely.

1 mark

- b. Expand $\frac{2}{3} - (1 - h) + \frac{(1 - h)^3}{3}$.

1 mark

- c. Find the maximum value of V .

1 mark

The pond is initially full. Water is drained from the pond at 2 litres per second. (Note: 1 litre = 1000 cm³)

- d. How long will it take to empty the pond?

1 mark

- e. Find the rate of decrease of the depth h with respect to t when $h = 0.5$ m.

3 marks

- f. Hence find the rate of decrease of the area A with respect to t when $h = 0.5$ m.

2 marks

The draining is stopped when $h = 0.5$ m, and pebbles of volume 0.831 m^3 are placed (submerged) at the bottom of the pond.

- g. i. Determine the height (3 decimal places in m) of the water surface from the bottom of the pond.


- ii. Does the rate of decrease of the depth h with respect to t change when draining is resumed? Explain.

$2 + 2 = 4$ marks

Total 13 marks

Question 3

The following drawing shows the cross-section of four successive identical wave crests. **The seabed is at $y = -2$.** Length is measured in metres.



- a. Find the exact values of a , b , c , h and k .

3 marks

- b. Write down the exact coordinates of point P, the intersection of the cosine curve and the semi-circle.

1 mark

- c. i. Find the equation (including its domain) of the semi-circle in the third wave crest.

- ii. Find the equation (including its domain) of the cosine curve in the fourth wave crest.

2 + 2 = 4 marks

d. i. Find the exact cross-sectional area of the second wave crest above the seabed.

d. ii. Find the volume of water (nearest m^3) above the seabed contained in the second wave crest when the wave front is 10 m long, assuming the wave crest has a uniform cross-section along the wave front.

3 + 1 = 4 marks

Total 12 marks


Question 4

Consider $f(x) = a \log_e(x) + \frac{1}{2}$.

a. Find $f^{-1}(x)$.

2 marks

b. Let $a = 2$. Sketch the graphs of $y = f(x)$ and $y = f^{-1}(x)$ on the same set of axes.



2 marks

c. i. Find the value of a (3 decimal places) such that the graphs of $y = f(x)$ and $y = f^{-1}(x)$ touch each other once only.

c. ii. Find the coordinates (3 decimal places) of the contact point.

3 + 1 = 4 marks

Total 8 marks

Question 5

A machine is calibrated to fill cans with 375.5 ml of soft drink. The volume of soft drink (x ml) in a can is a random variable X with a probability distribution given by probability density function $f(x) = e^{-\pi(x-375.5)^2}$.

a. Out of 1000 filled cans of soft drink, how many are expected to be over 375 ml?

2 marks

692 out of 1000 cans are over **375.3** ml without changing the original setting of the machine. The manager of the soft drink company changes the setting of the machine to fill cans with k ml so that 692 out of 1000 cans are over **375** ml.

b. Find the value of k .

3 marks

The company has 5 such machines at the new setting. The manager takes a can at random from each machine.

- c. What is the probability (3 decimal places) that 4 of them are under 375 ml?

2 marks

The manager makes another inspection and takes **two** cans at random from each machine.

- d. What is the probability (3 decimal places) that 8 of the 10 cans are under 375 ml?

1 mark

- e. The manager measured 3 of them and found 2 cans under 375 ml. What is the probability (3 decimal places) that 8 of the 10 cans are under 375 ml?

2 marks

Another company has only two machines, A and B. The volume of soft drink in a can filled by machine A is a random variable with a normal distribution, $\mu = 375.5$ ml and $\sigma = 0.3989$ ml. The volume of soft drink filled by machine B is a random variable with a normal distribution of the same mean $\mu = 375.5$ ml but different standard deviation $\sigma = 0.7978$ ml. The two machines produce **the same number** of cans of soft drink. All cans of soft drink produced by the two machines are now put together. Let the volume of drink in a can be random variable Y . **Information:** For a normal distribution,

$\Pr(\mu - \sigma < X < \mu + \sigma) = 0.683$, $\Pr(\mu - 2\sigma < X < \mu + 2\sigma) = 0.954$ and $\Pr(\mu - 3\sigma < X < \mu + 3\sigma) = 0.997$.

- f. i. Describe the shape of the probability distribution of Y .

- f. ii. It is true that $\Pr(374.15 < Y < 376.85) = 0.954$. Is Y normally distributed? Justify your answer.

1 + 3 = 4 marks

Total 14 marks

End of exam 2