

**Year 2009**

**VCE**

**Mathematical Methods**

**CAS**

**Trial Examination 2**



**KILBAHA MULTIMEDIA PUBLISHING**  
**PO BOX 2227**  
**KEW VIC 3101**  
**AUSTRALIA**

**TEL: (03) 9817 5374**  
**FAX: (03) 9817 4334**  
**kilbaha@gmail.com**  
**<http://kilbaha.googlepages.com>**

## **IMPORTANT COPYRIGHT NOTICE**

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Pty Ltd.
  - The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
  - For authorised copying within Australia please check that your institution has a licence from Copyright Agency Limited. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.
  - Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.
  - Schools which purchase a licence to use this material may distribute this electronic file to the students at the school for their exclusive use. This distribution can be done either on an Intranet Server or on media for the use on stand-alone computers.
  - Schools which purchase a licence to use this material may distribute this printed file to the students at the school for their exclusive use.
- 
- **The Word file (if supplied) is for use ONLY within the school.**
  - **It may be modified to suit the school syllabus and for teaching purposes.**
  - **All modified versions of the file must carry this copyright notice.**
  - **Commercial use of this material is expressly prohibited.**

**Victorian Certificate of Education  
2009**

**STUDENT NUMBER**

Figures  
Words

|  |  |
|--|--|
|  |  |
|  |  |

|  |  |  |  |  |  |
|--|--|--|--|--|--|
|  |  |  |  |  |  |
|  |  |  |  |  |  |

Letter

|  |
|--|
|  |
|--|

**MATHEMATICAL METHODS CAS  
Trial Written Examination 2**

Reading time: 15 minutes

Total writing time: 2 hours

**QUESTION AND ANSWER BOOK**

**Structure of book**

| Section | Number of questions | Number of questions to be answered | Number of marks |
|---------|---------------------|------------------------------------|-----------------|
| 1       | 22                  | 22                                 | 22              |
| 2       | 5                   | 5                                  | 58              |
|         |                     |                                    | Total 80        |

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers, a protractor, set-squares, aids for curve sketching, one bound reference, one approved **graphics** calculator ( memory DOES NOT need to be cleared ) and, if desired, one scientific calculator.
- Students are NOT permitted to bring into the examination room: blank sheets of paper and/or white out liquid/tape.

**Materials supplied**

- Question and answer booklet of 29 pages with a detachable sheet of miscellaneous formulas at the end of this booklet.
- Answer sheet for multiple choice questions.

**Instructions**

- Detach the formula sheet from the end of this book during reading time.
- Write your **student number** in the space provided above on this page.
- Check that your **name** and **student number** as printed on your answer sheet for multiple-choice questions are correct, and sign your name in the space provided to verify this.
- All written responses must be in English.

**At the end of the examination**

- Place the answer sheet for multiple-choice questions inside the front cover of this book.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.**

**SECTION 1****Instructions for Section I**

Answer **all** questions in pencil on the answer sheet provided for multiple-choice questions.

A correct answer scores 1 mark, an incorrect answer scores 0.

Marks will **not** be deducted for incorrect answers.

No mark will be given if more than one answer is completed for any question.

**Question 1**

If  $f(x) = g(x) \log_e(2x)$ ,  $g\left(\frac{e}{2}\right) = \frac{e}{2}$  and  $g'\left(\frac{e}{2}\right) = 1$  then  $f'\left(\frac{e}{2}\right)$  is equal to

- A. 1
- B.  $\frac{e}{2} + 1$
- C.  $\frac{e}{2}$
- D. 2
- E. 3

**Question 2**

Given the two linear simultaneous equations

$2x + (p+1)y = 4$  and  $px + 6y = q$ , then

- A. if  $p = 3$  and  $q = 6$  there is an infinite number of solutions.
- B. if  $p = 3$  and  $q = 6$  there is no solution.
- C. if  $p = 3$  and  $q \in \mathbb{R}$  there is a unique solution.
- D. if  $p = -4$  and  $q \in \mathbb{R}$  there is a unique solution.
- E. if  $p = -4$  and  $q \neq 6$  there is more than one solution.

**Question 3**


The average value of the function with the rule  $f(x) = x^3 + e^{2x}$  between  $x = 0$  and  $x = 2$  is equal to

- A.  $\frac{7 + e^4}{2}$
- B.  $\frac{7 + e^4}{4}$
- C.  $12 + 2e^4$
- D.  $5 + e^4$
- E.  $\frac{11 + 2e^4}{2}$

**Question 4**

The transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ , which maps the curve with equation  $y = x^4$  to the curve with equation  $y = (3x + 6)^4 - 2$ , could have the rule

- A.  $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 6 \\ -2 \end{bmatrix}$
- B.  $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 6 \\ 2 \end{bmatrix}$
- C.  $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} -2 \\ 2 \end{bmatrix}$
- D.  $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} \frac{1}{3} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 2 \\ 2 \end{bmatrix}$
- E.  $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} \frac{1}{3} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} -2 \\ -2 \end{bmatrix}$

**Question 5**

The diagram above show the graph of  $y = -x^2 + 3x - 3$  and the tangent to the graph at the point  $P$ , where  $x = 3$ . The tangent has the equation  $y = mx + c$ . The shaded area  $A$  is the area between the graph, the tangent and the  $y$ -axis. Which of the following is **FALSE**?

- A.  $c = 6$
- B.  $m = -3$
- C.  $-3m + 6 = -3$
- D.  $m < 0$  and  $c > 0$
- E.  $A = \int_0^3 (x^2 + (m-3)x + (c+3)) dx$

**Question 6**

If  $f(x) = \frac{1}{x^2}$  then the best approximation to  $\frac{1}{3.99^2}$  is given by

- A.  $f(4) - 0.01f'(4)$
- B.  $f(4) + 0.01f'(4)$
- C.  $f(2) - 0.01f'(4)$
- D.  $f(2) + 0.01f'(2)$
- E.  $f(2) - 0.01f'(2)$

**Question 7**

Which of the following is **NOT** a measure of the area bounded by the graph of  $y = x^2 - a^2$  and the  $x$ -axis?

- A.  $\left| \int_{-a}^a (x^2 - a^2) dx \right|$
- B.  $2 \int_{-a}^0 \sqrt{a^2 - x} dx$
- C.  $2 \int_0^a (a^2 - x^2) dx$
- D.  $\int_{-a}^a (a^2 - x^2) dx$
- E.  $2 \left| \int_{-a^2}^0 \sqrt{x + a^2} dx \right|$

**Question 8**

If  $a$  is a real positive constant, then the inverse of the function

$f : \mathbb{R}^+ \rightarrow \mathbb{R}$ ,  $f(x) = x^2 + a$  is

- A.  $f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f^{-1}(x) = \frac{1}{x^2 + a}$
- B.  $f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f^{-1}(x) = \sqrt{a} - \sqrt{x}$
- C.  $f^{-1} : (a, \infty) \rightarrow \mathbb{R}$ ,  $f^{-1}(x) = -\sqrt{x - a}$
- D.  $f^{-1} : (a, \infty) \rightarrow \mathbb{R}$ ,  $f^{-1}(x) = \sqrt{x - a}$
- E.  $f^{-1} : (0, \infty) \rightarrow \mathbb{R}$ ,  $f^{-1}(x) = \sqrt{x} - \sqrt{a}$

**Question 9**

If  $f(x) = f(-x)$  and  $\int_{-6}^6 f(x) dx = 10$ , then  $\int_0^6 (2f(x) - 1) dx$  is equal to

- A. 4
- B. 14
- C. 7
- D.  $10 - x$
- E.  $20 - x$

**Question 10**

The graph of the function  $f(x) = x^2 \log_e |x|$  has

- A. a local maximum at  $x = 0$  and local minimums at  $x = \pm \frac{1}{\sqrt{e}}$
- B. local minimums at  $x = \pm \frac{1}{\sqrt{e}}$
- C. a local minimum at  $x = 0$
- D. a point of inflexion at  $x = 0$
- E. a point of inflexion at  $x = 0$  and local minimums at  $x = \pm \frac{1}{\sqrt{e}}$



**Question 11**

Given the function  $f(x) = b + \frac{a}{x-b}$  where  $a$  and  $b$  are non-zero real constants,

and  $y = f^{-1}(x)$  is the inverse function.

Which of the following statements is **FALSE**?

- A. The graph of  $y = f(x)$  has the line  $x = b$  as a vertical asymptote and the line  $y = b$  as a horizontal asymptote.
- B. The domain and range of both  $f$  and  $f^{-1}$  are  $R \setminus \{b\}$ .
- C. The inverse function  $f^{-1} = f$ .
- D. The graph of  $y = f(x)$  passes through  $\left(0, b - \frac{a}{b}\right)$  and the graph of  $y = f^{-1}(x)$  passes through  $\left(b - \frac{a}{b}, 0\right)$ .
- E. The graphs of  $y = f(x)$  and  $y = f^{-1}(x)$  always intersect on the line  $y = x$  at the points  $(b \pm \sqrt{a}, b \pm \sqrt{a})$ .

**Question 12**

A certain curve has its gradient given by  $2 \cos\left(\frac{x}{2}\right)$ . If the curve crosses the x-axis

at  $x = \frac{5\pi}{3}$ , then it crosses the y-axis at

- A.  $-\frac{1}{2}$
- B.  $\frac{1}{2}$
- C.  $-2$
- D.  $\frac{\sqrt{3}}{2}$
- E.  $2$

**Question 13**

A curve has a vertical asymptote at  $x = a$  and a horizontal asymptote at  $y = b$ . Which one of the following could be the equation of the curve?

A.  $y = \frac{x-b}{x+a}$

B.  $y = \frac{x-b}{x-a}$

C.  $y = \frac{b}{x-a}$

D.  $y = \frac{bx}{x-a}$

E.  $y = b + \frac{1}{x+a}$

**Question 14**

Let  $f : [0, \pi] \rightarrow \mathbb{R}$ ,  $f(x) = 2 \cos\left(\frac{x}{2}\right) - 2$ . The graph of  $f$  is transformed by a reflection in the  $x$ -axis, followed by a dilation of factor 2 from the  $y$ -axis, then a dilation by a factor of 3 from the  $x$ -axis. The resulting graph is defined by

A.  $g : [0, \pi] \rightarrow \mathbb{R}$ ,  $g(x) = 6 - 6 \cos(x)$

B.  $g : \left[0, \frac{\pi}{3}\right] \rightarrow \mathbb{R}$ ,  $g(x) = 4 + 4 \cos\left(\frac{3x}{2}\right)$

C.  $g : [0, 3\pi] \rightarrow \mathbb{R}$ ,  $g(x) = 4 - 4 \cos\left(\frac{x}{6}\right)$

D.  $g : [0, 2\pi] \rightarrow \mathbb{R}$ ,  $g(x) = 6 - 6 \cos\left(\frac{x}{4}\right)$

E.  $g : [0, 2\pi] \rightarrow \mathbb{R}$ ,  $g(x) = 4 - 4 \cos\left(\frac{3x}{2}\right)$

**Question 15**

If  $\Pr(A) = a$ ,  $\Pr(B) = b$  and  $\Pr(A \cap B) = p$  where  $0 < a < 1$ ,  $0 < b < 1$  and  $0 < p < 1$ , then  $\Pr(A' \cap B')$  is equal to

- A.  $1 - p$
- B.  $1 - (a + b)$
- C.  $1 - (a + b) - ab$
- D.  $2 - (a + b) - (1 - a)(1 - b)$
- E.  $1 + p - (a + b)$

**Question 16**

The function  $f : (-\infty, a) \rightarrow \mathbb{R}$  with the rule  $f(x) = x^4 - 4x^2$ , will have an inverse provided that

- A.  $a < 2$
- B.  $a > 0$
- C.  $a < \sqrt{2}$
- D.  $a < -\sqrt{2}$
- E.  $a > -\sqrt{2}$

**Question 17**

Anthony and Betty play  $n$  games of table tennis. The probability that Anthony wins any game is 0.4, no game can be a draw. If the probability that Betty wins no games is less than 0.01, then the minimum number of games they played is equal to

- A. 5
- B. 6
- C. 7
- D. 9
- E. 10


**Question 18**

If  $X$  is a normally distributed random variable with mean  $\mu$  and variance  $\frac{9\mu^2}{4}$ , then  $\Pr(X > 2\mu)$  is closest to

- A. 0.023
- B. 0.252
- C. 0.504
- D. 0.672
- E. 0.748

**Question 19**

Shown below is the graph of a function  $f$ .




Let  $g(x) = \int_0^x f(t) dt$ , then

- A.  $g(0)=1$   $g'(0)=0$   $g'(3)=0$
- B.  $g(0)=1$   $g'(0)=0$   $g'(3)=1$
- C.  $g(0)=0$   $g'(0)=9$   $g'(3)=0$
- D.  $g(0)=0$   $g'(0)=9$   $g'(3)=18$
- E.  $g(0)=0$   $g'(0)=0$   $g'(3)=0$


**Question 20**

The graph of  $f(g(x))$  is shown.




The graphs of  $y = f(x)$  and  $y = g(x)$  are best represented respectively as


**A.**




**B.**




**C.**



**D.**



**E.**



**Question 21**

A regular hexagon, has each side length being  $L$  cm. The area  $A$  of the hexagon is given by  $\frac{3\sqrt{3}}{2}L^2$ . If each of the six sides of the hexagon increases at a rate of  $\sqrt{3}$  cm/s, then the rate in  $\text{cm}^2/\text{s}$  at which the area of the hexagon is increasing, when the side lengths are  $\sqrt{3}$  cm, is given by

- A.  $9\sqrt{3}$
- B. 9
- C.  $6\sqrt{3}$
- D.  $\frac{9\sqrt{3}}{2}$
- E.  $\frac{27}{2}$

**Question 22**

The random variable  $X$  has the following probability distribution.

|              |               |     |     |               |
|--------------|---------------|-----|-----|---------------|
| $X$          | -2            | -1  | 1   | 2             |
| $\Pr(X = x)$ | $\frac{a}{2}$ | $a$ | $b$ | $\frac{b}{2}$ |

Which of the following statements is **FALSE**?

- A.  $3(a + b) = 2$
- B.  $E(X) = \sum x \Pr(X = x) = 2(b - a)$
- C.  $E(X^2) = 2$
- D.  $\text{var}(X) = 2 - 4b^2 + 8ab - 4a^2$
- E.  $E\left(\frac{1}{X}\right) = \frac{1}{2(b - a)}$

**END OF SECTION 1**

**SECTION 2****Instructions for Section 2**

Answer **all** questions in the spaces provided.

A decimal approximation will not be accepted if an **exact** answer is required to a question.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

**Question 1**

Consider the function  $f : R \rightarrow R$ ,  $f(x) = x^3 - 3x^2 + cx + d$ , where  $c$  and  $d$  are real numbers.

**a.** The coordinates of the turning point on the graph of  $y = f(x)$  are  $(-1, 5)$  and  $(u, v)$ .

**i.** Show that in this case  $c = -9$ ,  $d = 0$ ,  $v = -27$  and determine the value of  $u$ .

---

---

---

---

---

---

---

---

2 marks

**ii.** For what values of  $d$  does the graph of  $y = x^3 - 3x^2 - 9x + d$  cross the  $x$ -axis at three distinct points?

---

---

---

2 marks

- b.** For what values of  $c$  and  $d$  does the graph of  $y = x^3 - 3x^2 + cx + d$  have two distinct turning points?

---

---

---

---

---

---

2 marks

- c.** If the graph of  $y = f(x)$  is translated  $p$  units to the left away from the  $y$ -axis, it becomes the graph of  $y = x^3$ . Find the values of  $p$ ,  $c$  and  $d$  in this case.

---

---

---

---

---

---

---

---


2 marks





**Question 2**

The diagram below shows a “triangular” shade cloth, which is designed to block the direct sunlight onto a children’s playground. The cloth lies in a horizontal plane and has vertical posts erected at points O, A and B. The point O is the origin and the co-ordinates of the points A, P and B are  $(8,0)$ ,  $(4,1.5)$  and  $(4,8)$  respectively. The axes are shown on the drawing and the units are in metres.



The curve OPA has the form  $y = a \sin(nx)$ .

- a. Explain why  $a = 1.5$  and  $n = \frac{\pi}{8}$ .

---



---



---



---



---

1 mark

The curve OB has the form  $y = 16(1 - e^{-kx})$ .

- b.** Find the value of  $k$ .

---

---

---

---

---

---

---

1 mark

- c.** The curve BA is the reflection of the curve OB in the line  $x = 4$ .

- i.** Write down **two** transformations, which take the curve OB into the curve BA.

---

---

---

1 mark

- ii.** Hence write down a **function** in terms of  $k$ , which describes the curve BA.

---

---

---

---

---

2 marks



**Question 3**

At a cinema, the time  $T$  minutes, that customers have to wait in order to buy their tickets has a probability density function given by

$$f(t) = \begin{cases} bt^2 & \text{for } 0 \leq t \leq 4 \\ c(8-t) & \text{for } 4 < t \leq 8 \\ 0 & \text{otherwise} \end{cases}$$

- a. Show that  $b = \frac{3}{160}$  and  $c = \frac{3}{40}$ .

---

---

---

---

---

---

---

3 marks

- b. Sketch the graph of  $y = f(t)$  on the axes provided, clearly labelling the scale.



1 mark

- c. Find the probability that a customer waits more than six minutes to buy a ticket.

---

---

---

---

2 marks

- d. Find the mean time that customers have to wait in order to buy their tickets, give your answer correct to one decimal place.

---

---

---

---

2 marks

- e. Find the median time, correct to two decimal places, that customers have to wait in order to buy their tickets.

---

---

---

---

---

---

---

---

2 marks

The running times of movies shown at the cinema are normally distributed with a mean of 94 minutes, with a standard deviation of 10 minutes.

- f.** Find the probability that the running time of a movie is more than 109 minutes, give your answer correct to three significant figures.

---

---

---

---

---

1 mark

- g.** A certain cinema complex has four different movies playing. Find the probability that at least two of the movies have a running time of more than 109 minutes, give your answer correct to three significant figures.

---

---

---

---

---

2 marks

Pete goes to the movies once every week. He likes to see only action or comedy movies. If he sees an action movie one week, the probability that he sees an action movie the following week is 0.45, while if he sees a comedy one week, the probability that he sees an action movie the following week is 0.35. Suppose he has just seen an action movie.

- h.** What is the probability that of the next three movies he sees, exactly two are comedies, give your answer correct to three decimal places.

---

---

---

---

---

---

---

2 marks

- i.** In the long term, what percentage of the movies that Pete sees are action movies? Give your answer correct to one decimal place.

---

---

---

---

---

---

1 mark


Total 16 marks



**Question 4**

The diagram below shows part of the graph of the function  $f : (0, \infty) \rightarrow \mathbb{R}$  where  $f(x) = \frac{4}{x^2}$ .

Let  $P(a, f(a))$  where  $a > 0$  be a point on the graph of  $y = \frac{4}{x^2}$ .



- a. Show that the distance  $s$  from the origin  $O$  to the point  $P$  is given by  $s = \frac{\sqrt{16 + a^6}}{a^2}$ .

---

---

---

---

---

---

---

2 marks

- b i.** Find, the exact value of  $a$ , for which the distance  $s$  is a minimum.  
Verify that it is a minimum.

---

---

---

---

---

---

---

---

---

---

3 marks

- ii.** Find the minimum distance, give an exact answer.

---

---

---

---

1 mark

- c.i.** Find in terms of  $a$ , the equation of the normal to the curve  $y = f(x)$  at the point  $P$ .

---

---

---

---

---

---

---

---

2 marks

- ii.** Find the value of  $a$ , for which the normal passes through the origin.

---

---

---

---

1 mark

Total 9 marks

**Question 5**

Given the function  $g : \mathbb{R} \rightarrow \mathbb{R}$ ,  $g(x) = \sqrt{3} \sin(2x) + \cos(2x)$

- a.** Find the general solution of  $g(x) = 0$ .

---

---

1 mark

- b.** State the smallest positive value of  $T$  for which  $g(x+T) = g(x)$ .

---

---

1 mark

Given the function  $f : [0, 2\pi] \rightarrow \mathbb{R}$ ,  $f(x) = \sqrt{3} \sin(2x) + \cos(2x)$ .

- c.** Find  $\{x : f(x) = 0\}$ .

---

---

---

---

1 mark

- d.i.** Find  $\{x : f'(x) = 0\}$ .

---

---

---

---

---

---

2 marks

- ii. Find the exact coordinates of the maximum and minimum turning points on the graph of  $y = f(x)$ .

---




---



---

1 mark

- e. Sketch the graph of  $y = f(x)$  on the axes below, clearly labelling the scale.



2 marks

- f. If  $f(x) = A \sin(2(x + \alpha))$  state the values of  $A$  and  $\alpha$ .

---



---



---

2 marks  
Total 10 marks

**END OF EXAMINATION**

**EXTRA WORKING SPACE**

[illegible]

# **MATHEMATICAL METHODS CAS**

## **Written examination 2**

### **FORMULA SHEET**

#### **Directions to students**

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

## Mathematical Methods and CAS Formulas

### Mensuration

|                                    |                        |                      |                         |
|------------------------------------|------------------------|----------------------|-------------------------|
| area of a trapezium:               | $\frac{1}{2}(a+b)h$    | Volume of a pyramid: | $\frac{1}{3}Ah$         |
| curved surface area of a cylinder: | $2\pi rh$              | volume of a sphere:  | $\frac{4}{3}\pi r^3$    |
| volume of a cylinder:              | $\pi r^2 h$            | area of triangle:    | $\frac{1}{2}bc \sin(A)$ |
| volume of a cone:                  | $\frac{1}{3}\pi r^2 h$ |                      |                         |

### Calculus

|                                                                |                                                     |
|----------------------------------------------------------------|-----------------------------------------------------|
| $\frac{d}{dx}(x^n) = nx^{n-1}$                                 | $\int x^n dx = \frac{1}{n+1}x^{n+1} + c, n \neq -1$ |
| $\frac{d}{dx}(e^{ax}) = ae^{ax}$                               | $\int e^{ax} dx = \frac{1}{a}e^{ax} + c$            |
| $\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$                        | $\int \frac{1}{x} dx = \log_e x  + c$               |
| $\frac{d}{dx}(\sin(ax)) = a \cos(ax)$                          | $\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$      |
| $\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$                         | $\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$       |
| $\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$ |                                                     |

product rule:  $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$

quotient rule:  $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

Chain rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

approximation:  $f(x+h) \approx f(x) + h f'(x)$

### Probability

$\Pr(A) = 1 - \Pr(A')$

$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$

$\Pr(A/B) = \frac{\Pr(A \cap B)}{\Pr(B)}$

mean:  $\mu = E(X)$

variance:  $\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

| probability distribution |                                     | mean                                      | variance                                                 |
|--------------------------|-------------------------------------|-------------------------------------------|----------------------------------------------------------|
| discrete                 | $\Pr(X = x) = p(x)$                 | $\mu = \sum x p(x)$                       | $\sigma^2 = \sum (x - \mu)^2 p(x)$                       |
| continuous               | $\Pr(a < X < b) = \int_a^b f(x) dx$ | $\mu = \int_{-\infty}^{\infty} x f(x) dx$ | $\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$ |



# ANSWER SHEET

**STUDENT NUMBER**

Figures  
Words

|  |  |
|--|--|
|  |  |
|  |  |

|  |  |  |  |  |  |
|--|--|--|--|--|--|
|  |  |  |  |  |  |
|  |  |  |  |  |  |

Letter

|  |
|--|
|  |
|--|

**SIGNATURE** \_\_\_\_\_

## SECTION 1

|    |   |   |   |   |   |
|----|---|---|---|---|---|
| 1  | A | B | C | D | E |
| 2  | A | B | C | D | E |
| 3  | A | B | C | D | E |
| 4  | A | B | C | D | E |
| 5  | A | B | C | D | E |
| 6  | A | B | C | D | E |
| 7  | A | B | C | D | E |
| 8  | A | B | C | D | E |
| 9  | A | B | C | D | E |
| 10 | A | B | C | D | E |
| 11 | A | B | C | D | E |
| 12 | A | B | C | D | E |
| 13 | A | B | C | D | E |
| 14 | A | B | C | D | E |
| 15 | A | B | C | D | E |
| 16 | A | B | C | D | E |
| 17 | A | B | C | D | E |
| 18 | A | B | C | D | E |
| 19 | A | B | C | D | E |
| 20 | A | B | C | D | E |
| 21 | A | B | C | D | E |
| 22 | A | B | C | D | E |

**Year 2009**  
**VCE**  
**Mathematical Methods**  
**CAS**  
**Solutions**  
**Trial Examination 2**



**KILBAHA MULTIMEDIA PUBLISHING**  
**PO BOX 2227**  
**KEW VIC 3101**  
**AUSTRALIA**

**TEL: (03) 9817 5374**  
**FAX: (03) 9817 4334**  
**kilbaha@gmail.com**  
**<http://kilbaha.googlepages.com>**

### **IMPORTANT COPYRIGHT NOTICE**

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Pty Ltd.
  - The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
  - For authorised copying within Australia please check that your institution has a licence from Copyright Agency Limited. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.
  - Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.
  - Schools which purchase a licence to use this material may distribute this electronic file to the students at the school for their exclusive use. This distribution can be done either on an Intranet Server or on media for the use on stand-alone computers.
  - Schools which purchase a licence to use this material may distribute this printed file to the students at the school for their exclusive use.
- 
- **The Word file (if supplied) is for use ONLY within the school.**
  - **It may be modified to suit the school syllabus and for teaching purposes.**
  - **All modified versions of the file must carry this copyright notice.**
  - **Commercial use of this material is expressly prohibited.**

## SECTION 1

## ANSWERS

|    |   |   |   |   |   |
|----|---|---|---|---|---|
| 1  | A | B | C | D | E |
| 2  | A | B | C | D | E |
| 3  | A | B | C | D | E |
| 4  | A | B | C | D | E |
| 5  | A | B | C | D | E |
| 6  | A | B | C | D | E |
| 7  | A | B | C | D | E |
| 8  | A | B | C | D | E |
| 9  | A | B | C | D | E |
| 10 | A | B | C | D | E |
| 11 | A | B | C | D | E |
| 12 | A | B | C | D | E |
| 13 | A | B | C | D | E |
| 14 | A | B | C | D | E |
| 15 | A | B | C | D | E |
| 16 | A | B | C | D | E |
| 17 | A | B | C | D | E |
| 18 | A | B | C | D | E |
| 19 | A | B | C | D | E |
| 20 | A | B | C | D | E |
| 21 | A | B | C | D | E |
| 22 | A | B | C | D | E |

## SECTION 1

### Question 1

Answer D

$f(x) = g(x) \log_e(2x)$  differentiating using the product rule

$$f'(x) = g'(x) \log_e(2x) + \frac{g(x)}{x}$$

$$f'\left(\frac{e}{2}\right) = g'\left(\frac{e}{2}\right) \log_e(e) + \frac{2}{e} g\left(\frac{e}{2}\right)$$

$$f'\left(\frac{e}{2}\right) = 1 \times 1 + \frac{e}{2} \times \frac{2}{e} = 2$$

### Question 2

Answer A

$2x + (p+1)y = 4$  and  $px + 6y = q$ , in matrix form as  $\begin{bmatrix} 2 & p+1 \\ p & 6 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ q \end{bmatrix}$

The determinant

$$\begin{vmatrix} 2 & p+1 \\ p & 6 \end{vmatrix} = 12 - p(p+1) = 12 - p^2 - p = -(p^2 + p - 12) = -(p-3)(p+4)$$

so when  $p = 3$  and  $p = -4$  there is no unique solution. If  $p = 3$  the equations become

$$(1) \quad 2x + 4y = 4 \quad \text{and} \quad (2) \quad 3x + 6y = q$$

$3 \times (1) \quad 6x + 12y = 12$  and  $2 \times (2) \quad 6x + 12y = 2q$  when  $q = 6$  the two equations are the same equation, so when  $p = 3$  and  $q = 6$  there is an infinite number of solutions.

### Question 3

Answer B

$$f(x) = x^3 + e^{2x} \quad \text{average value is} \quad \frac{1}{2} \int_0^2 (x^3 + e^{2x}) dx = \frac{7 + e^4}{4}$$

### Question 4

Answer E

$$y = x^4 \quad \text{into} \quad y = (3x+6)^4 - 2 \quad \text{or} \quad y + 2 = (3x+6)^4$$

$$y = y' + 2 \quad \text{and} \quad x = 3x' + 6 \quad \text{become}$$

$$x' = \frac{x}{3} - 2 \quad \text{and} \quad y' = y - 2 \quad \text{in matrix form}$$

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} \frac{1}{3} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} -2 \\ -2 \end{bmatrix}$$

**Question 5**

**Answer C**

Let  $y_1 = mx + c$  and  $y_2 = -x^2 + 3x - 3$ , the tangent to the graph at

the point  $P$ , where  $x = 3$ .  $\frac{dy_2}{dx} = -2x + 3 \quad \left. \frac{dy_2}{dx} \right|_{x=3} = -3 = m$  so **B.** is true.

At  $x = 3$   $y_2 = -9 + 9 - 3 = -3$   $P(3, -3)$  is on the tangent,

$y_1 = mx + c \Rightarrow -3 = -9 + c \Rightarrow c = 6$  so **A.** is true, also **D.** is true.

The area  $A = \int_a^b (y_1 - y_2) dx$   $a = 0$   $b = 3$ , so that  $A = \int_0^3 (x^2 + (m-3)x + (c+3)) dx$

**E.** is true, **C.** is false.

**Question 6**

**Answer A**

$f(x+h) \approx f(x) + hf'(x)$  with  $f(x) = \frac{1}{x^2}$   $x = 4$   $h = -0.01$

so that  $\frac{1}{3.99} \approx f(4) - 0.01f'(4)$

**Question 7**

**Answer B**

The required area is below the  $x$ -axis, so taking the absolute value, makes the area

positive. **A.** is true  $\left| \int_{-a}^a (x^2 - a^2) dx \right|$  this is also equal to **D.** which is true  $\int_{-a}^a (a^2 - x^2) dx$ ,

by symmetry **C.** is true  $2 \int_0^a (a^2 - x^2) dx$ . The graph of  $y = x^2 - a^2$ , this crosses the  $y$ -axis

at  $-a^2$ , now the inverse function is  $x = y^2 - a^2 \Rightarrow y^2 = x + a^2 \Rightarrow y = \sqrt{x + a^2}$ , the

area bounded by the curve and the  $y$ -axis is  $2 \left| \int_{-a^2}^0 \sqrt{x + a^2} dx \right|$ , so that **E.** is true, **B.** is false.

**Question 8**

**Answer C**

$f: y = x^2 + a \quad \text{dom } f = \mathbb{R}^- \quad \text{ran } f = (a, \infty)$

$f^{-1} \quad x = y^2 + a$  transposing

$y^2 = x - a \quad y = \pm \sqrt{x - a}$  but  $\text{ran } f^{-1} = \mathbb{R}^- \quad \text{dom } f^{-1} = (a, \infty)$  so we must take the

negative,  $f^{-1}: (a, \infty) \rightarrow \mathbb{R}^-$ ,  $f^{-1}(x) = -\sqrt{x - a}$

**Question 9**

**Answer A**

$f(x) = f(-x)$ ,  $f(x)$  is an even function, and  $\int_{-6}^6 f(x) dx = 10$ , then  $\int_0^6 f(x) dx = 5$

$$\int_0^6 (2f(x) - 1) dx = 2 \int_0^6 f(x) dx - [x]_0^6 = 2 \times 5 - (6 - 0) = 4$$

**Question 10**

**Answer B**

The function is not defined when  $x = 0$ ,  
all of **A**, **C**, **D**. and **E**. are false,  
The function is an even function,  
symmetrical about the y-axis.


$$y = x^2 \log_e(x)$$

$$\frac{dy}{dx} = 2x \log_e(x) + x = x(2 \log_e(x) + 1)$$

for turning points,  $\frac{dy}{dx} = 0$ , since  $x \neq 0$

$$\log_e(x) = -\frac{1}{2} \quad x = e^{-\frac{1}{2}} = \frac{1}{\sqrt{e}},$$

the graph has minimums at  $x = \pm \frac{1}{\sqrt{e}}$



**Question 11**

**Answer E**

$$f: y = b + \frac{a}{x-b}$$

$$f^{-1} \quad x = b + \frac{a}{y-b} \quad \Rightarrow x-b = \frac{a}{y-b} \quad \Rightarrow y-b = \frac{a}{x-b}$$

$$f^{-1}(x) = y = b + \frac{a}{x-b} \quad \text{so} \quad f^{-1} = f$$

The domain and range of both  $f$  and  $f^{-1}$  are  $R \setminus \{b\}$ .

Since  $a \neq 0$  and  $b \neq 0$ , the graph of  $y = f(x)$  passes through  $\left(0, b - \frac{a}{b}\right)$

and the graph of  $y = f^{-1}(x)$  passes through  $\left(b - \frac{a}{b}, 0\right)$ .

All of **A**, **B**, **C**, **D**. are true, however **E**. is false

The graph of  $y = f(x)$  and  $y = f^{-1}(x)$  always intersects on the line  $y = x$  at the points  $(b \pm \sqrt{a}, b \pm \sqrt{a})$  only if  $a > 0$ .

**Question 12**

**Answer C**

$$\frac{dy}{dx} = 2 \cos\left(\frac{x}{2}\right) \Rightarrow y = \int 2 \cos\left(\frac{x}{2}\right) dx = 4 \sin\left(\frac{x}{2}\right) + c \quad \text{to find } c, \text{ use } y\left(\frac{5\pi}{3}\right) = 0$$

$$0 = 4 \sin\left(\frac{5\pi}{6}\right) + c = 2 + c = 0 \Rightarrow c = -2$$

$$y = 4 \sin\left(\frac{x}{2}\right) - 2 \quad \text{now when } x = 0 \quad y = 4 \sin(0) - 2 = -2$$

**Question 13**

**Answer D**

$y = \frac{bx}{x-a} = \frac{bx-ab+ab}{x-a} = b + \frac{ab}{x-a}$  has  $y = b$  as a horizontal asymptote and  $x = a$  as a vertical asymptote.

**Question 14**

**Answer D**

Let  $f: [0, \pi] \rightarrow R$ ,  $f(x) = 2 \cos\left(\frac{x}{2}\right) - 2$ . The period is  $T = \frac{2\pi}{\frac{1}{2}} = 4\pi$

The graph of  $f$  is transformed by a reflection in the  $x$ -axis, the rule is

$g(x) = 2 - 2 \cos\left(\frac{x}{2}\right)$ , we only have one-quarter of a cycle

now a dilation of factor 2 from the  $y$ -axis, replace  $x$  with  $\frac{x}{2}$

$g: [0, 2\pi] \rightarrow R$ ,  $g(x) = 2 - 2 \cos\left(\frac{x}{4}\right)$  since we must have one-quarter of a cycle,

the new domain is  $[0, 2\pi]$

then a dilation by a factor of 3 from the  $x$ -axis, multiply  $y$  by 3

the equation becomes  $g: [0, 2\pi] \rightarrow R$ ,  $g(x) = 6 - 6 \cos\left(\frac{x}{4}\right)$

**Question 15**

**Answer E**

$$\Pr(A' \cap B') + b - p = 1 - a \quad \text{or}$$

$$\Pr(A' \cap B') + a - p = 1 - b$$

$$\Pr(A' \cap B') = 1 + p - (a + b)$$

|      |         |         |         |
|------|---------|---------|---------|
|      | $A$     | $A'$    |         |
| $B$  | $p$     | $b - p$ | $b$     |
| $B'$ | $a - p$ | $?$     | $1 - b$ |
|      | $a$     | $1 - a$ |         |



**Question 16**

**Answer D**

$$f(x) = x^4 - 4x^2$$

$$f'(x) = 4x^3 - 8x$$


$$f'(x) = 4x(x^2 - 2)$$

turning points at

$$x = 0, x = \pm\sqrt{2}$$

for the function to be one-one, we require

$$a < -\sqrt{2}$$



**Question 17**

**Answer B**

$X \stackrel{d}{=} Bi(n=?, p=0.6)$  Betty winning a game.

$$\Pr(X = 0) = 0.4^n \leq 0.01$$

$$n \log_e(0.4) \leq \log_e(0.01)$$

$$n \geq \frac{\log_e(0.01)}{\log_e(0.4)} = 5.02 \quad \text{so } n = 6$$

**Question 18**

**Answer B**

$$X \stackrel{d}{=} N\left(\mu_X = \mu, \sigma_X^2 = \frac{9\mu^2}{4}\right)$$

$$\Pr(X > 2\mu) = \Pr\left(Z > \frac{2\mu - \mu}{\frac{3}{2}\mu}\right) = \Pr\left(Z > \frac{2}{3}\right) = 0.252$$

**Question 19**

**Answer C**

$$\text{Let } g(x) = \int_0^x f(t) dt \text{ then } g'(x) = f(x)$$

$$g(0) = 0 \quad g'(0) = f(0) = 9 \quad g'(3) = f(3) = 0$$

**Question 20**

**Answer D**

Option **D.** has  $f(x) = e^x$   $g(x) = -x^2$  and  $f(g(x)) = e^{-x^2}$   
Which is the graph required.

**Question 21**

**Answer A**

$$A = \frac{3\sqrt{3}}{2} L^2 \quad \frac{dA}{dL} = 3\sqrt{3}L \quad \text{given} \quad \frac{dL}{dt} = \sqrt{3} \text{ cm/s}$$

$$\frac{dA}{dt} = \frac{dA}{dL} \cdot \frac{dL}{dt} = 3\sqrt{3}L \times \sqrt{3} = 9L$$

$$\left. \frac{dA}{dt} \right|_{L=\sqrt{3}} = 9\sqrt{3} \text{ cm}^2/\text{s}$$

**Question 22**

**Answer E**

$$\text{Since } \sum \Pr(X = x) = 1 \Rightarrow \frac{a}{2} + a + b + \frac{b}{2} = \frac{3a}{2} + \frac{3b}{2} = 1 \Rightarrow 3(a+b) = 2 \text{ **A.** is true}$$

$$E(X) = \sum x \Pr(X = x) = -2 \times \frac{a}{2} - a + b + 2 \times \frac{b}{2} = -2a + 2b = 2(b-a) \text{ **B.** is true}$$

$$E(X^2) = \sum x^2 \Pr(X = x) = (-2)^2 \times \frac{a}{2} + (-1)^2 a + (1)^2 b + (2)^2 \times \frac{b}{2} = 2a + a + b + 2b = 3(a+b) = 2$$

**C.** is true, since **A.** is true.

$$\text{var}(X) = E(X^2) - (E(X))^2 = 2 - 4(b-a)^2 = 2 - 4b^2 + 8ab - 4a^2 \text{ **D.** is true}$$

$$\text{E. is false, } E\left(\frac{1}{X}\right) = \sum \frac{1}{x} \Pr(X = x) = -\frac{a}{4} - a + b + \frac{b}{4} = \frac{5}{4}(b-a)$$

**END OF SECTION 1 SUGGESTED ANSWERS**

## SECTION 2

### Question 1

- a.i**  $f(x) = x^3 - 3x^2 + cx + d$   
 $f'(x) = 3x^2 - 6x + c$   
 but  $x = -1$  is a turning point so  $f'(x) = (x+1)(3x-9) = 3(x+1)(x-3)$   
 Expanding gives  $c = -9$ ,  
 also  $u = 3$ , A1  
 $f(-1) = 5 = -1 - 3 - c + d = -4 + 9 + d$  so that  
 $d = 0$  and  $f(3) = v = 27 - 27 - 27 = -27$   $v = -27$  A1
- ii.** The graph of  $y = x^3 - 3x^2 - 9x$  has a maximum value of 5, and a minimum value of  $-27$ , and crosses the x-axis at three distinct points. The graph of  $y = x^3 - 3x^2 - 9x + d$  will therefore cross the x-axis at three distinct points, provided that  $d \in (-5, 27)$  or  $-5 < d < 27$  A2
- b.**  $f(x) = x^3 - 3x^2 + cx + d$   
 $f'(x) = 3x^2 - 6x + c$ , for two distinct turning points, we require  
 $\Delta = 36 - 12c > 0$  M1  
 $c < 3$  and  $d \in R$  A1
- c.**  $f(x+p) = (x+p)^3 - 3(x+p)^2 + c(x+p) + d$   
 $f(x+p) = x^3 + x^2(3p-3) + x(3p^2-6p+c) + p^3-3p^2+cp+d = x^3$   
 therefore  $3p-3=0 \Rightarrow p=1$   
 and  $3p^2-6p+c=0$  since  $p=1$   $c=6-3=3$   
 $c=3$  A1  
 and  $p^3-3p^2+cp+d=0$   
 so  $d=-1$  A1  
 alternative method, if  $y = (x-1)^3 = x^3 - 3x^2 + 3x - 1 \rightarrow y = x^3$   
 so that  $p=1$   $c=3$  and  $d=-1$

d.  $A = \int_a^b f(x) dx$   $a = 0$   $b = 2$   $h = \frac{1}{2}$   $n = 4$   $f(x) = x^3 - 3x^2 + cx + d$

(1)  $L = 10 = h \left[ f(0) + f\left(\frac{1}{2}\right) + f(1) + f\left(\frac{3}{2}\right) \right]$  M1

(2)  $R = 6 = h \left[ f\left(\frac{1}{2}\right) + f(1) + f\left(\frac{3}{2}\right) + f(2) \right]$

(1)  $\Rightarrow 20 = f(0) + f\left(\frac{1}{2}\right) + f(1) + f\left(\frac{3}{2}\right)$

(2)  $\Rightarrow 12 = f\left(\frac{1}{2}\right) + f(1) + f\left(\frac{3}{2}\right) + f(2)$  subtracting gives M1

$8 = f(0) - f(2) = d - (8 - 12 + 2c + d) = 4 - 2c$

$2c = -4$

$c = -2$  A1

Now  $\int_0^2 (x^3 - 3x^2 - 2x + d) = 2d - 8 = 8$  A1

$d = 8$  A1

## Question 2

a. the amplitude is 1.5, so that  $a = 1.5$

one-half cycle is 8, so that  $T = \frac{2\pi}{n} = 16 \Rightarrow n = \frac{\pi}{8}$  A1

b.  $y = 16(1 - e^{-kx})$  passes through the origin  $O(0,0)$  and  $B(4,8)$

$8 = 16(1 - e^{-4k}) \Rightarrow \frac{1}{2} = 1 - e^{-4k}$

$e^{-4k} = \frac{1}{2} \quad e^{4k} = 2$

$4k = \log_e(2)$

$k = \frac{1}{4} \log_e(2)$  A1

**c.i** reflect in the y-axis  
 translate 8 units, to the right, away from the y-axis A1  
 or translate 8 units, to the right parallel to the x-axis.

**ii.**  $f : [4, 8] \rightarrow R$  ,  $f(x) = 16(1 - e^{k(x-8)})$  A2  
 must give domain.

**d.i**  $A = 2 \int_0^4 \left( 16(1 - e^{-kx}) - \frac{3}{2} \sin\left(\frac{\pi x}{8}\right) \right) dx$  A1

**ii.**  $A = 2 \left[ 16x + \frac{16}{k} e^{-kx} + \frac{12}{\pi} \cos\left(\frac{\pi x}{8}\right) \right]_0^4$  each term A1

$A = 2 \left[ \left( 64 + \frac{16}{k} e^{-4k} + \frac{12}{\pi} \cos\left(\frac{\pi}{2}\right) \right) - \left( 0 + \frac{16}{k} + \frac{12}{\pi} \cos(0) \right) \right]$  but  $e^{-4k} = \frac{1}{2}$  M1

$A = 2 \left[ 64 + \frac{8}{k} - \frac{16}{k} - \frac{12}{\pi} \right]$

$A = 128 - \frac{16}{k} - \frac{24}{\pi}$

$p = 128$   $q = -16$  and  $r = -24$  A2

### Question 3

**a.** the function is continuous  $f(4) = 16b = 4c \Rightarrow c = 4b$  A1  
 the total area under the curve is one.


$b \int_0^4 t^2 dt + c \int_4^8 (8-t) dt = 1$  A1

$\frac{64b}{3} + 8c = 1$  and  $c = 4b$  A1

solving gives  $b = \frac{3}{160}$  and  $c = \frac{3}{40}$

- b. must show point at  $(4, 0.3)$  and zero for  $t \geq 8$  and  $t \leq 0$

G1



c.  $\Pr(T > 6) = \frac{3}{40} \int_6^8 (8-t) dt$  or the area of a triangle as

M1

$$\Pr(T > 6) = \frac{1}{2} \times 2 \times 2c = \frac{3}{20}$$

$$\Pr(T > 6) = \frac{3}{20}$$

A1

d.  $E(T) = \frac{3}{160} \int_0^4 t^3 dt + \frac{3}{40} \int_4^8 t(8-t) dt$

M1

$$E(T) = 1.2 + 3.2 = 4.4 \text{ minutes}$$

A1

e. Since  $\frac{3}{160} \int_0^4 t^2 dt = 0.4$  the median time  $m$  is given by

$$\frac{3}{40} \int_4^m (8-t) dt = 0.1 \quad \text{M1}$$

$$\frac{-3(m^2 - 16m + 48)}{80} = 0.1$$

solving for  $m$  with  $4 < m < 8$

$$m = 4.35 \text{ minutes} \quad \text{A1}$$

f.  $X$  is the running time of the movie in minutes

$$X \stackrel{d}{=} N(\mu = 94, \sigma^2 = 10^2)$$

$$\Pr(X > 109) = \Pr\left(Z > \frac{109 - 94}{10}\right) = \Pr(Z > 1.5)$$

$$= 0.0668 \quad \text{A1}$$

g.  $Y \stackrel{d}{=} Bi(n = 4, p = 0.0668)$

$$\Pr(Y \geq 2) = 1 - [\Pr(Y = 0) + \Pr(Y = 1)]$$

$$\Pr(Y \geq 2) = 1 - [0.9332^4 + {}^4C_1 0.0668 \times 0.9332^3] \quad \text{M1}$$

$$\Pr(Y \geq 2) = 0.0244 \quad \text{A1}$$

h.  $\Pr(2 \text{ comedies}) = ACC + CAC + CCA$  M1

$$= 0.45 \times 0.55 \times 0.65 + 0.55 \times 0.35 \times 0.55 + 0.55 \times 0.65 \times 0.35$$

$$= 0.392 \quad \text{A1}$$

i.  $\frac{0.35}{0.35 + 0.55} = 0.389$

or alternatively

|   |      |      |
|---|------|------|
|   | A    | C    |
| A | 0.45 | 0.35 |
| C | 0.55 | 0.65 |

$$\begin{bmatrix} 0.45 & 0.35 \\ 0.55 & 0.65 \end{bmatrix}^{-100} = \begin{bmatrix} 0.389 & 0.389 \\ 0.611 & 0.611 \end{bmatrix}$$

in the long run, the percentage of movies which are actions are 38.9% A1

**Question 4**

**a.**  $P\left(a, \frac{4}{a^2}\right) O(0,0)$

$$s = d(OP) = \sqrt{(a-0)^2 + \left(\frac{4}{a^2} - 0\right)^2} \quad \text{M1}$$

$$s = \sqrt{a^2 + \frac{16}{a^4}} = \sqrt{\frac{16 + a^6}{a^4}} \quad \text{since } a > 0$$

$$s = \frac{1}{a^2} \sqrt{16 + a^6} \quad \text{A1}$$

**b.i.**  $\frac{ds}{da} = \frac{a^6 - 32}{a^3 \sqrt{16 + a^6}} = 0$  for minimum distance A1

$$a = \sqrt[6]{32} = 2^{\frac{5}{6}} \quad \text{A1}$$

if  $a > 2^{\frac{5}{6}}$  consider  $a = 1.8$   $\frac{ds}{da} = 0.049 > 0$

and if  $a < 2^{\frac{5}{6}}$  consider  $a = 1.7$   $\frac{ds}{da} = -0.25 < 0$

by the sign test it is a minimum. M1

**ii.**  $S_{\min} = \sqrt[3]{2} \cdot \sqrt{3}$  A1

**c.i** at the point  $P\left(a, \frac{4}{a^2}\right)$   $f'(x) = -8x^{-3}$   $m_T = -\frac{8}{a^3}$  A1

$$m_N = \frac{a^3}{8}$$

normal  $y - \frac{4}{a^2} = \frac{a^3}{8}(x - a)$  or  $y = \frac{a^3 x}{8} - \frac{a^4}{8} + \frac{4}{a^2}$  A1

**ii.** normal passes through origin (0,0) then

$$-\frac{a^4}{8} + \frac{4}{a^2} = 0$$


$$a^6 = 32$$

$$a = \sqrt[6]{32} = 2^{\frac{5}{6}} \quad \text{A1}$$



**Question 5**

- a.**  $\sqrt{3} \sin(2x) + \cos(2x) = 0$   
 $x = \frac{(6k-1)\pi}{12}$  where  $k \in \mathbb{Z}$  A1
- b.** since the period of both are  $\pi$ , it follows that  $g(x + \pi) = g(x)$   
 $T = \pi$  A1
- c.**  $f : [0, 2\pi] \rightarrow \mathbb{R}$ ,  $f(x) = \sqrt{3} \sin(2x) + \cos(2x)$   
 $\sqrt{3} \sin(2x) + \cos(2x) = 0$   
 $\sqrt{3} \sin(2x) = -\cos(2x)$   
 $\tan(2x) = -\frac{1}{\sqrt{3}}$   
 $x = \frac{5\pi}{12}, \frac{11\pi}{12}, \frac{17\pi}{12}, \frac{23\pi}{12}$  A1
- d.i.**  $f'(x) = 2\sqrt{3} \cos(2x) - 2 \sin(2x) = 0$  A1  
 $\sqrt{3} \cos(2x) = \sin(2x)$   
 $\tan(2x) = \sqrt{3}$   
 $x = \frac{\pi}{6}, \frac{2\pi}{3}, \frac{7\pi}{6}, \frac{5\pi}{3}$  A1
- ii.**  $\max \left( \frac{\pi}{6}, 2 \right)$  and  $\left( \frac{7\pi}{6}, 2 \right)$ ,  $\min \left( \frac{2\pi}{3}, -2 \right)$  and  $\left( \frac{5\pi}{3}, -2 \right)$  A1
- e.** graph on correct domain, correct x-intercepts G1  
 and correct max and min. G1



f. 
$$f(x) = \sqrt{3} \sin(2x) + \cos(2x) = 2 \sin\left(2x + \frac{\pi}{6}\right) = 2 \sin\left(2\left(x + \frac{\pi}{12}\right)\right)$$

translate  $2 \sin(2x)$ ,  $\frac{\pi}{12}$  to the left parallel to the x-axis

$$A = 2$$

A1

$$\alpha = \frac{\pi}{12}$$

A1

**END OF SECTION 2 SUGGESTED ANSWERS**