

**THE
HEFFERNAN
GROUP**

P.O. Box 1180
Surrey Hills North VIC 3127
Phone 03 9836 5021
Fax 03 9836 5025
info@theheffernangroup.com.au
www.theheffernangroup.com.au

Student Name.....

MATHEMATICAL METHODS (CAS) UNITS 3 & 4
TRIAL EXAMINATION 1

2012

Reading Time: 15 minutes

Writing time: 1 hour

Instructions to students

This exam consists of 11 questions.
All questions should be answered in the spaces provided.
There is a total of 40 marks available.
The marks allocated to each of the questions are indicated throughout.
Students may **not** bring any calculators or notes into the exam.
Where an exact answer is required a decimal approximation will not be accepted.
Where more than one mark is allocated to a question, appropriate working must be shown.
Diagrams in this trial exam are not drawn to scale.
A formula sheet can be found on page 11 of this exam.

This paper has been prepared independently of the Victorian Curriculum and Assessment Authority to provide additional exam preparation for students. Although references have been reproduced with permission of the Victorian Curriculum and Assessment Authority, the publication is in no way connected with or endorsed by the Victorian Curriculum and Assessment Authority.

© THE HEFFERNAN GROUP 2012

This Trial Exam is licensed on a non transferable basis to the purchasing school. It may be copied by the school which has purchased it. This license does not permit distribution or copying of this Trial Exam by any other party.

Question 1

- a. For $y = \frac{e^{2x}}{x^2 + 3}$ find $\frac{dy}{dx}$.

1 mark

- b. Let $f(x) = 2x \tan\left(\frac{x}{3}\right)$. Find $f'(\pi)$.

2 marks

Question 2

- a. Find an antiderivative of $\sin(3x - 1)$ with respect to x .

1 mark

- b. Evaluate $\int_0^2 \frac{1}{2x+1} dx$.

2 marks

Question 3

Let $f : R^+ \rightarrow R, f(x) = \log_e(x)$.

- a.** Show that $f\left(\frac{1}{u}\right) = -f(u)$ where $u \in R^+$.

1 mark

- b.** If $2f(u) = f(2v) + f(3v)$ where $u, v \in R^+$, then express u in terms of v .

2 marks

Question 4

- a. Find the period and the range of the function $g : \mathbb{R} \rightarrow \mathbb{R}, g(x) = 3 + \sin\left(\frac{2x}{3}\right)$.

2 marks

- b. Solve the equation $\cos(3x) = \frac{\sqrt{3}}{2}$ for $x \in \mathbb{R}$.

3 marks

Question 5

The discrete random variable X has a probability distribution given by

x	0	1	2	3
$\Pr(X = x)$	0.2	0.1	0.4	p

- a.** Find the value of p .

1 mark

- b.** Find $\Pr(X \leq 1 | X < 3)$.

2 marks

- c.** Find $E(X)$, the mean of X .

2 marks

Question 6

Let $f : \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = 1 - e^{2(x-1)}$.

Find f^{-1} , the inverse function of f .

3 marks

Question 8

For two events A and B , $\Pr(A) = \frac{1}{3}$ and $\Pr(B) = \frac{1}{2}$.

Find $\Pr(A \cup B')$ if

- a.** A and B are mutually exclusive.

2 marks

- b.** A and B are independent events.

2 marks

Question 9

June shops either at the market or locally every week.

If she shops at the market one week the probability that she shops there the following week is 0.6. If she shops locally one week then the probability that she shops at the market the following week is 0.7.

This week June shopped at the market.

What is the probability that June will shop locally exactly once in the coming three weeks?

3 marks

Question 10

A normal to the graph of $y = x^{\frac{3}{5}} + c$ at the point where $x = 1$ has an x-intercept at $(a, 0)$ where a and c are both non-zero real constants.
Find an expression for a in terms of c .

3 marks

Mathematical Methods (CAS) Formulas

Mensuration

area of a trapezium:	$\frac{1}{2}(a+b)h$	volume of a pyramid:	$\frac{1}{3}Ah$
curved surface area of a cylinder:	$2\pi rh$	volume of a sphere:	$\frac{4}{3}\pi r^3$
volume of a cylinder:	$\pi r^2 h$	area of a triangle:	$\frac{1}{2}bc \sin A$
volume of a cone:	$\frac{1}{3}\pi r^2 h$		

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1}x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a}e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e x + c$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a}\cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a}\sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$	
product rule: $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$	quotient rule: $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$	approximation: $f(x+h) \approx f(x) + hf'(x)$

Probability

$\Pr(A) = 1 - \Pr(A')$	$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$
$\Pr(A B) = \frac{\Pr(A \cap B)}{\Pr(B)}$	transition matrices: $S_n = T^n \times S_0$
mean: $\mu = E(X)$	variance: $\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

probability distribution		mean	variance
discrete	$\Pr(X = x) = p(x)$	$\mu = \sum x p(x)$	$\sigma^2 = \sum (x - \mu)^2 p(x)$
continuous	$\Pr(a < X < b) = \int_a^b f(x) dx$	$\mu = \int_{-\infty}^{\infty} x f(x) dx$	$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$

Reproduced with permission of the Victorian Curriculum and Assessment Authority, Victoria, Australia.

This formula sheet has been copied in 2012 from the VCAA website www.vcaa.vic.edu.au
The VCAA publish an exam issue supplement to the VCAA bulletin.