

**The Mathematical Association of Victoria**

**Trial Exam 2012**

**MATHEMATICAL METHODS (CAS)**

**WRITTEN EXAMINATION 1**

**STUDENT NAME** \_\_\_\_\_

**Reading time: 15 minutes**

**Writing time: 1 hour**

**QUESTION AND ANSWER BOOK**

**Structure of book**

| <i>Number of questions</i> | <i>Number of questions to be answered</i> | <i>Number of marks</i> |
|----------------------------|-------------------------------------------|------------------------|
| 11                         | 11                                        | 40                     |

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers.
- Students are NOT permitted to bring into the examination room: notes of any kind, blank sheets of paper, white out liquid/tape or a calculator of any type.

**Materials supplied**

- Question and answer book of 9 pages, with a detachable sheet of miscellaneous formulas at the back
- Working space is provided throughout the book.

**Instructions**

- Detach the formula sheet from the back of this book during reading time.
- Write your name in the space provided above on this page.
- All written responses must be in English.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.**

**This page is blank**

**Instructions**

Answer **all** questions in the spaces provided.

In all questions where a numerical answer is required an exact value must be given unless otherwise specified.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

**Question 1**

Let  $g(x) = \cos(x)$  and  $f(x) = \log_e(2x - 2)$ .

- a. State the maximal domain of  $f$ .

\_\_\_\_\_ 1 mark

- b. Find the rule for  $g(f(x))$ .


\_\_\_\_\_ 1 mark

- c. Find  $g'(f(x))f'(x)$ .

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_ 2 marks

**Question 2**

Sketch the graph of  $y = \left(\frac{1}{4}\right)^{x-1} - 1$  on the set of axes below. Clearly label any asymptotes with their equations and axes-intercepts with their coordinates.



3 marks

**Question 3**

Solve the equation  $2\log_e(x-2) - \log_e(x+1) = \log_e(2)$  for  $x$ .

---

---

---

---

---

---

---

---

---

---

3 marks

**Question 4**

a. Differentiate  $(x+2)\sqrt{x-1}$ , giving your answer as a single fraction.

---

---

---

---

---

---

---

3 marks

b. Hence, find an antiderivative of  $\frac{x}{\sqrt{x-1}}$ .

---

---

---

2 marks



**Question 6**

If  $h(x) = e^{(x-1)}$ , show that  $h(x+y) \times h(x-y) = (h(x))^2$ .

---

---

---

---


---

---

2 marks

**Question 7**

On the axes provided, sketch the graph of the function  $f : [-\pi, 3\pi] \rightarrow \mathbb{R}$  where  $f(x) = \frac{1}{2} \sin\left(\frac{x}{2}\right)$ . Label the endpoints with their coordinates.



2 marks

**Question 8**

Find the values of  $a \in \left[0, \frac{2}{3}\right]$  for which  $x = \frac{1}{6}$  is the solution to the equation  $2 \sin(3\pi(x - a)) = \sqrt{3}$ .

---

---

---

---

---

---

---

---

3 marks

**Question 9**

If  $X \sim \text{Bi}(5, 0.1)$ , find  $\Pr(X = 3)$ .

---

---

---

2 marks

**Question 10**

The probability density function of a random variable  $X$  has a density function given by

$$f(x) = \begin{cases} |x-1| & 0 \leq x \leq 1 \\ 0.1 & 1 < x \leq a \\ 0 & \text{elsewhere} \end{cases}$$

a. Find the value of  $a$ .

---

---

---

---

1 mark

- b. Find  $\Pr(X > 0.5)$ .

---



---



---

1 mark

- c. Find  $\Pr(X > 2 \mid X > 0.5)$ .

---



---



---



---



---




---

2 marks

### Question 11

The diagram represents two sails of a toy yacht, where triangle  $ABC$  and triangle  $BCD$  are the two sails. The length of  $AC$  is  $a$  metres and  $\angle BAC = \theta^\circ$ .

Also  $AB = 2BD$  and  $\angle CBD = \angle CBA = \frac{\pi}{2}$ .



- a. Write  $BC$  and  $AD$  in terms of  $a$  and  $\theta$ .

---



---



---

2 marks

- b.** Write the total sail area,  $T \text{ m}^2$ , in terms of  $a$  and  $\theta$ .

---

---

---

1 mark

- c. i.** Find  $\frac{dT}{d\theta}$ .

---

---

---

- ii. Hence,** find the maximum total sail area, when  $a = 4 \text{ m}$ .

---

---

---

---

---

---

---

---

---

1 + 3 = 4 marks

**END OF QUESTION AND ANSWER BOOK**

# **MATHEMATICAL METHODS (CAS)**

## **Written examinations 1 and 2**

### **FORMULA SHEET**

#### **Directions to students**

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

**This page is blank**

## Mathematical Methods (CAS)

### Formulas

#### Mensuration

|                                    |                        |                      |                        |
|------------------------------------|------------------------|----------------------|------------------------|
| area of a trapezium:               | $\frac{1}{2}(a+b)h$    | volume of a pyramid: | $\frac{1}{3}Ah$        |
| curved surface area of a cylinder: | $2\pi rh$              | volume of a sphere:  | $\frac{4}{3}\pi r^3$   |
| volume of a cylinder:              | $\pi r^2 h$            | area of a triangle:  | $\frac{1}{2}bc \sin A$ |
| volume of a cone:                  | $\frac{1}{3}\pi r^2 h$ |                      |                        |

#### Calculus

|                                                                |                                                     |
|----------------------------------------------------------------|-----------------------------------------------------|
| $\frac{d}{dx}(x^n) = nx^{n-1}$                                 | $\int x^n dx = \frac{1}{n+1}x^{n+1} + c, n \neq -1$ |
| $\frac{d}{dx}(e^{ax}) = ae^{ax}$                               | $\int e^{ax} dx = \frac{1}{a}e^{ax} + c$            |
| $\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$                        | $\int \frac{1}{x} dx = \log_e x  + c$               |
| $\frac{d}{dx}(\sin(ax)) = a \cos(ax)$                          | $\int \sin(ax) dx = -\frac{1}{a}\cos(ax) + c$       |
| $\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$                         | $\int \cos(ax) dx = \frac{1}{a}\sin(ax) + c$        |
| $\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$ |                                                     |

product rule:  $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$

quotient rule:  $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

chain rule:  $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

approximation:  $f(x+h) \approx f(x) + hf'(x)$

#### Probability

$\Pr(A) = 1 - \Pr(A')$

$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$

$\Pr(A|B) = \frac{\Pr(A \cap B)}{\Pr(B)}$

transition matrices:  $S_n = T^n \times S_0$

mean:  $\mu = E(X)$

variance:  $\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

| probability distribution |                                     | mean                                      | variance                                                 |
|--------------------------|-------------------------------------|-------------------------------------------|----------------------------------------------------------|
| discrete                 | $\Pr(X = x) = p(x)$                 | $\mu = \sum x p(x)$                       | $\sigma^2 = \sum (x - \mu)^2 p(x)$                       |
| continuous               | $\Pr(a < X < b) = \int_a^b f(x) dx$ | $\mu = \int_{-\infty}^{\infty} x f(x) dx$ | $\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$ |

END OF FORMULA SHEET