

insightTM

YEAR 12 *Trial Exam Paper*

2014

MATHEMATICAL METHODS (CAS)

Written examination 2

Worked solutions

This book presents:

- worked solutions, giving you a series of points to show you how to work through the questions
- mark allocations
- tips on how to approach the questions

This trial examination produced by Insight Publications is NOT an official VCAA paper for the 2014 Mathematical Methods (CAS) Written examination 2.

The Publishers assume no legal liability for the opinions, ideas or statements contained in this Trial Exam.

This examination paper is licensed to be printed, photocopied or placed on the school intranet and used only within the confines of the purchasing school for examining their students. No trial examination or part thereof may be issued or passed on to any other party including other schools, practising or non-practising teachers, tutors, parents, websites or publishing agencies without the written consent of Insight Publications.

Copyright © Insight Publications 2014.

SECTION 1

Question 1

Answer is D

Worked solution

The range is given by $[5 - 2, 5 + 2] = [3, 7]$.


The period is given by $\frac{2\pi}{4\pi} = \frac{1}{2}$.

Question 2

Answer is D.

Worked solution

Using CAS, the range is $(-\infty, 2]$.



Tip

- To find the range always draw the graph.

Question 3**Answer is E.****Worked solution**

For $f(x)$ to be defined, $x > k$, so the domain is (k, ∞) .

And $\sqrt{x-k} > 0$, so the range is R^+ .

**Tip**

- If it helps, an arbitrary value of k can be chosen and then a graph produced.

Question 4**Answer is D.****Worked solution**


Using the chain rule for $y = g(f(x))$, $\frac{dy}{dx} = g'(f(x)) \times f'(x)$.

Now $f(x) = \sin(5x)$ and $f'(x) = 5 \cos(5x)$, so $\frac{d}{dx}[g(\sin(5x))] = g'(\sin(5x)) \times 5 \cos(5x)$.

Question 5**Answer is B.****Worked solution**

The graph is a tan graph with a period of $\frac{\pi}{2}$ that has been shifted to the left $\frac{\pi}{4}$ units.

Checking using CAS gives




Question 6*Answer is A.***Worked solution**

$$f(3) = 27 - 27 = 0$$

$$f(1) = 1 - 3 = -2$$

$$\text{Average rate of change} = \frac{f(3) - f(1)}{3 - 1} = \frac{0 - (-2)}{2} = 1$$

$$\text{Average value of the function is } \frac{1}{3-1} \int_1^3 x^3 - 3x^2 \, dx = -3 \text{ (found using CAS).}$$


**Tip**

- Be careful! Option E is the answer to the **average value** of the function—this is easily and readily confused.

Question 7**Answer is A.****Worked solution**

For an inverse function to exist, the function $f(x)$ must be one-to-one; i.e. for each x value there is exactly one y value.

The graph of $f(x)$ is



To be one-to-one $x \in [0, \infty)$.

**Tip**

- For the graph to be one-to-one it must pass the horizontal line test.

Question 8

Answer is D.

Worked solution

Expanding the matrix gives

The screenshot shows a TI-84 Plus calculator interface. The top screen displays the expression:

$$\begin{bmatrix} 0 & 1 & 1 \\ -1 & 0 & 1 \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

The bottom screen shows the result:

$$\begin{bmatrix} y+z \\ -x+z \\ z \end{bmatrix}$$

The calculator interface includes a top menu bar with 'Edit', 'Action', and 'Interactive' options. Below this is a toolbar with various mathematical functions like '0.5 1/2', '1/2', 'f dx/dx', 'Simp', 'f dx/dx', and 'f dx/dx'. The bottom of the screen shows a keypad with categories like 'Math1', 'Math2', 'Math3', 'Trig', 'Var', 'abc', and 'Alg'. The 'Alg' category is currently selected, showing options like 'Standard', 'Real', and 'Rad'.



Tip

- Ensure the equations are first re-written in the form


$$0x + 0y + z = 1$$

$$-1x + 0y + z = 2$$

$$0x + y + z = 5$$

Question 9*Answer is C.***Worked solution**


Using CAS gives



Question 10*Answer is E.***Worked solution**

A suitable way to do this question is to choose an arbitrary value for a ; e.g. $a = 4$.

A sample graph would be



Owing to the condition that $u < t$, the gradient is positive for $x \in (-\infty, u) \cup (0, t)$.

Question 11**Answer is A.****Worked solution**

For this approximation $h = 0.2$, $x = 9$ and $f'(x) = \frac{3}{2}\sqrt{x}$.

So $f(9+h) \approx f(9) + hf'(9)$

$$\begin{aligned} &= \left(9^{\frac{1}{2}}\right)^3 + 0.2 \times \frac{3}{2}\sqrt{9} \\ &= 3^3 + 0.2 \times \frac{9}{2} = 27.9 \end{aligned}$$

Question 12**Answer is A.****Worked solution**

Rearranging the equation $y = 1 - 2\sin(3x + \pi)$ gives $\frac{y-1}{-2} = \sin\left(3\left(x + \frac{\pi}{3}\right)\right)$.

So $y = \frac{y'-1}{-2}$ and $x = 3\left(x' + \frac{\pi}{3}\right)$.

Therefore, $y' = -2y + 1$ and $x' = \frac{x}{3} - \frac{\pi}{3}$.

The expansion of the matrix in option A; i.e. $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{1}{3} & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} -\frac{\pi}{3} \\ 1 \end{bmatrix}$, gives

$x' = \frac{x}{3} - \frac{\pi}{3}$ and $y' = -2y + 1$, as required.

Question 13*Answer is E.***Worked solution**

$$X \sim \text{Bi}$$

$$E(X) = np = 20$$

$$\text{var}(X) = npq = 12 = 20q$$

$$\text{So } q = \frac{12}{20} = 0.6 \Rightarrow p = 0.4$$

$$\therefore n = \frac{20}{0.4} = 50$$

Question 14*Answer is A.***Worked solution**

$$\sum \Pr(X = x) = 1 \Rightarrow 10a = 1, \therefore a = 0.1$$

$$E(X) = \sum x \Pr(X = x)$$

$$= 1 \times 0.2 + 2 \times 0.3 + 3 \times 0.1$$

$$= 1.1$$

Question 15***Answer is D.*****Worked solution**

The transformation represents a reflection in the y -axis and a translation of 2 units up.
 Operating this transformation on the tangent line gives

$$y = 2(-x) - 5 + 2$$

$$y = -2x - 3$$

Question 16***Answer is B.*****Worked solution**

Using the given information

$$\int_{-4}^5 f(x) dx = 4$$

$$\Rightarrow F(5) - F(-4) = 4 \quad (1)$$

$$\begin{aligned} \text{So } \int_{-1}^2 (f(3x-1)) dx &= \frac{1}{3} (F(3(2)-1) - F(3(-1)-1)) \\ &= \frac{1}{3} (F(5) - F(-4)) \\ &= \frac{1}{3} (4) \quad (\text{from equation 1}) \\ &= \frac{4}{3} \end{aligned}$$

Question 17*Answer is C.***Worked solution**

$$\Pr(G' | G) = 0.65 \text{ and } \Pr(G' | B) = 0.3$$

So the matrix is set up as
$$\begin{bmatrix} \Pr(G' | G) & \Pr(G' | B) \\ \Pr(B' | G) & \Pr(B' | B) \end{bmatrix} = \begin{bmatrix} 0.65 & 0.3 \\ 0.35 & 0.7 \end{bmatrix}.$$

Question 18*Answer is C.***Worked solution**

$$\begin{aligned} [f(u)]^2 - 2 &= (e^x + e^{-x})^2 - 2 \\ &= e^{2x} + 2 + e^{-2x} - 2 \\ &= e^{2x} + e^{-2x} \\ &= f(2u) \end{aligned}$$

Question 19*Answer is E.***Worked solution**


$$\Pr(X > 71) = \Pr\left(Z > \frac{71-50}{7}\right) = \Pr(Z > 3)$$

Using symmetry, $\Pr(Z > 3) = \Pr(Z < -3)$.

Question 20*Answer is E.***Worked solution**

First, choose a value for θ , say $\theta = \frac{\pi}{6}$.

Using CAS gives




Question 21

Answer is E.

Worked solution


The graph of $y = -\log_e(x)$ is shown below.



By choosing a value greater than e the x -intercept of the tangent line to the graph is negative, which means that option E is false.

Question 22*Answer is E.***Worked solution**

A graph of the general curve looks like



So for $f(x) < 0$, then $a + b < 0$; i.e. $a < -b$.

**Tip**

- *If it helps, choose arbitrary values for a and b to produce a sketch.*


SECTION 2

Question 1a.

Worked solution

$$\text{Mean} = \int_1^{\infty} \left(x \times \frac{1}{4} e^{\frac{-(x-1)}{4}} \right) dx$$

Using CAS, we get



This gives the mean of 5 years.


Mark allocation: 2 marks

- 1 mark for writing: $\text{Mean} = \int_1^{\infty} \left(x \times \frac{1}{4} e^{\frac{-(x-1)}{4}} \right) dx$.
- 1 mark for answer of 5 years.

Question 1b.**Worked solution**

Median = m , such that $\int_1^m \frac{1}{4} e^{\frac{-(x-1)}{4}} dx = 0.5$.

Using CAS to solve for m gives $m = 3.77$ years.


**Mark allocation: 2 marks**

- 1 mark for writing $\int_1^m \frac{1}{4} e^{\frac{-(x-1)}{4}} dx = 0.5$.
- 1 mark for answer of $m = 3.77$ years.

Question 1c.**Worked solution**

$$\begin{aligned}
 \text{var}(X) &= E(X^2) - (E(X))^2 \\
 &= \int_1^{\infty} x^2 f(x) dx - \left(\int_1^{\infty} x f(x) dx \right)^2 \\
 &= \int_1^{\infty} x^2 \frac{1}{4} e^{\frac{-(x-1)}{4}} dx - \left(\int_1^{\infty} x \frac{1}{4} e^{\frac{-(x-1)}{4}} dx \right)^2
 \end{aligned}$$

Using CAS, this gives $\text{var}(X) = 16$ years.

**Mark allocation: 2 marks**


- 1 mark for stating $\text{var}(X) = \int_1^{\infty} x^2 \frac{1}{4} e^{\frac{-(x-1)}{4}} dx - \left(\int_1^{\infty} x \frac{1}{4} e^{\frac{-(x-1)}{4}} dx \right)^2$.
- 1 mark for answer $\text{var}(X) = 16$ years.

Question 1d.

Worked solution

$$\begin{aligned}\Pr(X > 5) &= \int_5^\infty \frac{1}{4} e^{\frac{-(x-1)}{4}} dx \\ &= \frac{1}{e} \quad (\text{Found using CAS.})\end{aligned}$$

Using CAS gives



Mark allocation: 2 marks

- 1 mark for stating $\Pr(X > 5) = \int_5^{\infty} \frac{1}{4} e^{\frac{-(x-1)}{4}} dx$.
- 1 mark for answer $\frac{1}{e}$.



Tip

- *The answer requires an exact answer, so be sure to have your CAS calculator in exact/standard mode. A decimal answer, no matter how accurate, will not be accepted in this instance.*

Question 1e.**Worked solution**

$$Y \sim \text{Bi}\left(n = 4, p = \Pr(X > 5) = \frac{1}{e}\right)$$

$$E(Y) = n \times p = 4 \times \frac{1}{e} = \frac{4}{e}$$

Mark allocation: 2 marks


- 1 mark for identifying the binomial and the parameters.
- 1 mark for answer $E(Y) = \frac{4}{e}$.

**Tip**

- Although not explicitly asked for, this question also requires an exact answer.

Question 1f.**Worked solution**

To be operational, the television requires at least two switches to be working; i.e. $\Pr(Y \geq 2)$.
Using CAS, this gives $\Pr(Y \geq 2) = 0.4687$.

**Mark allocation: 2 marks**


- 1 mark for stating $\Pr(Y \geq 2)$.
- 1 mark for answer 0.4687.

Question 1g.**Worked solution**

$$N \sim (n = 12, p = 0.468662)$$

$$\Pr(N = 7) = 0.1666 \quad (\text{Found using CAS.})$$

Using CAS gives

**Mark allocation: 2 marks**

- 1 mark for recognising the binomial with $n = 12$ and $p = 0.4688$.
- 1 mark for answer 0.1666.

**Tip**

- To answer correct to 4 decimal places requires that your calculations be carried out to at least 5 decimal places.

Question 2a.**Worked solution**

Let $4a\sqrt{x} - x = 0$.

So $4a\sqrt{x} = x$

$$16a^2x = x^2$$

$$16a^2x - x^2 = 0$$

$$x(16a^2 - x) = 0$$

$$x = 0 \text{ or } x = 16a^2$$

Since $c \neq 0$, $c = 16a^2$.

Mark allocation: 2 marks


- 1 mark for setting $4a\sqrt{x} - x = 0$.
- 1 mark for $c = 16a^2$.

Question 2b.**Worked solution**

f_a is strictly decreasing for $f'_a \leq 0$.

Using CAS, we get $f'(x) = 0$ for $x = 4a^2$.

So, it is strictly decreasing for $x \in [4a^2, \infty)$.

**Mark allocation: 2 marks**

- 1 mark for finding $x = 4a^2$.
- 1 mark for $x \in [4a^2, \infty)$.

**Tip**

- For strictly increasing/decreasing conditions, turning points must be included.

Question 2c.**Worked solution**

At $x = c$, $x = 16a^2$ and $y = 0$ (as determined from part a).

Using CAS, we get $f'(x) = \frac{2a - \sqrt{x}}{\sqrt{x}}$.

$$\begin{aligned} \text{At } x = 16a^2, \quad f'(16a^2) &= \frac{-(4|a| - 2a)}{4|a|} \\ &= -\frac{1}{2}, \text{ since } a > 0. \end{aligned}$$

$$\text{So, } m = -\frac{1}{2}.$$

The image shows two screenshots of a CAS interface, likely TI-Nspire, used for solving a problem. Both screenshots show the same interface with a toolbar at the top and a keypad at the bottom.

Left Screenshot: The main display area shows the equation $\text{solve}\left(\frac{2 \cdot a - \sqrt{x}}{\sqrt{x}} = 0, x\right)$ in the top line. Below it, the solution is shown as $\{x = 4 \cdot a^2\}$. The next line shows the expression $\frac{2 \cdot a - \sqrt{x}}{\sqrt{x}} | x = 16a^2$. The final line shows the simplified result $-\frac{(4 \cdot |a| - 2 \cdot a)}{4 \cdot |a|}$.

Right Screenshot: The main display area shows the expression $\frac{2 \cdot a - \sqrt{x}}{\sqrt{x}} | x = 16a^2$ in the top line. Below it, the simplified result is shown as $-\frac{(4 \cdot |a| - 2 \cdot a)}{4 \cdot |a|}$. The next line shows the further simplified result $-(4a - 2a) / (4a)$. The final line shows the final result $-\frac{1}{2}$.

So equation of the tangent line is

$$y - y_1 = m(x - x_1)$$

$$y - 0 = -\frac{1}{2}(x - 16a^2)$$

$$y = -\frac{1}{2}(x - 16a^2)$$

Mark allocation: 3 marks

- 1 mark for finding $m = -\frac{1}{2}$.
- 1 mark for finding y -intercept.
- 1 mark for finding tangent line equation.

**Tip**

- *The question specifically stated ‘show that ...’, therefore your working must show step-by-step how you obtained your answer.*

Question 2d.**Worked solution**

$$\begin{aligned}
 y &= -\frac{1}{2}(x - 16a^2) \\
 &= -\frac{1}{2}x + 8a^2
 \end{aligned}$$

The tangent line drawn to $f_a(x)$ at $x = c$ passes through the point $(0, 8a^2)$.

So in order for the tangent drawn to $g_a(x)$ to pass through the origin, the graph tangent line must be translated down by $8a^2$, so $b = 8a^2$.

Mark allocation: 2 marks

- 1 mark for finding y -intercept $(0, 8a^2)$.
- 1 mark for finding $b = 8a^2$.

Question 2e.i.**Worked solution**

The area under the curve is given by $\int_0^{16a^2} 4a\sqrt{x} - x \, dx$.

Evaluating this using CAS gives the area is equal to $\frac{128a^4}{3}$ square units, since $a > 0$.

The image shows two screenshots of a CAS interface, likely TI-Nspire, demonstrating the evaluation of the integral $\int_0^{16a^2} 4a\sqrt{x} - x \, dx$.

Left Screenshot: The input is $\int_0^{16a^2} 4a\sqrt{x} - x \, dx$. The CAS displays the antiderivative $-128a^4 + \frac{512a^3 \cdot |a|}{3}$ and then evaluates it to $\frac{128a^4}{3}$.

Right Screenshot: The input is $\int_0^{16a^2} 4a\sqrt{x} - x \, dx$. The CAS displays the antiderivative $-128a^4 + \frac{512a^3 \cdot |a|}{3}$ and then evaluates it to $\frac{128a^4}{3}$.

Both screenshots show the same CAS interface with a toolbar at the top and a keypad at the bottom.

Mark allocation: 2 marks

- 1 mark for setting up the integral.
- 1 mark for the answer $\frac{128a^4}{3}$ square units.

**Tip**

- Make use of the dilation factor and understand what effect this has on the graph and the resulting area.

Question 2e.ii.**Worked solution**

$f\left(\frac{x}{2}\right)$ represents a dilation of factor of 2 parallel to the x -axis. This means the area under the curve is doubled, so the area equals $\frac{256a^4}{3}$ square units.


Mark allocation: 2 marks

- 1 mark for identifying the transformation correctly.
- 1 mark for the answer $\frac{256a^4}{3}$ square units.

Question 2f.**Worked solution**

Find d such that $\frac{1}{d} \int_0^d h_a(x) dx = 0$; i.e. $\int_0^d h_a(x) dx = 0$, since $d \neq 0$.


Using CAS to solve, gives



So $d = \frac{512a^2}{9}$.

Mark allocation: 2 marks

- 1 mark for setting up $\frac{1}{d} \int_0^d h_a(x) dx = 0$.
- 1 mark for the answer $d = \frac{512a^2}{9}$.

Question 3a.**Worked solution****Mark allocation: 2 marks**

- 1 mark for correct shapes and labelling x-intercept correctly.
- 1 mark for labelling turning points correctly.

Question 3b.**Worked solution**

Firework is extinguished when $F = 0$.

When $F = 0$, then

$$\Rightarrow -pt(pt - 24) = 0$$

$$\Rightarrow t = 0 \text{ or } pt = 24$$

$$\text{So } t = \frac{24}{p}.$$


Mark allocation: 2 marks

- 1 mark for setting $F = 0$.
- 1 mark for the answer $t = \frac{24}{p}$.

Question 3c.i.**Worked solution**

Maximum brightness is achieved when $F'(x) = 0$.

Using CAS gives



So the maximum brightness will occur when $\frac{-(4p^2t + 24pt - 48p)}{(t+2)^3} = 0$.

Mark allocation: 2 marks

- 1 mark for finding the derivative.
- 1 mark for setting it equal to zero.

Question 3c.ii.

The image shows two screenshots of a TI-84 Plus calculator interface, likely a TI-84 Plus CE, used for solving a problem. Both screens show the 'Edit Action Interactive' window.

Left Screenshot: The top line shows the derivative of a function $f(x)$ with respect to x , $\frac{d}{dx}(f(x))$. Below this, the expression $-(4 \cdot p^2 \cdot x + 24 \cdot p \cdot x - 48 \cdot p)$ is shown over $(x+2)^3$. The 'solve' button is pressed, resulting in the solution $x = \frac{48 \cdot p}{4 \cdot p^2 + 24 \cdot p}$.

Right Screenshot: The top line shows the expression $-(4 \cdot p^2 \cdot x + 24 \cdot p \cdot x - 48 \cdot p)$ being solved for x , resulting in $x = \frac{48 \cdot p}{4 \cdot p^2 + 24 \cdot p}$. The 'simplify' button is pressed, resulting in the simplified expression $\frac{12}{p+6}$.


So the maximum occurs when $t = \frac{12}{p+6}$.

Mark allocation: 2 marks

- 1 mark for finding t in terms of p .
- 1 mark for simplifying to get $t = \frac{12}{p+6}$.

Question 3d.**Worked solution**

Maximum brightness is at $x = \frac{12}{p+6}$ and $F\left(\frac{12}{p+6}\right) = \frac{36p}{p+12}$, which was found using CAS.


**Mark allocation: 2 marks**

- 1 mark for finding $F\left(\frac{12}{p+6}\right)$.
- 1 mark for the answer $\frac{36p}{p+12}$.

Question 3e.**Worked solution**

For $F \leq 11$, equation becomes $\frac{36p}{p+12} \leq 11$.

Using CAS to find this value of p gives



So the value of K is $\frac{132}{25} = 5.28$.


Mark allocation: 2 marks

- 1 mark for setting $\frac{36p}{p+12} \leq 11$.
- 1 mark for the answer $\frac{132}{25}$ or 5.28.

Question 3f.**Worked solution**

Maximum brightness occurs at $t = \frac{12}{p+6}$.

We want $t \leq 1.5$, so equation becomes $\frac{12}{p+6} \leq 1.5$.



So the values for p that satisfy the conditions are $2 \leq p \leq 5.28$.

Mark allocation: 2 marks

- 1 mark for finding $p \geq 2$.
- 1 mark for $2 \leq p \leq 5.28$.

Question 4a.**Worked solution**

The x -intercepts of the graph are at $(0, 0)$ and $(2, 0)$. These are found when $y = 0$.

Setting $y = 0$ gives

$$\frac{\pi x}{4} \cos(nx) = 0$$

$$\Rightarrow \frac{\pi x}{4} = 0 \text{ or } \cos(nx) = 0$$

$$\Rightarrow x = 0 \text{ or } nx = \frac{\pi}{2}$$

$$\text{So when } x = 2 \Rightarrow 2n = \frac{\pi}{2}.$$

$$\therefore n = \frac{\pi}{4}$$

Mark allocation: 2 marks


- 1 mark for setting $\frac{\pi x}{4} \cos(nx) = 0$.
- 1 mark for correctly simplifying to give $n = \frac{\pi}{4}$.

Question 4b.**Worked solution**

Using CAS, the second x -intercept occurs at $(6, 0)$; so $b = 6$.

Mark allocation: 1 mark


- 1 mark for the correct answer.

Question 4c.i.**Worked solution****Mark allocation: 3 marks**

- 1 mark for shape passing through (0, 4) and (2, 4).
- 1 mark for labelling endpoints correctly.
- 1 mark for labelling turning point correctly.

Question 4c.ii.**Worked solution**

Using CAS, at $x=1$, $y = \frac{-\sqrt{2}\pi}{8} + 4$.

**Mark allocation: 1 mark**


- 1 mark for the correct answer.

**Tip**

- Note that an exact value is required here.

Question 4c.iii.**Worked solution**

Use CAS to find when $f(x) = 6$.



So $f(x) \geq 6$ for $3.1815 \leq x \leq 4.3$, so for 1.119 km.

Mark allocation: 2 marks

- 1 mark for finding the points and intersection.
- 1 mark for the correct answer 1.119.

**Tips**

- Be careful of the endpoint! The graph stops at $x = 4.3$, so the upper endpoint of the interval is 4.3.
- To give an answer that is correct to 3 decimal places requires that your working be done to at least 4 decimal places.


Question 4d.i.**Worked solution**

Use CAS to find p for when $\frac{d}{dx}\left(4 - \frac{\pi x}{4}\cos(px)\right) = 0$ with $x = 1$.

Edit Action Interactive
 0.5 1/2 $\frac{d}{dx}$ $\frac{d}{dx^4}$ Simp $\frac{d}{dx}$ $\frac{d}{dx}$ $\frac{d}{dx}$
 2888, x=-1.095410747, x=
 Define $f(x) = 4 - \frac{\pi x}{4}\cos(px)$
 done
 $\text{solve}\left(\frac{d}{dx}(f(x)) = 0, x\right)$
 $\{p \cdot x \cdot \tan(p \cdot x) - 1 = 0\}$
 Math1 a b c d e f
 Math2 g h i j k l
 Math3 m n o p q r
 Trig s t u v w x
 Var y z CAPS
 abc
 Alg Standard Real Rad

Edit Action Interactive
 0.5 1/2 $\frac{d}{dx}$ $\frac{d}{dx^4}$ Simp $\frac{d}{dx}$ $\frac{d}{dx}$ $\frac{d}{dx}$
 $\text{solve}\left(\frac{d}{dx}(f(x)) = 0, x\right)$
 $\{p \cdot x \cdot \tan(p \cdot x) - 1 = 0\}$
 $p \cdot x \cdot \tan(p \cdot x) - 1 = 0 \mid x = 1$
 $p \cdot \tan(p) - 1 = 0$
 $\text{solve}(p \cdot \tan(p) - 1 = 0, p)$
 0333589, p=0.860333589,
 Math1 Line $\sqrt{\square}$ π $\Rightarrow$
 Math2 Define f g i ∞
 Math3 solve(dSlv ' $\{\square, \square\}$ $\mid$
 Trig < > () { } []
 Var $\leq$ $\geq$ = $\neq$ <
 abc $\leftarrow$ $\rightarrow$ $\frac{d}{dx}$ $\frac{d}{dx^4}$ ans EXE
 Alg Standard Real Rad

Although this gives a number of values for p , $p = 0.8603$ gives the correct outcome. This can be verified by checking the graph $y = 4 - \frac{\pi x}{4} \cos(px)$ with $p = 0.8603$.



Mark allocation: 2 marks


- 1 mark for setting $\frac{d}{dx} \left(4 - \frac{\pi x}{4} \cos(px) \right) = 0$ with $x = 1$.
- 1 mark for the answer $p = 0.8603$.

Question 4d.ii.**Worked solution**


For the maximum to occur at the endpoint, this requires that the maximum turning point occurs at $x \geq 4.3$ and that $f(4.3) > f(0) = 4$.

Use CAS to find p such that $\frac{d}{dx}\left(4 - \frac{\pi x}{4} \cos(px)\right) = 0$ with $x = 4.3$.

This gives a number of values for p . Verifying the correct value for p by checking the graph gives $p = 0.7966$. (Note: The value 0.7967 gives a turning point just before the endpoint.)



Using CAS to solve $f(4.3) > 4$ for p gives $p > 0.3653$.



So for the maximum to occur at $x = 4.3$, then $0.3653 < p < 0.7966$.

Mark allocation: 4 marks

- 1 mark for setting $\frac{d}{dx}\left(4 - \frac{\pi x}{4} \cos(px)\right) = 0$ with $x = 4.3$.
- 1 mark for $p < 0.7966$.
- 1 mark for $f(4.3) > 4$.
- 1 mark for $0.3653 < p < 0.7966$.



Tip

- Don't assume that the normal rules for rounding apply. Always check your answer by sketching a graph and seeing whether it complies with the requirements. In this case, a 'rounded down' answer of 0.7966 was required.

END OF WORKED SOLUTIONS BOOK