

# TEST 9

Functions, graphs, algebra, calculus, probability and statistics  
Technology-free end-of-year examination  
Total marks: 35  
Suggested writing time: 55 minutes

## Specific instructions to students

- Answer all of the questions in the spaces provided.
- Show all workings in questions where more than one mark is available.
- An exact value must be provided in questions where a numerical answer is required, unless otherwise specified.

### QUESTION 1

$\Pr(A)$  and  $\Pr(B)$  are independent events. If  $\Pr(A \cap B) = 0.06$  and  $\Pr(B) = 0.2$ , what is  $\Pr(A)$ ?

$A$  and  $B$  are **independent** if  $\Pr(A \cap B) = \Pr(A) \times \Pr(B)$ .

$$\Rightarrow 0.06 = \Pr(A) \times 0.2$$

$$\Pr(A) = \frac{0.06}{0.2} = 0.3$$

1 mark

### QUESTION 2

- a Two independent events  $A$  and  $B$  have probabilities respectively of 0.2 and 0.9. Find  $\Pr(A \cup B)$ .

Use the addition formula  $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$ .

$$\Pr(A \cup B) = 0.2 + 0.9 - \Pr(A \cap B) \text{ where}$$

$$\Pr(A \cap B) = 0.2 \times 0.9 = 0.18 \text{ (independent)}$$

$$\Pr(A \cup B) = 0.2 + 0.9 - 0.18$$

$$= 1.1 - 0.18$$

$$= 0.92$$

2 marks

- b Two events  $C$  and  $D$  have probabilities respectively of 0.1 and 0.2. If  $\Pr(D | C) = 0.3$ , find  $\Pr(C \cap D)$ .

## Conditional probability

$$\Pr(D | C) = \frac{\Pr(D \cap C)}{\Pr(C)}$$

$$0.3 = \frac{\Pr(D \cap C)}{0.1}$$

$$\Pr(D \cap C) = \Pr(C \cap D) = 0.3 \times 0.1 = 0.03$$

2 marks

(Total: 4 marks)

### QUESTION 3

Differentiate the function  $f(x) = \frac{1}{\sqrt{1-x^2}}$ .

$$f(x) = \frac{1}{\sqrt{1-x^2}} = (1-x^2)^{-\frac{1}{2}}$$

Using the chain rule,

$$f'(x) = -\frac{1}{2}(1-x^2)^{-\frac{3}{2}} \times (-2x)$$

$$= \frac{x}{(1-x^2)^{\frac{3}{2}}}$$

2 marks

### QUESTION 4

- a Let  $f(x) = \sin(3x) \cos(3x)$ . Find  $f'(x)$ .

$$f(x) = \sin(3x) \cos(3x).$$

Using the product rule,

$$f'(x) = \cos(3x) \times 3 \cos(3x) + \sin(3x) \times [-3 \sin(3x)]$$

$$= 3 \cos^2(3x) - 3 \sin^2(3x)$$

2 marks

- b Hence find  $f'\left(\frac{\pi}{12}\right)$ .

$$f'(x) = 3 \cos^2(3x) - 3 \sin^2(3x)$$

$$f'\left(\frac{\pi}{12}\right) = 3 \cos^2\left(\frac{\pi}{4}\right) - 3 \sin^2\left(\frac{\pi}{4}\right)$$

$$= \frac{3}{2} - \frac{3}{2} = 0$$

1 mark

(Total: 3 marks)

### QUESTION 5

Find an anti-derivative with respect to  $x$  of  $\cos(4-2x)$ .

$$\int \cos(4-2x) dx = -\frac{1}{2} \sin(4-2x)$$

1 mark

### QUESTION 6

Consider the following functions.

$$f: D \rightarrow \mathbf{R}, f(x) = \log_e(\log_e(x)) \text{ and}$$

$$g: (b, \infty) \rightarrow \mathbf{R}, g(x) = \frac{1}{4}x$$

- a For the maximal domain of  $f(x)$ , find  $D$ .

For  $f(x) = \log_e(\log_e(x))$  to exist, we need,

$\text{ran}(\text{inner}) \subseteq \text{dom}(\text{outer})$ .

$$\mathbf{R} \not\subset (0, \infty)$$

$f(x)$  will exist if we restrict range inner,  $\mathbf{R}$ , to  $(0, \infty)$ .

So  $\text{dom}(\text{inner}) = (1, \infty)$ .

$$\text{dom } f(x) = \text{dom}(\text{inner}) = (1, \infty)$$

$$D = (1, \infty)$$

2 marks

- b If  $f(g(x))$  is defined over the domain of  $g$ , find the smallest possible value of  $b$ .

Domain of  $g = (b, \infty)$ .

For  $f(g(x))$  to exist, test  $\text{ran}(g) \subseteq \text{dom}(f)$ .

$$\left(\frac{1}{4}b, \infty\right) \subseteq (1, \infty)$$

$$\text{So } \frac{1}{4}b = 1$$

$$b = 4$$

So the least value of  $b = 4$ .

2 marks

(Total: 4 marks)

### QUESTION 7

A probability density function is defined by

$$f(x) = \begin{cases} k \sin(x), & 0 \leq x \leq \pi \\ 0, & \text{otherwise} \end{cases}$$

- a Find the value of  $k$ .

For a PDF,

$$\int_0^\pi k \sin(x) dx = 1$$

$$-k [\cos(x)]_0^\pi = 1$$

$$-k (\cos(\pi) - \cos(0)) = 1$$

$$-k (-1 - 1) = 1$$

$$k = \frac{1}{2}$$

2 marks

- b Hence, find  $\Pr\left(X < \frac{\pi}{3}\right)$ .

$$f(x) = \begin{cases} \frac{1}{2} \sin(x), & 0 \leq x \leq \pi \\ 0, & \text{otherwise} \end{cases}$$

$$\Pr\left(X < \frac{\pi}{3}\right) = \int_0^{\frac{\pi}{3}} \frac{1}{2} \sin(x) dx$$

$$= -\frac{1}{2} [\cos(x)]_0^{\frac{\pi}{3}}$$

$$= -\frac{1}{2} \left[ \cos\left(\frac{\pi}{3}\right) - \cos(0) \right]$$

$$= -\frac{1}{2} \left( \frac{1}{2} - 1 \right)$$

$$= \frac{1}{4}$$

2 marks

(Total: 4 marks)

### QUESTION 8

Solve the equation  $\sin^2(2x) = \cos^2(2x)$  for  $x \in [0, \pi]$ .

$$\sin^2(2x) = \cos^2(2x) \Rightarrow \tan^2(2x) = 1$$

$$\Rightarrow \tan(2x) = \pm 1$$

$$2x = \frac{\pi}{4}, \pi - \frac{\pi}{4}, \pi + \frac{\pi}{4}, 2\pi - \frac{\pi}{4}$$

$$= \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$$

$$x = \frac{\pi}{8}, \frac{3\pi}{8}, \frac{5\pi}{8}, \frac{7\pi}{8}$$

2 marks

**QUESTION 9**

The probability of scoring each goal in netball practice is independent and Claire finds that the probability of her scoring a goal is 0.8.

- a State an expression for the probability that, in 10 shots, Claire scores exactly 2 goals.

$$\begin{aligned}\text{Bi}(n, p) &= \text{Bi}(10, 0.8) \\ \Pr(X = 2) &= {}^{10}C_2 \times (0.2)^2 \times (0.8)^8\end{aligned}$$

2 marks

- b Find the probability that in 4 shots, Claire scores at least 1 goal.

$$\begin{aligned}\text{Bi}(n, p) &= \text{Bi}(4, 0.8) \\ \Pr(X \geq 1) &= 1 - \Pr(X = 0) \\ &= 1 - {}^4C_0 (0.2)^0 (0.8)^4 \\ &= 1 - (0.2)^4 \\ &= 1 - 0.0016 \\ &= 0.9984\end{aligned}$$

2 marks

(Total: 4 marks)

**QUESTION 10**

Solve the following equations for  $x$ .

- a  $2(x + 1)^2 - 7 = 0$

$$\begin{aligned}2(x + 1)^2 - 7 = 0 &\Rightarrow x + 1 = \pm\sqrt{\frac{7}{2}} \\ x &= -1 \pm \sqrt{\frac{7}{2}}\end{aligned}$$

2 marks

- b  $5(x - 1)^3 = 320$

$$\begin{aligned}5(x - 1)^3 = 320 &\Rightarrow (x - 1)^3 = 64 \\ &\Rightarrow x - 1 = 4 \\ x &= 5\end{aligned}$$

2 marks

- c  $\log_{10}(x + 5) = 2$

$$\begin{aligned}\log_{10}(x + 5) = 2 &\Leftrightarrow 10^2 = x + 5 \\ \therefore x &= 10^2 - 5 \\ x &= 95\end{aligned}$$

2 marks

- d  $\sqrt{3} \tan(3x) = -1, x \in [0, \pi]$

$$\begin{aligned}\sqrt{3} \tan(3x) &= -1 \\ \tan(3x) &= -\frac{1}{\sqrt{3}} \\ \text{Reference angle} &= \frac{\pi}{6} \\ \Rightarrow 3x &= \pi - \frac{\pi}{6}, 2\pi - \frac{\pi}{6}, 3\pi - \frac{\pi}{6} \\ &= \frac{5\pi}{6}, \frac{11\pi}{6}, \frac{17\pi}{6} \\ x &= \frac{5\pi}{18}, \frac{11\pi}{18}, \frac{17\pi}{18}\end{aligned}$$

2 marks

- e  $\log_{10}(x + 5) + \log_{10}(x) = 1$

$$\begin{aligned}\log_{10}(x + 5) + \log_{10}(x) &= 1 \\ \Rightarrow \log_{10}(x(x + 5)) &= 1 \\ \Rightarrow \log_{10}(x^2 + 5x) &= 1 \\ \Leftrightarrow x^2 + 5x &= 10^1 \\ x^2 + 5x - 10 &= 0 \\ \Rightarrow x &= \frac{-5 \pm \sqrt{25 + 40}}{2} \\ \text{Testing in the original equation, we reject the} & \\ \text{negative value.} & \\ \text{Answer is } x &= \frac{-5 + \sqrt{65}}{2}\end{aligned}$$

2 marks

(Total: 10 marks)