

Victorian Certificate of Education
2018

SUPERVISOR TO ATTACH PROCESSING LABEL HERE

STUDENT NUMBER

									Letter
--	--	--	--	--	--	--	--	--	--------

MATHEMATICAL METHODS

Written examination 1

Wednesday 7 November 2018

Reading time: 9.00 am to 9.15 am (15 minutes)

Writing time: 9.15 am to 10.15 am (1 hour)

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
9	9	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or correction fluid/tape.

Materials supplied

- Question and answer book of 14 pages
- Formula sheet
- Working space is provided throughout the book.

Instructions

- Write your **student number** in the space provided above on this page.
- Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.
- All written responses must be in English.

At the end of the examination

- You may keep the formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

THIS PAGE IS BLANK

Instructions

Answer **all** questions in the spaces provided.

In all questions where a numerical answer is required, an exact value must be given, unless otherwise specified.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Question 1 (3 marks)

a. If $y = (-3x^3 + x^2 - 64)^3$, find $\frac{dy}{dx}$.

1 mark

b. Let $f(x) = \frac{e^x}{\cos(x)}$.

Evaluate $f'(\pi)$.

2 marks

TURN OVER

Question 2 (3 marks)

The derivative with respect to x of the function $f: (1, \infty) \rightarrow \mathbb{R}$ has the rule $f'(x) = \frac{1}{2} - \frac{1}{(2x-2)}$.

Given that $f(2) = 0$, find $f(x)$ in terms of x .

Question 3 (5 marks)


Let $f: [0, 2\pi] \rightarrow \mathbb{R}$, $f(x) = 2\cos(x) + 1$.

a. Solve the equation $2\cos(x) + 1 = 0$ for $0 \leq x \leq 2\pi$.

2 marks

- b. Sketch the graph of the function f on the axes below. Label the endpoints and local minimum point with their coordinates.

3 marks



TURN OVER

Question 4 (2 marks)

Let X be a normally distributed random variable with a mean of 6 and a variance of 4. Let Z be a random variable with the standard normal distribution.

a. Find $\Pr(X > 6)$.

1 mark

b. Find b such that $\Pr(X > 7) = \Pr(Z < b)$.

1 mark

Question 5 (3 marks)

Let $f: (2, \infty) \rightarrow \mathbb{R}$, where $f(x) = \frac{1}{(x-2)^2}$.
State the rule and domain of f^{-1} .

Question 6 (4 marks)

Two boxes each contain four stones that differ only in colour.

Box 1 contains four black stones.

Box 2 contains two black stones and two white stones.

A box is chosen randomly and one stone is drawn randomly from it.

Each box is equally likely to be chosen, as is each stone.

- a.** What is the probability that the randomly drawn stone is black? 2 marks

- b.** It is not known from which box the stone has been drawn.

Given that the stone that is drawn is black, what is the probability that it was drawn from Box 1?

2 marks

Question 7 (5 marks)

Let P be a point on the straight line $y = 2x - 4$ such that the length of OP , the line segment from the origin O to P , is a minimum.

- a.** Find the coordinates of P .

3 marks

- b.** Find the distance OP . Express your answer in the form $\frac{a\sqrt{b}}{b}$, where a and b are positive integers.

2 marks

CONTINUES OVER PAGE

TURN OVER

Question 8 (7 marks)

Let $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x^2 e^{kx}$, where k is a positive real constant.


- a.** Show that $f'(x) = xe^{kx}(kx + 2)$.

1 mark

- b.** Find the value of k for which the graphs of $y = f(x)$ and $y = f'(x)$ have exactly one point of intersection.

2 marks

Let $g(x) = -\frac{2xe^{kx}}{k}$. The diagram below shows sections of the graphs of f and g for $x \geq 0$.



Let A be the area of the region bounded by the curves $y = f(x)$, $y = g(x)$ and the line $x = 2$.

c. Write down a definite integral that gives the value of A .

1 mark


d. Using your result from **part a.**, or otherwise, find the value of k such that $A = \frac{16}{k}$.

3 marks

TURN OVER

Question 9 (8 marks)

Consider a part of the graph of $y = x \sin(x)$, as shown below.



- a. i. Given that $\int (x \sin(x)) dx = \sin(x) - x \cos(x) + c$, evaluate $\int_{n\pi}^{(n+1)\pi} (x \sin(x)) dx$ when n is a positive **even** integer or 0. Give your answer in simplest form.

2 marks

- ii. Given that $\int (x \sin(x)) dx = \sin(x) - x \cos(x) + c$, evaluate $\int_{n\pi}^{(n+1)\pi} (x \sin(x)) dx$ when n is a positive **odd** integer. Give your answer in simplest form.

1 mark

- b.** Find the equation of the tangent to $y = x \sin(x)$ at the point $\left(-\frac{5\pi}{2}, \frac{5\pi}{2}\right)$. 2 marks

- c.** The translation T maps the graph of $y = x \sin(x)$ onto the graph of $y = (3\pi - x) \sin(x)$, where

$$T: \mathbb{R}^2 \rightarrow \mathbb{R}^2, T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} a \\ 0 \end{bmatrix}$$


and a is a real constant.

State the value of a .

1 mark

- d. Let $f: [0, 3\pi] \rightarrow \mathbb{R}$, $f(x) = (3\pi - x) \sin(x)$ and $g: [0, 3\pi] \rightarrow \mathbb{R}$, $g(x) = (x - 3\pi) \sin(x)$.

The line l_1 is the tangent to the graph of f at the point $\left(\frac{\pi}{2}, \frac{5\pi}{2}\right)$ and the line l_2 is the tangent to the graph of g at $\left(\frac{\pi}{2}, -\frac{5\pi}{2}\right)$, as shown in the diagram below.



Find the total area of the shaded regions shown in the diagram above.

2 marks

**Victorian Certificate of Education
2018**

MATHEMATICAL METHODS

Written examination 1

FORMULA SHEET

Instructions

This formula sheet is provided for your reference.
A question and answer book is provided with this formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Mathematical Methods formulas

Mensuration

area of a trapezium	$\frac{1}{2}(a+b)h$	volume of a pyramid	$\frac{1}{3}Ah$
curved surface area of a cylinder	$2\pi rh$	volume of a sphere	$\frac{4}{3}\pi r^3$
volume of a cylinder	$\pi r^2 h$	area of a triangle	$\frac{1}{2}bc \sin(A)$
volume of a cone	$\frac{1}{3}\pi r^2 h$		

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$		
$\frac{d}{dx}((ax+b)^n) = an(ax+b)^{n-1}$	$\int (ax+b)^n dx = \frac{1}{a(n+1)}(ax+b)^{n+1} + c, n \neq -1$		
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$		
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e(x) + c, x > 0$		
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$		
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$		
$\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$			
product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$	quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$		

Probability

$\Pr(A) = 1 - \Pr(A')$		$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$	
$\Pr(A B) = \frac{\Pr(A \cap B)}{\Pr(B)}$			
mean	$\mu = E(X)$	variance	$\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

Probability distribution		Mean	Variance
discrete	$\Pr(X = x) = p(x)$	$\mu = \sum x p(x)$	$\sigma^2 = \sum (x - \mu)^2 p(x)$
continuous	$\Pr(a < X < b) = \int_a^b f(x) dx$	$\mu = \int_{-\infty}^{\infty} x f(x) dx$	$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$

Sample proportions

$\hat{p} = \frac{X}{n}$		mean	$E(\hat{p}) = p$
standard deviation	$\text{sd}(\hat{p}) = \sqrt{\frac{p(1-p)}{n}}$	approximate confidence interval	$\left(\hat{p} - z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \right)$