

2019 VCE Mathematical Methods Trial Examination 1



Kilbaha Multimedia Publishing
PO Box 2227
Kew Vic 3101
Australia

Tel: (03) 9018 5376
Fax: (03) 9817 4334
kilbaha@gmail.com
<https://kilbaha.com.au>

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Multimedia Publishing.
- The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
- For authorised copying within Australia please check that your institution has a licence from <https://www.copyright.com.au>. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.

For details of the CAL licence for educational institutions contact

CAL, Level 11, 66 Goulburn Street, Sydney, NSW, 2000

Tel: +612 9394 7600 or 1800 066 844

Fax: +612 9394 7601

Email: memberservices@copyright.com.au

- All of these pages must be counted in Copyright Agency Limited (CAL) surveys
- This file must not be uploaded to the Internet.

These questions have no official status.

While every care has been taken, no guarantee is given that these questions are free from error. Please contact us if you believe you have found an error.

CAUTION NEEDED!

All Web Links when created linked to appropriate Web Sites. Teachers and parents must always check links before using them with students to ensure that students are protected from unsuitable Web Content. Kilbaha Multimedia Publishing is not responsible for links that have been changed in this document or links that have been redirected

Victorian Certificate of Education 2019

STUDENT NUMBER

Figures
Words

Letter

--

MATHEMATICAL METHODS

Trial Written Examination 1

Reading time: 15 minutes

Total writing time: 1 hour

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
10	10	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software) notes of any kind, blank sheets of paper, and/or correction fluid/tape.

Materials supplied

- Question and answer book of 20 pages.
- Detachable sheet of miscellaneous formulas at the end of this booklet.
- Working space is provided throughout the booklet.

Instructions

- Detach the formula sheet from the end of this book during reading time.
- Write your **student number** in the space provided above on this page.
- Unless otherwise indicated, the diagrams in this booklet are **not** drawn to scale.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Instructions

Answer **all** questions in the spaces provided.

In all questions where a numerical answer is required an exact value must be given unless otherwise specified.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Question 1 (4 marks)

- a. Let $y = \frac{\sin(2x)}{x^2}$ if $\frac{dy}{dx} = \frac{2f(x)}{x^3}$, find the rule for the function $f(x)$.

2 marks

- b. Let $g(x) = \tan\left(\frac{2}{x}\right)$. Evaluate $g'\left(\frac{6}{\pi}\right)$.

2 marks


Question 3 (3 marks)

- a. Find the general solution of $2 - 4\cos\left(\frac{\pi x}{3}\right) = 0$

1 mark

- b. Sketch the graph of the function $f : [0, 6] \rightarrow \mathbb{R}$ $f(x) = 2 - 4\cos\left(\frac{\pi x}{3}\right)$,
stating the coordinates of all axial intercepts, endpoints and turning points.

2 marks



The tangent to the graph of the curve $y = \sqrt{5 + cx}$ at the point $x = 2$, makes an angle of 135° with the positive direction of the x -axis. Determine the value of c .

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

a. Find $\frac{d}{dx}[\log_e(x^2 + 4)]$.

1 mark

- 3 marks

Question 6 (6 marks)

Sometimes in packets of biscuits, some of the biscuits are broken.

One packet of these biscuits contains 12 biscuits and suppose that $\frac{1}{6}$ of the biscuits are broken.

A random sample of the 12 biscuits in a packet is chosen from the population.

Let $\hat{P}$ represent the sample proportion of broken biscuits in these packets.

- a. Find $\Pr\left(\hat{P} = \frac{1}{12}\right)$, giving your answer in the form $c\left(\frac{a}{b}\right)^n$ where a , b , c and n are positive integers.

2 marks

- b. For a sample of 144 packets, find the standard deviation of $\hat{P}$.

1 mark

- c. Jenny accidentally drops a second packet of these biscuits on the floor and several more of the biscuits became broken. The first packet is known to have two broken biscuits. Jenny then randomly selects one of these packets of biscuits, then takes out and eats two biscuits from this packet. If probability that both the biscuits are not broken are is $\frac{23}{66}$, determine the number of broken biscuits in the packet that was dropped onto the floor.

2 marks

- d. The weights of these packets of biscuits are normally distributed with a mean of 250 grams, with a standard deviation of 8 grams. Let Z be a random variable with the standard normal distribution. If the probability that a packet of biscuits weighs between 244 and 256 grams, is equal to $2 \Pr(w \leq Z \leq 0)$, find the value of w .

1 mark

Question 7 (3 marks)

Consider the function $f : R^+ \rightarrow R$, $f(x) = \frac{1}{x}$.

- a.** Find the equation of the tangent to the graph of f at the point where $x = a$, and $a > 0$.
Give your answer in the form $y = mx + c$.


2 marks

- b.** If O is the origin and this tangent crosses the x -axis at the point R and the y -axis at the point S , find the coordinates of the points R and S and find the area of the triangle ORS .

1 mark

Question 8 (3 marks)

An isosceles triangle has two vertices on the graph of the function $f(x) = e^{-x^2}$ one at the point $P(u, f(u))$ the other at $Q(-u, f(u))$ where $u > 0$ and one at the origin as shown in the diagram below.



Find the value of u , for which the area of the triangle is a maximum.

Question 9 (4 marks)

- a.** The transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ with the rule $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} h \\ k \end{bmatrix}$ maps the graph of $y = x^2$ onto the graph of $y = 11 + 2x - x^2$. Find the values of a , b , h and k .

1 mark

Consider the functions with the rules $f(x) = \frac{1}{\sqrt{x+4}}$ and $g(x) = 11 + 2x - x^2$, defined on their maximal domains.

- b.** Show that $f(g(x))$ does not exist.


1 mark

- 2 marks

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 10 (6 marks)

The graph of the function $f : [0, \pi] \rightarrow \mathbb{R}$, $f(x) = e^{-2x} \sin(2x)$ is shown below.



- a. Find the x -coordinates of the stationary points on the graph of f .

2 marks

- b. Differentiate $e^{-2x} (\cos(2x) + \sin(2x))$ with respect to x .

1 mark

[illegible]

MATHEMATICAL METHODS

Written examination 1

FORMULA SHEET

Directions to students

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

Mathematical Methods formulas

Mensuration

area of a trapezium	$\frac{1}{2}(a+b)h$	volume of a pyramid	$\frac{1}{3}Ah$
curved surface area of a cylinder	$2\pi rh$	volume of a sphere	$\frac{4}{3}\pi r^3$
volume of a cylinder	$\pi r^2 h$	area of triangle	$\frac{1}{2}bc \sin(A)$
volume of a cone	$\frac{1}{3}\pi r^2 h$		

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1}x^{n+1} + c, n \neq -1$
$\frac{d}{dx}((ax+b)^n) = na(ax+b)^{n-1}$	$\int (ax+b)^n dx = \frac{1}{a(n+1)}(ax+b)^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a}e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e(x) + c, x > 0$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$	
product rule $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$	quotient rule $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$	

Probability

$\Pr(A) = 1 - \Pr(A')$		$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$	
$\Pr(A B) = \frac{\Pr(A \cap B)}{\Pr(B)}$			
mean	$\mu = E(X)$	variance	$\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$

Probability distribution		Mean	Variance
discrete	$\Pr(X = x) = p(x)$	$\mu = \sum x p(x)$	$\sigma^2 = \sum (x - \mu)^2 p(x)$
continuous	$\Pr(a < X < b) = \int_a^b f(x) dx$	$\mu = \int_{-\infty}^{\infty} x f(x) dx$	$\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$

Sample proportions

$\hat{p} = \frac{X}{n}$		mean	$E(\hat{p}) = p$
standard deviation	$\text{sd}(\hat{p}) = \sqrt{\frac{p(1-p)}{n}}$	approximate confidence interval	$\left(\hat{p} - z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \right)$

END OF FORMULA SHEET