

Victorian Certificate of Education  
2019

SUPERVISOR TO ATTACH PROCESSING LABEL HERE

STUDENT NUMBER

Letter

## MATHEMATICAL METHODS

## Written examination 1

Wednesday 6 November 2019

Reading time: 9.00 am to 9.15 am (15 minutes)

Writing time: 9.15 am to 10.15 am (1 hour)

## QUESTION AND ANSWER BOOK

## Structure of book

| <i>Number of questions</i> | <i>Number of questions to be answered</i> | <i>Number of marks</i> |
|----------------------------|-------------------------------------------|------------------------|
| 9                          | 9                                         | 40                     |

- Students are to write in blue or black pen.
- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or correction fluid/tape.

**Materials supplied**

- Question and answer book of 13 pages
- Formula sheet
- Working space is provided throughout the book.

**Instructions**

- Write your **student number** in the space provided above on this page.
- Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.
- All written responses must be in English.

**At the end of the examination**

- You may keep the formula sheet.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.**

THIS PAGE IS BLANK

DO NOT WRITE IN THIS AREA

### Instructions

Answer **all** questions in the spaces provided.

In all questions where a numerical answer is required, an exact value must be given, unless otherwise specified.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

#### Question 1 (4 marks)

Let  $f: \left(\frac{1}{3}, \infty\right) \rightarrow \mathbb{R}$ ,  $f(x) = \frac{1}{3x-1}$ .

a. i. Find  $f'(x)$ .

1 mark

---

---

---

---

ii. Find an antiderivative of  $f(x)$ .

1 mark

---

---

---

---

b. Let  $g: \mathbb{R} \setminus \{-1\} \rightarrow \mathbb{R}$ ,  $g(x) = \frac{\sin(\pi x)}{x+1}$ .

Evaluate  $g'(1)$ .

2 marks

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

TURN OVER

**Question 2** (4 marks)

a. Let  $f: R \setminus \left\{\frac{1}{3}\right\} \rightarrow R, f(x) = \frac{1}{3x-1}$ .

Find the rule of  $f^{-1}$ .

2 marks

---

---

---

---

---

---

---

b. State the domain of  $f^{-1}$ .

1 mark

---

c. Let  $g$  be the function obtained by applying the transformation  $T$  to the function  $f$ , where

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} c \\ d \end{bmatrix}$$

and  $c, d \in R$ .

Find the values of  $c$  and  $d$  given that  $g = f^{-1}$ .

1 mark

---

---

---

**Question 3** (3 marks)

The only possible outcomes when a coin is tossed are a head or a tail. When an unbiased coin is tossed, the probability of tossing a head is the same as the probability of tossing a tail.

Jo has three coins in her pocket; two are unbiased and one is biased. When the biased coin is tossed, the probability of tossing a head is  $\frac{1}{3}$ .

Jo randomly selects a coin from her pocket and tosses it.

- a. Find the probability that she tosses a head.

2 marks

---

---

---

---

---

---

- b. Find the probability that she selected an unbiased coin, given that she tossed a head.

1 mark

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

TURN OVER

**Question 4** (4 marks)

- a. Solve  $1 - \cos\left(\frac{x}{2}\right) = \cos\left(\frac{x}{2}\right)$  for  $x \in [-2\pi, \pi]$ .

2 marks

---



---



---



---




---



---

- b. The function  $f: [-2\pi, \pi] \rightarrow \mathbb{R}$ ,  $f(x) = \cos\left(\frac{x}{2}\right)$  is shown on the axes below.



Let  $g: [-2\pi, \pi] \rightarrow \mathbb{R}$ ,  $g(x) = 1 - f(x)$ .

Sketch the graph of  $g$  on the axes above. Label all points of intersection of the graphs of  $f$  and  $g$ , and the endpoints of  $g$ , with their coordinates.

2 marks

**Question 5** (5 marks)

Let  $f: R \setminus \{1\} \rightarrow R$ ,  $f(x) = \frac{2}{(x-1)^2} + 1$ .

- a. i. Evaluate  $f(-1)$ .

1 mark

---




---



---

- ii. Sketch the graph of  $f$  on the axes below, labelling all asymptotes with their equations.

2 marks



- b. Find the area bounded by the graph of  $f$ , the  $x$ -axis, the line  $x = -1$  and the line  $x = 0$ .

2 marks

---



---



---



---



---



---



---

**TURN OVER**

**Question 6** (3 marks)

Fred owns a company that produces thousands of pegs each day. He randomly selects 41 pegs that are produced on one day and finds eight faulty pegs.

- a. What is the proportion of faulty pegs in this sample?

1 mark

---

- b. Pegs are packed each day in boxes. Each box holds 12 pegs. Let  $\hat{P}$  be the random variable that represents the proportion of faulty pegs in a box.

The actual proportion of faulty pegs produced by the company each day is  $\frac{1}{6}$ .

Find  $\Pr\left(\hat{P} < \frac{1}{6}\right)$ . Express your answer in the form  $a(b)^n$ , where  $a$  and  $b$  are positive rational numbers and  $n$  is a positive integer.

2 marks

---

---

---

---


---

---

---

---

The graph of the relation  $y = \sqrt{1 - x^2}$  is shown on the axes below.  $P$  is a point on the graph of this relation,  $A$  is the point  $(-1, 0)$  and  $B$  is the point  $(x, 0)$ .




- b.** Find the maximum area of the triangle  $ABP$ . 3 marks

Find the maximum area of the triangle  $ABP$ .

**Question 8** (4 marks)

The function  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x)$  is a polynomial function of degree 4. Part of the graph of  $f$  is shown below. The graph of  $f$  touches the  $x$ -axis at the origin.



- a. Find the rule of  $f$ .

1 mark

---



---



---



---

Let  $g$  be a function with the same rule as  $f$ .

Let  $h: D \rightarrow \mathbb{R}$ ,  $h(x) = \log_e(g(x)) - \log_e(x^3 + x^2)$ , where  $D$  is the maximal domain of  $h$ .

- b. State  $D$ .

1 mark

---



---



---

c. State the range of  $h$ .

2 marks

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

TURN OVER

**Question 9** (9 marks)

Consider the functions  $f: \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = 3 + 2x - x^2$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$ ,  $g(x) = e^x$ .

- a.** State the rule of  $g(f(x))$ . 1 mark

---

- b.** Find the values of  $x$  for which the derivative of  $g(f(x))$  is negative. 2 marks

---

---

---

---

---

---

---

- c.** State the rule of  $f(g(x))$ . 1 mark

---

- d.** Solve  $f(g(x)) = 0$ . 2 marks

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

- e. Find the coordinates of the stationary point of the graph of  $f(g(x))$ .

2 marks

---

---

---

---

---

---

---

- f. State the number of solutions to  $g(f(x)) + f(g(x)) = 0$ .

1 mark

---

---



**Victorian Certificate of Education  
2019**

**MATHEMATICAL METHODS**

**Written examination 1**

**FORMULA SHEET**

**Instructions**

This formula sheet is provided for your reference.  
A question and answer book is provided with this formula sheet.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.**

## Mathematical Methods formulas

### Mensuration

|                                   |                        |                     |                         |
|-----------------------------------|------------------------|---------------------|-------------------------|
| area of a trapezium               | $\frac{1}{2}(a+b)h$    | volume of a pyramid | $\frac{1}{3}Ah$         |
| curved surface area of a cylinder | $2\pi rh$              | volume of a sphere  | $\frac{4}{3}\pi r^3$    |
| volume of a cylinder              | $\pi r^2 h$            | area of a triangle  | $\frac{1}{2}bc \sin(A)$ |
| volume of a cone                  | $\frac{1}{3}\pi r^2 h$ |                     |                         |

### Calculus

|                                                                |                                                                  |               |                                                                                        |
|----------------------------------------------------------------|------------------------------------------------------------------|---------------|----------------------------------------------------------------------------------------|
| $\frac{d}{dx}(x^n) = nx^{n-1}$                                 | $\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$             |               |                                                                                        |
| $\frac{d}{dx}((ax+b)^n) = an(ax+b)^{n-1}$                      | $\int (ax+b)^n dx = \frac{1}{a(n+1)}(ax+b)^{n+1} + c, n \neq -1$ |               |                                                                                        |
| $\frac{d}{dx}(e^{ax}) = ae^{ax}$                               | $\int e^{ax} dx = \frac{1}{a} e^{ax} + c$                        |               |                                                                                        |
| $\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$                        | $\int \frac{1}{x} dx = \log_e(x) + c, x > 0$                     |               |                                                                                        |
| $\frac{d}{dx}(\sin(ax)) = a \cos(ax)$                          | $\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$                   |               |                                                                                        |
| $\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$                         | $\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$                    |               |                                                                                        |
| $\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$ |                                                                  |               |                                                                                        |
| product rule                                                   | $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$           | quotient rule | $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$ |
| chain rule                                                     | $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$                    |               |                                                                                        |

**Probability**

|                                           |              |                                                   |                                                              |
|-------------------------------------------|--------------|---------------------------------------------------|--------------------------------------------------------------|
| $\Pr(A) = 1 - \Pr(A')$                    |              | $\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$ |                                                              |
| $\Pr(A B) = \frac{\Pr(A \cap B)}{\Pr(B)}$ |              |                                                   |                                                              |
| mean                                      | $\mu = E(X)$ | variance                                          | $\text{var}(X) = \sigma^2 = E((X - \mu)^2) = E(X^2) - \mu^2$ |

| Probability distribution |                                     | Mean                                      | Variance                                                 |
|--------------------------|-------------------------------------|-------------------------------------------|----------------------------------------------------------|
| discrete                 | $\Pr(X = x) = p(x)$                 | $\mu = \sum x p(x)$                       | $\sigma^2 = \sum (x - \mu)^2 p(x)$                       |
| continuous               | $\Pr(a < X < b) = \int_a^b f(x) dx$ | $\mu = \int_{-\infty}^{\infty} x f(x) dx$ | $\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx$ |

**Sample proportions**

|                         |                                                |                                 |                                                                                                                   |
|-------------------------|------------------------------------------------|---------------------------------|-------------------------------------------------------------------------------------------------------------------|
| $\hat{p} = \frac{X}{n}$ |                                                | mean                            | $E(\hat{p}) = p$                                                                                                  |
| standard deviation      | $\text{sd}(\hat{p}) = \sqrt{\frac{p(1-p)}{n}}$ | approximate confidence interval | $\left( \hat{p} - z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}, \hat{p} + z \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \right)$ |