



YEAR 12 *Trial Exam Paper*

2021

MATHEMATICAL METHODS

Written examination 2

Worked solutions

This book presents:

- worked solutions
- mark allocations
- tips.

This trial examination produced by Insight Publications is NOT an official VCAA paper for the 2021 Mathematical Methods written examination 2. The Publishers assume no legal liability for the opinions, ideas or statements contained in this trial examination. This examination paper is licensed to be printed, photocopied or placed on the school intranet and used only within the confines of the purchasing school for examining their students. No trial examination or part thereof may be issued or passed on to any other party, including other schools, practising or non-practising teachers, tutors, parents, websites or publishing agencies, without the written consent of Insight Publications.

Copyright © Insight Publications 2021

SECTION A – Multiple-choice questions

Question	Answer
1	<i>A</i>
2	<i>D</i>
3	<i>B</i>
4	<i>E</i>
5	<i>C</i>
6	<i>D</i>
7	<i>A</i>
8	<i>A</i>
9	<i>B</i>
10	<i>E</i>
11	<i>C</i>
12	<i>D</i>
13	<i>B</i>
14	<i>B</i>
15	<i>B</i>
16	<i>B</i>
17	<i>B</i>
18	<i>C</i>
19	<i>A</i>
20	<i>E</i>


Question 1**Answer: A****Explanatory notes**

Note that


$$\begin{aligned}
 f(x) &= -x^2 + 2x + 2 \\
 &= -[x^2 - 2x] + 2 \\
 &= -[(x-1)^2 - 1] + 2 \\
 &= -(x-1)^2 + 3
 \end{aligned}$$

and so the turning point of the quadratic occurs at the point with coordinates $(1, 3)$.


This can also be found using CAS:



The function is plotted below:



Alternatively, the minimum and maximum can be found using CAS:

The range of f is $(-13, 3]$.**Tips**


- Consider drawing a quick sketch.
- $fMax$ and $fMin$ could be used to find the extrema of the function.
- Note that the minimum occurs at an open endpoint and that the maximum does not occur at an endpoint.

Question 2


Answer: D

Explanatory notes

Use CAS (either the *expand* or *propFrac* commands) to find that $f(x) = \frac{2x+1}{3-x} = -2 - \frac{7}{x-3}$.



A screenshot of a CAS interface window titled "1.1" and "*Doc". The window shows the command `expand((2*x+1)/(3-x))` entered in the input field. The result, $\frac{-7}{x-3} - 2$, is displayed on the right side of the window.



A screenshot of a CAS interface window titled "1.1" and "*Doc". The window shows the command `propFrac((2*x+1)/(3-x))` entered in the input field. The result, $\frac{-7}{x-3} - 2$, is displayed on the right side of the window.

The asymptotes are $x = 3$ (the vertical asymptote) and $y = -2$ (the horizontal asymptote).



Tip


- Use CAS to simplify algebraic fractions. Long division can be used but is time-consuming and may lead to errors.

Question 3**Answer: B****Explanatory notes**

Note that

$$\begin{aligned}
 2 \sin\left(3x - \frac{\pi}{3}\right) - \sqrt{3} &= 0 \\
 \sin\left(3x - \frac{\pi}{3}\right) &= \frac{\sqrt{3}}{2} \\
 3x - \frac{\pi}{3} &= \frac{\pi}{3} + 2k\pi, \frac{2\pi}{3} + 2k\pi \\
 3x &= \frac{2\pi}{3} + 2k\pi, \pi + 2k\pi \\
 &= \frac{2\pi + 6k\pi}{3}, \frac{\pi + 2k\pi}{3} \\
 x &= \frac{2\pi(1+3k)}{9}, \frac{\pi(2k+1)}{3}
 \end{aligned}$$


This can also be found using CAS:



Note that the form of one solution is different to how it's given by the CAS. The CAS gives $\frac{(2 \times n1 - 1) \times \pi}{3}$ which matches the solution $\frac{\pi(1+2k)}{3}$ in option B. The reason these are equivalent is that they both give odd numbers times $\frac{\pi}{3}$. The CAS uses even numbers, subtracting one each time (... , 2 - 1, 4 - 1, 6 - 1, ...) while the written form adds one each time (... , 0 + 1, 2 + 1, 4 + 1, ...).

Question 4**Answer: E****Explanatory notes**

Use the remainder theorem, $p\left(-\frac{3}{2}\right) = -2$, and CAS to find the remainder:

**Tip**

- Recall that the remainder when $p(x)$ is divided by $ax + b$ is $p\left(-\frac{b}{a}\right)$. This is best calculated using CAS to avoid errors.

Question 5**Answer: C****Explanatory notes**

Let $F(x)$ be an antiderivative of $f(x)$. The gradient of $F(x)$ is

- positive if $x < 0$
- zero when $x = 0$
- positive if $x > 0$
- tends to zero as x becomes larger.

Graph C is the only graph which meets all four conditions.

Question 6**Answer: D****Explanatory notes**


The average value of $f(x)$ on the interval $[1, 3]$ is $\frac{1}{3-1} \int_1^3 f(x) dx = \frac{1}{2} \times 6 = 3$.

Suppose that $f(x) = 3$ is a constant function. Then $g(x)$ is also a constant function.

Now note that point $(1, 3)$ is mapped to point $(-2, 10)$ and point $(3, 3)$ is mapped to point $(-6, 10)$:

$$T\left(\begin{bmatrix} 1 \\ 3 \end{bmatrix}\right) = \begin{bmatrix} -2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 3 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 10 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 3 \\ 3 \end{bmatrix}\right) = \begin{bmatrix} -2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 \\ 3 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -6 \\ 10 \end{bmatrix}$$



Therefore

$$\int_{-6}^{-2} g(x) dx = 4 \times 10 = 40.$$

Alternatively, consider that the new integral with the transformed function has the same shape but is twice as wide, three times as tall, reflected in the y -axis and moved up 1. So its area is six times the original area plus the area of a rectangle that is 1 high and 4 wide:

$$A = (6 \times 6) + (4 \times 1) = 40.$$

Question 7**Answer: A****Explanatory notes**

For a binomial distribution, $E(X) = np$ and $\text{Var}(X) = np(1-p)$.


Since $E(X) = 15$ and $sd(X) = \sqrt{6}$ we have two equations:

$$np = 15$$

$$np(1-p) = 6$$

These can be solved by hand or using CAS to give $n = 25$ and $p = \frac{3}{5}$.

Therefore $\Pr(X \geq 15) = 0.5858$ (correct to four decimal places).

**Question 8****Answer: A****Explanatory notes**

Let $Z \sim N(0,1)$.

$$\text{Thus } \Pr(X > c) = \Pr\left(Z > \frac{c-75}{4}\right) = \Pr\left(Z < \frac{75-c}{4}\right).$$

Therefore

$$\frac{75-c}{4} = -1.5$$

$$75-c = -6$$

$$c = 81$$

Question 9**Answer: B****Explanatory notes**First find the value of a .

$$a + 2a + \frac{2}{5}a + \frac{3}{5-a} + 4a = 1$$

$$a = \frac{1}{16}$$

Then

$$\begin{aligned} E(2X+1) &= 1 \times a + 3 \times 2a + 5 \times \frac{2}{5}a + 7 \times \left(\frac{3}{5} - a\right) + 9 \times 4a \\ &= \frac{263}{40} \end{aligned}$$

A screenshot of a CAS calculator window. The top bar shows '1.1', '*Doc', 'RAD', and a close button. The main display area shows the expression $a + 3 \cdot 2 \cdot a + \frac{5 \cdot 2}{5} \cdot a + 7 \cdot \left(\frac{3}{5} - a\right) + 9 \cdot 4 \cdot a$ with $a = \frac{1}{16}$ substituted. The result is $\frac{263}{40}$.

Question 10**Answer: E****Explanatory notes**Solve $\int_0^5 f(x) dx = 1$ using CAS:

A screenshot of a CAS calculator window. The top bar shows '1.1', '*Doc', 'RAD', and a close button. The main display area shows the function definition $f(x) := \frac{a \cdot x \cdot (5-x)}{50} \cdot e^{\frac{-x}{10}}$ and the equation to solve $\text{solve}\left(\int_0^5 f(x) dx = 1, a\right)$. The result for a is $a = \frac{-e^{\frac{1}{2}}}{10 \cdot \left(3 \cdot e^{\frac{1}{2}} - 5\right)}$. The word 'Done' is in the top right corner.

This is equivalent to $a = \frac{\sqrt{e}}{50 - 30\sqrt{e}}$.

Question 11**Answer: C****Explanatory notes**Let $b = e^x$.Then the equation can be written $b^2 + ab - \frac{a}{2} = 0$.This has two solutions: $b = \frac{-a \pm \sqrt{a(a+2)}}{2}$ when $a < -2$ and $a > 0$.However, we also require that $b > 0$ since the equation $b = e^x$ has no solutions otherwise.

A screenshot of a CAS (Computer Algebra System) window. The window title is '*Doc' and it shows 'RAD' mode. The main display shows the quadratic formula solution for b: $b = \frac{\sqrt{a \cdot (a+2)} - a}{2}$ or $b = \frac{-(\sqrt{a \cdot (a+2)} + a)}{2}$. Below this, two solve commands are shown with their respective conditions for b > 0. The first solve command is $\text{solve}\left(\frac{\sqrt{a \cdot (a+2)} - a}{2} > 0, a\right)$ with the condition $a \leq -2 \text{ or } a > 0$. The second solve command is $\text{solve}\left(\frac{-a - \sqrt{a \cdot (a+2)}}{2} > 0, a\right)$ with the condition $a \leq -2$.

Checking with the CAS, it can be seen that both solutions for b are positive only when $a \leq -2$. Since the solutions are equal when $a = -2$, the equation has two solutions when $a < -2$.

Question 12**Answer: D****Explanatory notes**The coordinates of the intercepts are $A(-b, 0)$ and $B(0, \sqrt{b})$. The gradient of line AB is

therefore $\frac{\sqrt{b}}{b} = \frac{1}{\sqrt{b}}$.

At any point $(a, f(a))$, the gradient of the tangent to f is $\frac{1}{2\sqrt{a+b}}$.Equate the gradient of the tangent to f with the gradient of line AB at any point:

$$\frac{1}{2\sqrt{a+b}} = \frac{1}{\sqrt{b}} \text{ if } x = -\frac{3b}{4}.$$

Question 13**Answer: B****Explanatory notes**

The period of the function is π and so $\frac{2\pi}{b} = \pi \Rightarrow b = 2$.

The amplitude is 2 and so $a = \pm 2$.

Since the y -intercept occurs when $y = -\sqrt{3}$ it could be the case that

$$f(x) = 2 \cos\left(2x - \frac{5\pi}{6}\right) \text{ or } f(x) = -2 \cos\left(2x + \frac{\pi}{6}\right).$$

Only the first option matches any of the answers provided.

Question 14**Answer: B****Explanatory notes**

Use the quotient and the product rules:

$$f'(x) = \frac{(g(x) + xg'(x))h(x) - xg(x)h'(x)}{(h(x))^2}$$

$$f'(3) = \frac{(g(3) + 3g'(3))h(3) - 3g(3)h'(3)}{(h(3))^2}$$

$$= \frac{(2-3) \times 5 - 3 \times 2 \times 2}{5^2}$$

$$= \frac{-5-12}{25}$$

$$= -\frac{17}{25}$$



Question 15**Answer: B****Explanatory notes**

The equation of the perpendicular line is $y = \frac{x}{2}$.

The perpendicular line intersects the graph of $y = f(x)$ when $x = -\frac{1}{2}$ and $x = 2$.


Using CAS, the area of the shaded region is

$$\int_{-\frac{1}{2}}^2 \left(f(x) - \frac{x}{2} \right) dx = \frac{125}{48}$$



Question 16**Answer: B****Explanatory notes**

The turning points of $f(x) = 2x^3 + 3x^2 - 36x + 4$ occur when $x = -3$ and $x = 2$.



Therefore f will have an inverse function if D is a subset of $(-\infty, -3]$, $[-3, 2]$ or $[2, \infty)$.

Of the options given, only $[0, 2]$ is a subset of $[-3, 2]$.

Question 17**Answer: B****Explanatory notes**

The probability that all three marbles are red given that at least two of them are red is

$$\begin{aligned} \Pr(3 \text{ Red} | \text{At least 2 Red}) &= \frac{\Pr(3 \text{ Red})}{\Pr(\text{At least 2 Red})} \\ &= \frac{\Pr(RRR)}{\Pr(RRB, RBR, BRR, RRR)} \end{aligned}$$

We can calculate these probabilities:

$$\Pr(RRR) = \frac{6}{10} \times \frac{5}{9} \times \frac{4}{8} = \frac{1}{6}$$

$$\Pr(RRB) = \frac{6}{10} \times \frac{5}{9} \times \frac{4}{8} = \frac{1}{6}$$

$$\Pr(RBR) = \frac{6}{10} \times \frac{4}{9} \times \frac{5}{8} = \frac{1}{6}$$

$$\Pr(BRR) = \frac{4}{10} \times \frac{6}{9} \times \frac{5}{8} = \frac{1}{6}$$

$$\text{Therefore } \frac{\Pr(RRR)}{\Pr(RRB, RBR, BRR, RRR)} = \frac{\frac{1}{6}}{4 \times \frac{1}{6}} = \frac{1}{4}.$$

Alternatively, consider that once two red marbles are removed there are an even number of red and blue marbles. All four possible outcomes with at least two red marbles must have the same probability. So the answer is one quarter.

Question 18**Answer: C****Explanatory notes**

Since A and B are independent, $\Pr(A \cap B) = \Pr(A) \times \Pr(B) = ab$ and so $\Pr(A \cap B') = a - ab$.

Therefore

$$\begin{aligned}\Pr(A \cup B') &= \Pr(A) + \Pr(B') - \Pr(A \cap B') \\ &= a + 1 - b - a + ab \\ &= 1 - b + ab\end{aligned}$$

Question 19**Answer: A****Explanatory notes**

Find the value of $\hat{p}$ first:

$$\hat{p} = \frac{0.3828 + 0.5283}{2} = 0.45555.$$

Now solve the equation $\hat{p} + 1.96\sqrt{\frac{\hat{p}(1-\hat{p})}{n}} = 0.5283$ for n .

Thus $n = 180$ (to the nearest integer). This is the number of students in the sample.

The number of students who study chemistry from this random sample is $180 \times 0.4555 = 82$ (to the nearest integer).

The screenshot shows a TI-84 Plus calculator screen with the following steps:

- Line 1: $p := \frac{0.3828 + 0.5283}{2}$ resulting in 0.45555 .
- Line 2: $\text{solve}\left(p + 1.96 \cdot \sqrt{\frac{p \cdot (1-p)}{n}} = 0.5283, n\right)$ resulting in $n = 180.028059587$.
- Line 3: $p \cdot n | n = 180.02805958664$ resulting in 82.0117825447 .

**Tip**

- Read the question carefully. The question asks for the number of students studying chemistry, not the number of students in the sample.

Question 20**Answer: E****Explanatory notes**

If $F(t)$ is an antiderivative of $\frac{1}{\sqrt{t^2+1}}$, then

$$f(x) = \int_1^x \frac{1}{\sqrt{t^2+1}} dt = F(x) - F(1).$$

Since $F(1)$ is a constant,

$$f'(x) = F'(x) = \frac{1}{\sqrt{x^2+1}}.$$

Therefore

$$\begin{aligned} f'\left(\frac{1}{\sqrt{3}}\right) &= \frac{1}{\sqrt{\frac{1}{3}+1}} \\ &= \frac{1}{\sqrt{\frac{4}{3}}} \\ &= \frac{\sqrt{3}}{2} \end{aligned}$$

Alternatively, use CAS:

The screenshot shows a CAS window with a title bar containing "1.1", "*Doc", "RAD", and a close button. The main display area shows the expression:

$$\frac{d}{dx} \left(\int_1^x \frac{1}{\sqrt{t^2+1}} dt \right) \Big|_{x=\frac{1}{\sqrt{3}}} = \frac{\sqrt{3}}{2}$$

Below the expression, there is a vertical cursor line.


SECTION B

Question 1a.

Worked solution

Use the factor command in CAS to find that

$$f(x) = -\frac{1}{18}x(x+2)(7x^2+4x-29)$$




Mark allocation: 1 mark

- 1 mark for the correct answer

Question 1b.

Worked solution

Use CAS to solve the equation $f(x) = 0$.



Therefore $x = 0$, $x = -2$ or $x = \frac{-2 \pm 3\sqrt{23}}{7}$.

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 1c.**Worked solution**

Use CAS to find the equation of the tangent line:

The CAS interface shows the following steps:

- factor($f(x)$)**:
$$\frac{-x \cdot (x+2) \cdot (7x^2 + 4x - 29)}{18}$$
- solve($f(x)=0, x$)**:
$$x = \frac{-(3\sqrt{23} + 2)}{7} \text{ or } x = -2 \text{ or } x = 0 \text{ or } x = \frac{3\sqrt{23}}{7}$$
- tangentLine($f(x), x, 1$)**:
$$x + 2$$

Therefore $l_1(x) = x + 2$.

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 1d.**Worked solution**

The coordinates of B are $(-2, 0)$.

The gradient of the tangent to f at the point $B(-2, 0)$ is -1 . The gradient from **part b.** is $+1$.

So $m_1 \times m_2 = 1 \times -1 = -1$.

Therefore the tangent to the graph of f at A is the perpendicular to the graph of f at B .

The CAS interface shows the following steps:

- solve($f(x)=0, x$)**:
$$x = \frac{-(3\sqrt{23} + 2)}{7} \text{ or } x = -2 \text{ or } x = 0 \text{ or } x = \frac{3\sqrt{23}}{7}$$
- tangentLine($f(x), x, 1$)**:
$$x + 2$$
- $\frac{d}{dx}(f(x))|_{x=-2}$** :
$$-1$$

Mark allocation: 1 mark

- 1 mark for finding that the gradient of f at $(-2, 0)$ is -1 , from which the conclusion follows

Question 1e.**Worked solution**

The equation of the perpendicular to the graph of f at the origin is

$$l_2(x) = -\frac{9x}{29}.$$

The tangent and perpendicular meet when $l_1(x) = l_2(x) \Rightarrow x = -\frac{29}{19}$. Therefore, the coordinates of the point of intersection are $C\left(-\frac{29}{19}, \frac{9}{19}\right)$.

A calculator window titled "1.1" and "*Doc" with a "RAD" indicator. It shows the following steps:

- $\text{normalLine}(f(x), x, 0)$ resulting in $-\frac{9 \cdot x}{29}$
- $\text{solve}\left(\frac{-9 \cdot x}{29} = x + 2, x\right)$ resulting in $x = -\frac{29}{19}$
- $\frac{-29}{19} + 2$ resulting in $\frac{9}{19}$

Mark allocation: 2 marks

- 1 mark for finding the equation of the normal: $l_2(x) = -\frac{9x}{29}$
- 1 mark for finding the point of intersection: $C\left(-\frac{29}{19}, \frac{9}{19}\right)$

Question 1f.**Worked solution**

The acute angle between lines l_1 and l_2 is

$$\begin{aligned} \tan^{-1}(1) - \tan^{-1}\left(-\frac{9}{29}\right) &= 45^\circ + 17.24^\circ \\ &= 62.24^\circ \end{aligned}$$

A calculator window titled "1.1" and "*Doc" with a "RAD" indicator. It shows the following steps:

- $\text{solve}\left(\frac{-9 \cdot x}{29} = x + 2, x\right)$ resulting in $x = -\frac{29}{19}$
- $\frac{-29}{19} + 2$ resulting in $\frac{9}{19}$
- $\frac{\left(\tan^{-1}(1) - \tan^{-1}\left(\frac{-9}{29}\right)\right) \cdot 180}{\pi}$ resulting in 62.2414593989


Mark allocation: 2 marks

- 1 mark for using the difference of two inverse tangent functions
- 1 mark for the correct answer

Question 1g.**Worked solution**

The area of the shaded region is


$$\int_{-\frac{29}{19}}^0 (l_1(x) - l_2(x)) dx + \int_0^1 (l_1(x) - f(x)) dx = 2.35$$



A calculator window titled '*Doc' with 'RAD' mode selected. The expression $\int_{-\frac{29}{19}}^0 \left(x+2-\frac{-9 \cdot x}{29}\right) dx + \int_0^1 (x+2-f(x)) dx$ is entered. The result 2.35409356725 is displayed.

Alternatively, the area can be described as the sum of the area of a triangle and an integral:

$$\frac{1}{2} \times 2 \times \frac{29}{19} + \int_0^1 (l_1(x) - f(x)) dx = 2.35$$



A calculator window titled '*Doc' with 'RAD' mode selected. The first expression $\int_{-\frac{29}{19}}^0 \left(x+2-\frac{-9 \cdot x}{29}\right) dx + \int_0^1 (x+2-f(x)) dx$ is entered, resulting in 2.35409356725. The second expression $\frac{29}{19} + \int_0^1 (x+2-f(x)) dx$ is entered, also resulting in 2.35409356725.

Mark allocation: 2 marks

- 1 mark for the appropriate integrals (or a triangle plus an integral)
- 1 mark for the correct answer

Question 2a.**Worked solution**

Using CAS, the coordinates of the turning point are $\left(\sqrt{e}, \frac{1}{2e}\right)$.

CAS interface showing the definition of $f(x) := \frac{\ln(x)}{x^2}$. The derivative is solved for $\frac{d}{dx}(f(x)) = 0$, yielding $x = e^{\frac{1}{2}}$. The function value at this point is $f(x)|_{x=e^{\frac{1}{2}}} = \frac{e^{-1}}{2}$.

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 2b.i.**Worked solution**

Use CAS or find the derivative of $\frac{\log_e(x)}{x}$ by hand:

$$\frac{d}{dx} \left(\frac{\log_e(x)}{x} \right) = \frac{1}{x^2} - \frac{\log_e(x)}{x^2}$$

CAS interface showing the derivative of $f(x) = \frac{\ln(x)}{x}$. The derivative is solved for $\frac{d}{dx}(f(x)) = 0$, yielding $x = e^{\frac{1}{2}}$. The derivative value at this point is $\frac{d}{dx} \left(\frac{\ln(x)}{x} \right) \bigg|_{x=e^{\frac{1}{2}}} = \frac{1}{x^2} - \frac{\ln(x)}{x^2}$.

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 2b.ii.**Worked solution**

From **part b.i.** we have

$$\frac{\log_e(x)}{x^2} = \frac{1}{x^2} - \frac{d}{dx} \left(\frac{\log_e(x)}{x} \right)$$

and so

$$\begin{aligned} \int \frac{\log_e(x)}{x^2} dx &= \int \left(\frac{1}{x^2} - \frac{d}{dx} \left(\frac{\log_e(x)}{x} \right) \right) dx \\ &= -\frac{1}{x} - \frac{\log_e(x)}{x} + c \end{aligned}$$

Mark allocation: 2 marks

- 1 mark for rearranging and attempting to integrate both sides
- 1 mark for the correct answer (the constant of integration, c , may be omitted)

Question 2b.iii.**Worked solution**

Using the result from **part b.ii.** we have

$$\begin{aligned} \int_1^{e^2} f(x) dx &= \left[-\frac{1}{x} - \frac{\log_e(x)}{x} \right]_1^{e^2} \\ &= \left(-\frac{1}{e^2} - \frac{\log_e(e^2)}{e^2} \right) - \left(-\frac{1}{1} - \frac{\log_e(1)}{1} \right) \\ &= \left(-\frac{1}{e^2} - \frac{2}{e^2} \right) - (-1) \\ &= 1 - \frac{3}{e^2} \end{aligned}$$

Mark allocation: 2 marks

- 1 mark for substituting values from the previous result
- 1 mark for the correct answer

Question 2b.iv.**Worked solution**

The average value of f on the interval $[1, e^2]$ is

$$\begin{aligned}\frac{1}{e^2-1} \int_1^{e^2} f(x) dx &= \frac{1}{e^2-1} \left(1 - \frac{3}{e^2} \right) \\ &= \frac{1}{e^2-1} \left(\frac{e^2-3}{e^2} \right) \\ &= \frac{e^2-3}{e^2(e^2-1)}\end{aligned}$$


Mark allocation: 1 mark

- 1 mark for the correct answer

Question 2c.i.**Worked solution**

$$f'(x) = \frac{1}{x^3} - \frac{2 \log_e(x)}{x^3}$$

This can be found using CAS:

**Mark allocation: 1 mark**

- 1 mark for the correct answer

Question 2c.ii.**Worked solution**

The minimum value occurs when $x = e^{\frac{5}{6}}$ and is $-\frac{2}{3}e^{-\frac{5}{2}}$ or $-\frac{2}{3e^{\frac{5}{2}}}$.

This can be found using CAS:

The image shows a CAS interface with the following content:

- Top bar: 1.1, *Doc, RAD, and a close button.
- Input: $fMin\left(\frac{1}{x^3} - \frac{2 \cdot \ln(x)}{x^3}, x\right)$
- Result: $x = e^{\frac{5}{6}}$
- Result: $\frac{1}{x^3} - \frac{2 \cdot \ln(x)}{x^3} \Big|_{x=e^{\frac{5}{6}}} = \frac{-2 \cdot e^{-\frac{5}{2}}}{3}$

Mark allocation: 1 mark

- 1 mark for the correct answer

**Tip**

- *Read the question carefully – it is not asking you to find the minimum of f .*

Question 2d.i.**Worked solution**

From the previous question we know that the gradient is a minimum when $x = e^{\frac{5}{6}}$. Therefore the coordinates of the point on the graph of f where the gradient is a minimum are

$$A\left(e^{\frac{5}{6}}, \frac{5}{6}e^{-\frac{5}{3}}\right).$$

This can be found using CAS:

A screenshot of a CAS interface showing the calculation of the minimum gradient and the corresponding function value. The top line shows the derivative $\frac{1}{x^3} - \frac{2 \cdot \ln(x)}{x^3}$ evaluated at $x = e^{\frac{5}{6}}$, resulting in $\frac{-2 \cdot e^{\frac{-5}{2}}}{3}$. The bottom line shows the function $f(x)$ evaluated at $x = e^{\frac{5}{6}}$, resulting in $\frac{5 \cdot e^{\frac{-5}{3}}}{6}$.

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 2d.ii.**Worked solution**

Use CAS to find the point where the tangent to f at A intersects the x -axis:

A screenshot of a CAS interface showing the calculation of the x-intercept of the tangent line at point A. The top line shows the equation $\text{solve}\left(\text{tangentLine}\left(f(x), x, e^{\frac{5}{6}}\right) = 0, x\right)$. The bottom line shows the result $x = \frac{9 \cdot e^{\frac{5}{6}}}{4}$.

Therefore, $b = \frac{9}{4}e^{\frac{5}{6}}$

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 2e.**Worked solution**

The area of the shaded region is made up of an area under the curve and a triangle under the tangent line:

$$\int_1^{e^{\frac{5}{6}}} f(x) dx + \frac{1}{2} \left(\frac{9}{4} e^{\frac{5}{6}} - e^{\frac{5}{6}} \right) \times \frac{5}{6} e^{-\frac{5}{3}} = 1 - \frac{21}{16} e^{-\frac{5}{6}}$$

The screenshot shows a calculator interface with the following expression displayed:

$$\int_1^{e^{\frac{5}{6}}} f(x) dx + \frac{1}{2} \cdot \left(\frac{9}{4} \cdot e^{\frac{5}{6}} - e^{\frac{5}{6}} \right) \cdot \frac{5}{6} \cdot e^{-\frac{5}{3}}$$

The result shown at the bottom is:

$$1 - \frac{21 \cdot e^{-\frac{5}{6}}}{16}$$

Mark allocation: 3 marks

- 1 mark for determining the integral with correct terminals
- 1 mark for determining the area of the triangle
- 1 mark for the correct answer

Question 3a.**Worked solution**

The amplitude is $\frac{1}{2}$ and so $a = \frac{1}{2}$.

The period is 4 and so $\frac{2\pi}{b} = 4 \Rightarrow b = \frac{\pi}{2}$.

The vertical asymptote is $c = \frac{3+2}{2} = \frac{5}{2}$.

Therefore $f(x) = \frac{1}{2} \cos\left(\frac{\pi}{2}x\right) + \frac{5}{2}$.

Mark allocation: 1 mark

- 1 mark for the correct values of a , b and c

Question 3b.i.**Worked solution**

$$\begin{aligned} d(x) &= f(x) - g(x) \\ &= \frac{1}{2} \cos\left(\frac{\pi}{2}x\right) + \frac{5}{2} - (-\sin(\pi x) + 1) \\ &= \frac{1}{2} \cos\left(\frac{\pi}{2}x\right) + \sin(\pi x) + \frac{3}{2} \end{aligned}$$

Mark allocation: 1 mark

- 1 mark for showing the result

Question 3b.ii.**Worked solution**


The period of $d(x)$ is 4. This can be seen from the graph, or from the lowest common multiple of the periods of the two parts of the graph.

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 3b.iii.**Worked solution**

Use CAS to show that the range of $d(x)$ is $[0.132, 2.868]$:

**Mark allocation: 2 marks**

- 1 mark for each endpoint (up to 2 marks)

Question 3c.**Worked solution**

The average value of $d(x)$ over one period is


$$\begin{aligned} \frac{1}{4} \int_0^4 \left(\frac{3}{2} + \frac{1}{2} \cos\left(\frac{\pi}{2}x\right) + \sin(\pi x) \right) dx &= \frac{1}{4} \left[\frac{3}{2}x + \frac{1}{\pi} \sin\left(\frac{\pi}{2}x\right) - \frac{1}{\pi} \cos(\pi x) \right]_0^4 \\ &= \frac{1}{4} \left(6 - \frac{1}{\pi} \right) - \frac{1}{4} \left(-\frac{1}{\pi} \right) \\ &= \frac{3}{2} \end{aligned}$$

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 3d.**Worked solution**

Using CAS we find that $d(x) = \frac{3}{2}$ when $x = 1, 2.161, 3, 3.839$ and 5 .


**Mark allocation: 2 marks**

- 1 mark for the correct integer values
- 1 mark for the correctly rounded non-integer values

Question 3e.i.**Worked solution**

Use CAS to calculate that the graphs of $d(x)$ and $d(-x)$ meet at points

$(0, 2), \left(1, \frac{3}{2}\right), (2, 1), \left(3, \frac{3}{2}\right)$ and $(4, 2)$.

**Mark allocation: 2 marks**

- 1 mark for the coordinates $(0, 2), (2, 1), (4, 2)$
- 1 mark for the coordinates $\left(1, \frac{3}{2}\right), \left(3, \frac{3}{2}\right)$

Question 3e.ii.**Worked solution**

The area bounded by the graphs of $d(x)$ and $d(-x)$ is

$$\int_0^1 d(x) - d(-x) dx + \int_1^2 d(-x) - d(x) dx + \int_2^3 d(x) - d(-x) dx + \int_3^4 d(-x) - d(x) dx = \frac{16}{\pi}$$

Alternatively, using the symmetry of the graph

$$2 \left(\int_0^1 d(x) - d(-x) dx + \int_1^2 d(-x) - d(x) dx \right) = \frac{16}{\pi}$$

Mark allocation: 2 marks


- 1 mark for appropriate integrals with terminals
- 1 mark for the correct answer

Question 4a.**Worked solution**

Given that $X \sim N(5.2, 1.1^2)$, then

$$\begin{aligned}\Pr(X > 5.7 | X > 5.0) &= \frac{\Pr(X > 5.7 \cap X > 5.0)}{\Pr(X > 5.0)} \\ &= \frac{\Pr(X > 5.7)}{\Pr(X > 5.0)} \\ &= 0.5676\end{aligned}$$

This is found using CAS:


**Mark allocation: 2 marks**

- 1 mark for correctly setting up conditional probability
- 1 mark for the correct answer

Question 4b.**Worked solution**

Use CAS to find that

$$\begin{aligned}E(X) &= \int_1^9 x \times f(x) dx \\ &= 3.8328\end{aligned}$$

**Mark allocation: 2 marks**

- 1 mark for the correct formula for the expected value
- 1 mark for the correct answer

Question 4c.**Worked solution**

One approach is to use the formula for variance:

$$\begin{aligned}\text{Var}(X) &= E(X^2) - (E(X))^2 \\ &= 2.3630\end{aligned}$$

Therefore $\text{sd}(X) = 1.5372$.

This is found using CAS:

$$\begin{aligned}\int_1^9 (x \cdot f(x)) dx &= 3.83283170321 \\ \int_1^9 (x^2 \cdot f(x)) dx - \left(\int_1^9 (x \cdot f(x)) dx \right)^2 &= 2.36302541829 \\ \sqrt{2.36302541829} &= 1.53721352398\end{aligned}$$

Mark allocation: 2 marks

- 1 mark for finding the variance
- 1 mark for the correct answer

Question 4d.**Worked solution**

$$\begin{aligned}\Pr(X > 7 | X > 5) &= \frac{\Pr(X > 7)}{\Pr(X > 5)} \\ &= 0.1334\end{aligned}$$

This is found using CAS:

$$\begin{aligned}\sqrt{2.36302541829} &= 1.53721352398 \\ \frac{\int_7^9 f(x) dx}{\int_5^9 f(x) dx} &= 0.133393393791\end{aligned}$$


Mark allocation: 1 mark

- 1 mark for the correct answer

Question 4e.**Worked solution**

Let $W \sim \text{Bi}(30, 0.1134)$.

Then $\Pr(W \geq 5) = 0.25$, correct to two decimal places.


**Mark allocation: 1 mark**

- 1 mark for the correct answer

Question 4f.**Worked solution**

$$\begin{aligned} \Pr(\hat{P} > 0.1 | \hat{P} < 0.2) &= \Pr(W > 3 | W < 6) \\ &= \frac{\Pr(3 < W < 6)}{\Pr(W < 6)} \\ &= \frac{\Pr(4 \leq W \leq 5)}{\Pr(W \leq 5)} \\ &= 0.37 \end{aligned}$$

correct to two decimal places.

**Mark allocation: 2 marks**

- 1 mark for expressing the conditional probability statement in terms of a random variable (W , for example)
- 1 mark for the correct answer

Question 4g.**Worked solution**

First determine the value of $\hat{p}$:

$$\begin{aligned}\hat{p} &= \frac{0.10641 + 0.29395}{2} \\ &= 0.20018\end{aligned}$$


Using the left-hand value of the confidence interval we have

$$0.10641 = \hat{p} - s \sqrt{\frac{\hat{p}(1-\hat{p})}{30}}$$

giving $s = 1.2836$.

Let $Z \sim N(0,1)$. Thus $\Pr(Z < -s) = 0.0996$.

Then $c = 1 - 2 \times 0.0996 = 0.8008 = 80$ to the nearest integer.



The screenshot shows a TI-84 Plus calculator screen with the following calculations:

- $p := \frac{0.10641 + 0.29395}{2} = 0.20018$
- $\text{solve}\left(0.10641 = p - s \cdot \sqrt{\frac{p \cdot (1-p)}{30}}, s\right)$ resulting in $s = 1.28356560504$
- $\text{normCdf}(-\infty, -s, 0, 1) | s = 1.2835656050353$ resulting in 0.099647060871
- $1 - 2 \cdot 0.0996 = 0.8008$

Alternatively, $c = \Pr(-s < Z < s) = 0.8007$

Therefore, $c = 80$ to the nearest integer.


Mark allocation: 2 marks

- 1 mark for attempting to find s
- 1 mark for the correct answer

Question 5a.**Worked solution**

Since $f(0) = a^2$, the gradient of the line $g(x)$ is $-a^2$ and so $g(x) = -a^2x + a^2$.


The CAS screen below shows that the function $f(x)$ has been defined and the value of $f(0)$ found.

**Mark allocation: 1 mark**

- 1 mark for the correct answer

Question 5b.**Worked solution**

$f(x) = g(x)$ when $x = 0$, $x = 1$ or $x = 2a$. Therefore the coordinates of the point A are $A(2a, -a^2(2a-1)) = A(2a, a^2(1-2a))$.


**Mark allocation: 1 mark**

- 1 mark for the correct answer

Question 5c.i.**Worked solution**

$h(x)$ is the tangent to $f(x)$ at $x = 2a$. The equation is therefore

$$h(x) = a^2(8a - 3) - a(5a - 2)x.$$

**Mark allocation: 1 mark**

- 1 mark for the correct answer

Question 5c.ii.**Worked solution**

The x -intercept occurs when $x = b$. Therefore

$$h(b) = 0$$

$$\begin{aligned} b &= \frac{a^2(8a - 3)}{a(5a - 2)} \\ &= \frac{a(8a - 3)}{5a - 2} \end{aligned}$$

Mark allocation: 1 mark

- 1 mark for showing the required result

Question 5c.iii.**Worked solution**

The maximum value of b is $\frac{9}{25}$ which occurs when $a = \frac{3}{10}$.

TI-84 Plus calculator screen showing the function $b(a) := \frac{a \cdot (8 \cdot a - 3)}{5 \cdot a - 2}$. The $fMax$ function is used with arguments $b(a), a, 0, \frac{2}{5}$. The results are $a = \frac{3}{10}$ and $b(a) = \frac{9}{25}$.

Mark allocation: 2 marks

- 1 mark for calculating the value of a
- 1 mark for calculating the value of b

Question 5c.iv.**Worked solution**

Solving $a = \frac{a(8a-3)}{5a-2}$ gives $a = 0$ or $a = \frac{1}{3}$.

By considering the graph of $b(a)$, it is seen that $0 \leq a \leq b$ if $0 \leq a \leq \frac{1}{3}$.

TI-84 Plus calculator screen showing the $fMax$ function with arguments $b(a), a, 0, \frac{2}{5}$. The $solve(b(a)=a, a)$ function is also shown, resulting in $a = 0$ or $a = \frac{1}{3}$.

Mark allocation: 1 mark

- 1 mark for the correct answer

Question 5d.i.**Worked solution**

The gradient of the line segment through $f(0)$ and $(1, 0)$ is $-a^2$.

$\frac{d}{dx}(f(x)) = -(a-x)(a-3x+2)$, so the gradient of the tangents to f is equal to $-a^2$ when

$$x = \frac{2a+1 \pm \sqrt{4a^2-2a+1}}{3}$$

The screenshot shows the following steps in a CAS interface:

- $f(x) = -(a-x)(a-3x+2)$
- $\frac{d}{dx}(f(x)) = -a^2$
- $\text{solve}(-(a-x)(a-3x+2) = -a^2, x)$
- $x = \frac{2a+1 - \sqrt{4a^2-2a+1}}{3}$ or $x = \frac{2a+1 + \sqrt{4a^2-2a+1}}{3}$

Mark allocation: 2 marks

- 1 mark for equating $\frac{d}{dx}(f(x))$ to $-a^2$
- 1 mark for finding $x = \frac{2a+1 \pm \sqrt{4a^2-2a+1}}{3}$

Question 5d.ii.**Worked solution**

Let $g_m(x)$ be the tangent to f when $x = \frac{1}{3}(1 + 2a - \sqrt{4a^2 - 2a + 1})$.

Let $g_n(x)$ be the tangent to f when $x = \frac{1}{3}(1 + 2a + \sqrt{4a^2 - 2a + 1})$.

Then $\frac{g_n(0) + g_m(0)}{2} = \frac{1}{27}(16a^3 + 15a^2 - 6a + 2)$.

dx

solve $\left(-(a-m) \cdot (a-3 \cdot m+2) = -a^2, m\right)$

$m = \frac{-\left(\sqrt{4 \cdot a^2 - 2 \cdot a + 1} - 2 \cdot a - 1\right)}{3}$ or $m = \frac{\sqrt{4 \cdot a^2 - 2 \cdot a + 1} + 2 \cdot a + 1}{3}$

$gm(x) := \text{tangentLine}(f(x), x, m) | m = \frac{-\left(\sqrt{4 \cdot a^2 - 2 \cdot a + 1} - 2 \cdot a - 1\right)}{3}$ Done

$gn(x) := \text{tangentLine}(f(x), x, m) | m = \frac{\sqrt{4 \cdot a^2 - 2 \cdot a + 1} + 2 \cdot a + 1}{3}$ Done

$\frac{gm(0) + gn(0)}{2}$ $\frac{16 \cdot a^3 + 15 \cdot a^2 - 6 \cdot a + 2}{27}$

Mark allocation: 1 mark

- 1 mark for finding $\frac{g_n(0) + g_m(0)}{2} = \frac{1}{27}(16a^3 + 15a^2 - 6a + 2)$

Question 5d.iii.**Worked solution**

$$d_m = d_n \text{ when } \frac{g_n(0) + g_m(0)}{2} = a^2.$$

That is

$$\frac{1}{27}(16a^3 + 15a^2 - 6a + 2) = a^2 \text{ giving } a = \frac{1}{4}.$$

$$\text{The vertical distance is } d_m = a^2 - g_m(0) = \frac{\sqrt{3}}{36} \text{ when } a = \frac{1}{4}.$$

The calculator screen shows the following steps:

- Equation entered: $\text{solve}\left(\frac{16 \cdot a^3 + 15 \cdot a^2 - 6 \cdot a + 2}{27} = a^2, a\right)$
- Solutions found: $a = \frac{-1}{2} \text{ or } a = \frac{1}{4} \text{ or } a = 1$
- Final calculation: $a^2 - g_m(0) | a = \frac{1}{4} = \frac{\sqrt{3}}{36}$

Mark allocation: 2 marks

- 1 mark for finding the equation $\frac{1}{27}(16a^3 + 15a^2 - 6a + 2) = a^2$
- 1 mark for finding that $a = \frac{1}{4}$ and $d_m = \frac{\sqrt{3}}{36}$

END OF WORKED SOLUTIONS