

The Mathematical Association of Victoria

Trial Examination 2021

MATHEMATICAL METHODS

Written Examination 2

STUDENT NAME _____

Reading time: 15 minutes

Writing time: 2 hours

QUESTION AND ANSWER BOOK

Structure of examination

<i>Section</i>	<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
A	20	20	20
B	4	4	60
			Total 80

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers, a protractor, set-squares, aids for curve sketching, one bound reference, one approved technology (calculator or software) and, if desired, one scientific calculator. Calculator memory DOES NOT need to be cleared. For approved computer-based CAS, full functionality may be used.
- Students are NOT permitted to bring into the examination room: blank sheets of paper and/or white out liquid/tape.

Materials supplied

- Question and answer book of 24 pages.
- Formula sheet.
- Answer sheet for multiple-choice questions.

Instructions

- Write your **name** in the space provided above on this page.
- Write your **name** on the multiple-choice answer sheet.
Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.
- All written responses must be in English.

At the end of the examination

- Place the answer sheet for multiple-choice questions inside the front cover of this book.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

THIS PAGE IS BLANK

SECTION A- Multiple-choice questions**Instructions for Section A**

Answer **all** questions in pencil on the answer sheet provided for multiple – choice questions.

Choose the response that is **correct** for the question.

A correct answer scores 1, an incorrect answer scores 0.

Marks will **not** be deducted for incorrect answers.

No marks will be given if more than one answer is completed for any question.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Question 1

For the graph of $y = -2 \cos\left(\frac{x}{3}\right) - \frac{1}{2}$, the amplitude and period are respectively

- A. -2 and $-\frac{1}{2}$
- B. -2 and $\frac{2\pi}{3}$
- C. -2 and 6π
- D. 2 and 6π
- E. 2 and $\frac{\pi}{6}$

Question 2

The function $g : D \rightarrow R, g(x) = \frac{1}{1+3x}$ has range $[-0.5, -0.2)$.

The domain D is

- A. $[-2, -1)$
- B. $(-2, -1]$
- C. $[-2, 2.5)$
- D. $(-2, 2.5]$
- E. $(2, 1]$

SECTION A - continued
TURN OVER

Question 3

Let $f(x) = x^5 + mx^3 - nx^2 - 1$, where $m, n \in R$. If $x + 2$ is a factor of f and when f is divided by $x - 1$ the remainder is 5, the values of m and n are respectively


- A. $-\frac{13}{12}$ and $-\frac{73}{12}$
- B. $-\frac{59}{12}$ and $-\frac{25}{12}$
- C. $-\frac{5}{4}$ and $-\frac{23}{4}$
- D. $\frac{13}{12}$ and $-\frac{47}{12}$
- E. $-\frac{59}{12}$ and $-\frac{143}{12}$

Question 4

The following set of simultaneous equations will have no solutions for

$$\begin{aligned} nx - 2y &= m \\ n^2x + 6y &= m + 1 \end{aligned}$$

- A. $n \in R \setminus \{-3\}$, $m \in R \setminus \left\{-\frac{1}{4}\right\}$
- B. $n = -3$, $m = -\frac{1}{4}$ only
- C. $n = -3$, $m \in R \setminus \left\{-\frac{1}{4}\right\}$ only
- D. $n = 0$, $m = -\frac{1}{4}$ or $n = -3$, $m = -\frac{1}{4}$
- E. $n = 0$, $m \in R \setminus \left\{-\frac{1}{4}\right\}$ or $n = -3$, $m \in R \setminus \left\{-\frac{1}{4}\right\}$

Question 5

An equation for the inverse function of the above graph, $y = f(x)$ could be

- A. $f^{-1}(x) = \frac{1}{(x-1)^2} - 3$
- B. $f^{-1}(x) = \frac{2}{(x-1)^2} - 3$
- C. $f^{-1}(x) = 1 + \sqrt{\frac{1}{x+3}}$
- D. $f^{-1}(x) = 1 - \sqrt{\frac{2}{x+3}}$
- E. $f^{-1}(x) = 1 + \sqrt{\frac{2}{x+3}}$

Question 6

Let $f: \left[-\frac{\pi}{3}, \pi\right] \rightarrow \mathbb{R}$, $f(x) = -2\sin(3x) + \sqrt{3}$. The maximum rate of change of f with respect to x occurs when

- A. $x = 0, \frac{\pi}{3}, \frac{2\pi}{3}$
- B. $x = \frac{-\pi}{3}, \frac{\pi}{3}, \pi$
- C. $x = \frac{\pi}{3}$ only
- D. $x = \frac{-\pi}{6}, \frac{\pi}{2}$
- E. $x = \frac{\pi}{6}, \frac{5\pi}{6}$

SECTION A - continued
TURN OVER

Question 7

Consider the functions

$$f: \left[0, \frac{\pi}{2}\right) \rightarrow \mathbb{R}, f(x) = \tan(x) \text{ and } g(x) = \sqrt{2x+1} \text{ over its maximal domain.}$$

The rule and domain, respectively, for the function $h(x) = g(f(x))$ are

- A. $h(x) = \sqrt{2 \tan(x) + 1}$ and $x \in \left[0, \frac{\pi}{2}\right)$
- B. $h(x) = \sqrt{2 \tan(x)} + 1$ and $x \in \left[0, \frac{\pi}{2}\right)$
- C. $h(x) = \sqrt{2 \tan(x) + 1}$ and $x \in \left[-\frac{1}{2}, \infty\right)$
- D. $h(x) = \sqrt{2 \tan(x) + 1}$ and $x \in [0, \infty)$
- E. $h(x) = \tan(\sqrt{2x+1})$ and $x \in \left[-\frac{1}{2}, \infty\right)$

Question 8

A sequence of transformations applied to create the image rule $y = -\sqrt{3x+1} - \frac{1}{4}$ from the original function $y = -\sqrt{x}$, in an appropriate order, could be


- A. a reflection in the x -axis, then a translation 1 unit to the right, and then a dilation by a factor of 3 from the x -axis and finally a translation 4 units down.
- B. a translation of 1 unit to the left, then a dilation by a factor of 3 from the y -axis and finally a translation of 4 units down.
- C. a translation of 1 unit left, then a dilation by a factor of $\frac{1}{3}$ from the y -axis and finally a translation of $\frac{1}{4}$ of a unit down.
- D. a reflection in the x -axis, a translation of 1 unit left, then a dilation by a factor of $\frac{1}{3}$ from the y -axis and finally a translation of $\frac{1}{4}$ of a unit down.
- E. a translation of $\frac{1}{3}$ of a unit left, then a dilation by a factor of $\frac{1}{3}$ from the y -axis and finally a translation of $\frac{1}{4}$ of a unit down.

SECTION A - continued

Question 9

When the transformation described by $T: R^2 \rightarrow R^2, T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 3 & 0 \\ 0 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is applied to the function $y = -x^4$ its image rule is

- A. $y = -3(1-2x)^4$
 B. $y = -\frac{(3x+1)^4}{2}$
 C. $y = \frac{(3x+1)^4}{2}$
 D. $y = \frac{2(x-1)^4}{81}$
 E. $y = -\frac{2(x-1)^4}{81}$

Question 10

The area enclosed by the two graphs shown above with equations $y = f(x) = e^{2x}$ and $y = g(x) = 3e^x + 4$ and the line $x = 0$ can be found by evaluating

- A. $\int_{\log_e(4)}^0 (f(x) - g(x)) dx$
 B. $\int_0^{2\log_e(2)} (f(x) - g(x)) dx$
 C. $\int_{-\infty}^{2\log_e(2)} (g(x) - f(x)) dx$
 D. $\int_0^{\log_e(2)} (g(x) - f(x)) dx$
 E. $\int_{-1}^{2\log_e(2)} (f(x) - g(x)) dx$

SECTION A - continued
TURN OVER

Question 11

If $g(x) = \frac{1}{2^x}$ then $g(1-x)$ equals

- A. $g(-x) + \frac{1}{2}$
- B. $\frac{1}{2}g(x)$
- C. $2g(-x)$
- D. $\frac{1}{2}g(-x)$
- E. $2g(x)$

Question 12


The solution to the equation $\frac{1}{2}e^{x+2} \times e^{2x-1} = 1$ is

- A. $x = \frac{1}{3}\log_e(2) - 1$
- B. $x = \log_e(2) - \frac{1}{3}$
- C. $x = \frac{\log_4(2)}{3\log_2(e)} - \frac{1}{3}$
- D. $x = \frac{1}{3\log_2(e)} - 1$
- E. $x = \frac{1}{3\log_2(e)} - \frac{1}{3}$


Question 13

The graph of the inverse f^{-1} where $f(x) = -3\log_e(2x - 2)$ is


A.




B.




C.



D.



E.



**SECTION A - continued
TURN OVER**

Question 14


A 20 cm piece of wire is cut into two pieces. One of the pieces is made into a square with side length x cm. The other is made into a rectangle where the length is double the width.

The minimum total area, in cm^2 , that can be formed is

- A. $\frac{40}{17}$
- B. $\frac{200}{17}$
- C. $\frac{1600}{289}$
- D. 5
- E. 25

Question 15

The area bounded by the graph of $y = g(x) = 5x^2 + 4$, the x -axis and the lines $x = 0$ and $x = 1$ is estimated by taking the average of the areas of the left-endpoint and right-endpoint rectangles with width $\frac{1}{3}$ of a unit. The left-endpoint rectangles are shown in the diagram below.



The approximation, using the average of the areas, written as a percentage, correct to one decimal place, of the exact area is

- A. 69.9
- B. 86.9
- C. 101.6
- D. 116.3
- E. 304.9

Question 16

Consider the function h with rule

$$h(x) = \begin{cases} \log_e(1-x), & -3 \leq x \leq 0 \\ \log_e(x+1), & 0 < x \leq 3 \end{cases}.$$

The average value of the function can be found by evaluating

- A. $\frac{h(3) - h(-3)}{6}$
- B. $\frac{1}{3} \int_{-3}^0 (h(x)) dx$
- C. $\int_{-3}^3 (xh(x)) dx$
- D. $\frac{1}{6} \int_{-3}^3 (xh(x)) dx$
- E. $\frac{1}{3} \int_{-3}^0 (\log_e(1-x)) dx + \frac{1}{3} \int_0^3 (\log_e(x+1)) dx$

Question 17

A probability distribution function for a discrete random variable X is defined as $\Pr(X = x) = \frac{20}{859}x^2 - \frac{20}{859x}$.

A possible set of values for X is

- A. $\{1, 4, 5\}$
- B. $\{-1, 4, 5\}$
- C. $\{-1, 4, -5\}$
- D. $\{1, -4, 5\}$
- E. $\{1, 4, -5\}$

Question 18

A random sample of 300 Mathematical Methods students were selected and it was found that 80% of them did more than one hour of mathematics every day. The 90% confidence interval for the proportion of Mathematical Methods students who do less than one hour of mathematics per day, correct to three decimal places, is

- A. (0.155, 0.245)
- B. (0.162, 0.238)
- C. (0.762, 0.838)
- D. (0.755, 0.845)
- E. (0.741, 0.859)

SECTION A - continued
TURN OVER

Question 19

The birth weight of new born babies is normally distributed.

0.575% of new borns have a weight less than 2 kg and 0.518% greater than 4.8 kg.

The rule for the distribution is closest to

A. $f(x) = \frac{1}{0.55\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-3.39}{0.55}\right)^2}$

B. $f(x) = \frac{1}{3.39\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-0.55}{3.39}\right)^2}$

C. $f(x) = \frac{1}{0.55\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-3.38}{0.55}\right)^2}$

D. $f(x) = \frac{1}{3.38\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-0.55}{3.38}\right)^2}$

E. $f(x) = \frac{1}{0.87\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-3.38}{0.87}\right)^2}$

Question 20

Let p be the probability function for the number of times, x , a prime number is rolled on a 10-sided fair die in n trials. The sides of the die are numbered from one to ten. One is not a prime number. Let f be the probability function for the number of times, m , a number greater than four is rolled in n trials.

$f(m)$ is given by

A. $p\left(1 - \frac{m}{n}\right)$

B. $1 - p(x)$

C. $p(n - m)$

D. $1 - p(m)$

E. $p(m - n)$

END OF SECTION A

THIS PAGE IS BLANK

TURN OVER

SECTION B**Instructions for Section B**

Answer **all** questions in the spaces provided.

In all questions where a numerical answer is required an exact value must be given unless otherwise specified.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Question 1 (13 marks)

- a. If $\frac{2x+1}{2x-3} = a + \frac{b}{2x-3}$ show that $a=1$ and $b=4$. 1 mark

Let $f(x) = \frac{2x+1}{2x-3}$.

- b. i. State the equations of any asymptotes of the graph of f . 1 mark

- ii. State the maximal domain and range of f . 2 marks

Let $f_1 : \left[-\frac{1}{2}, \frac{1}{6}\right] \rightarrow R, f_1(x) = \frac{2x+1}{2x-3}$.

- c. Explain why $f_1(f_1(x))$ exists. 1 mark


SECTION B - Question 1 - continued

Let $g(x) = f_1(f_1(x)) = \frac{1-6x}{2x-11}$.

- d. i. State the coordinates of the axial intercepts of the graph of g .

2 marks

- ii. Sketch the graph of $y = g(x)$, labelling axial intercepts and endpoints with their coordinates. 2 marks



- e. i. Write down a suitable integral statement that would find the area bounded by the graph of g and the lines $y = x$ and $x = \frac{1}{6}$.

2 marks

- ii. Hence find this area, described in **part e.i.**, giving your answer correct to three decimal places.

1 mark

- iii. State the area enclosed between the graph of g^{-1} and the lines $y = x$ and $y = \frac{1}{6}$, correct to three decimal places.

1 mark

SECTION B - continued
TURN OVER

Question 2 (15 marks)

The graph of $d(t) = e^{-kt} \sin(kt)$, can be used to describe drug absorption in the bloodstream, where d is in mg/litre and time t is in minutes, $t \geq 0$. k is a positive real constant.

- a. Find $d'(t)$. 1 mark

- b. i. Hence show that the general solution for the t -ordinates of the stationary points in the graph of $d(t)$ are found by solving $\tan(kt) = 1$. 2 marks

- ii. State the general solution for the equation $d'(t) = 0$. Give your answer in terms of k . 1 mark

- iii. Hence find the t -ordinate, in terms of k , of the maximum stationary point of the graph of

$$d : \left[0, \frac{2}{k} \right] \rightarrow \mathbb{R}, d(t) = e^{-kt} \sin(kt).$$
1 mark

SECTION B - Question 2 - continued

Juniper is in hospital and needs a particular drug for her pain.

Let $d_j : [0, 10] \rightarrow \mathbb{R}, d_j(t) = 10e^{-0.2t} \sin(0.2t)$ where the particular drug in Juniper's bloodstream is measured in mg/litre at time, t minutes.

- c. After how many minutes will Juniper's bloodstream contain the maximum amount of the drug?
State the maximum amount of the drug.

2 marks

- d. Find the average rate of change of the amount of the drug in Juniper's blood stream, in mg/L/min, over the interval $t \in [0, 10]$. Give your answer correct to three decimal places.

2 marks

- e. Show that the graphs of
 $d_j : [0, 10] \rightarrow \mathbb{R}, d_j(t) = 10e^{-0.2t} \sin(0.2t)$

and

$$p : [0, 10] \rightarrow \mathbb{R}, p(t) = 10e^{-0.2t}$$

intersect at only one point and that it is not at the stationary point found in **part c**.

2 marks

SECTION B - Question 2 - continued
TURN OVER

- f.** Find the difference in time between the maximum stationary point on the graph of

$d : \left[0, \frac{2}{k}\right] \rightarrow \mathbb{R}, d(t) = e^{-kt} \sin(kt)$ and the point where the graph of $y = e^{-kt}$ is tangential to d for

(i) $k = 0.2$ and (ii) $k = 0.02$.

2 marks

- g.** Find the difference in time between the maximum stationary point on the graph of

$d : \left[0, \frac{2}{k}\right] \rightarrow \mathbb{R}, d(t) = e^{-kt} \sin(kt)$ and the point where the graph of $y = e^{-kt}$ is tangential to

d . Give your answer in terms of k .

2 marks

SECTION B - continued

Question 3 (17 marks)

Carlin, the manager of Office Supplies, is always looking for ways to improve customer service. He found that the waiting time, T minutes, for customers to be served has a probability density function w defined by

$$w(t) = \begin{cases} \frac{3}{64}t(4-t)^2 & 0 \leq t \leq 4 \\ 0 & \text{elsewhere} \end{cases}.$$

- a. i.** Find the expected value of T . 1 mark

- ii.** Find the median value of T . Give your answer in minutes correct to two decimal places. 1 mark

- iii.** Find the standard deviation of T . 2 marks

Carlin introduces a customer self-service option and finds that the mean waiting time is halved. The waiting time, T_1 minutes, for customers to be served now has a probability density function w_1 defined by

$$w_1(t) = \begin{cases} at(b-t)^2 & 0 \leq t \leq b \\ 0 & \text{elsewhere} \end{cases} \quad \text{where } a \text{ and } b \text{ are real constants.}$$

- b. i.** Find the values of a and b . 2 marks

- ii.** By what percentage was the standard deviation reduced? 1 mark

SECTION B - Question 3 - continued
TURN OVER

Carlin noticed that their 9 kg Ecostore laundry powder was not selling. A customer survey revealed that the most common complaint was that less than 9 kg of laundry powder was found in the container. Carlin rang the factory that supplies the laundry powder and they claimed that at least 80% of their 9 kg Ecostore laundry powder containers weighed more than 9 kg.

Carlin organized quality control inspectors to visit the factory. They decided to take a random sample of 1000 containers and they found that 867 of them weighed at least 9 kg.

- c. i. Find a 95% confidence interval for the proportion of containers that weigh at least 9 kg.
Give your answer correct to three decimal places. 1 mark

- ii. According to the 95% confidence interval, is the factory misleading Office Supplies?
Explain. What should the factory be claiming? 1 mark

The life time of the ink cartridges in Office Supplies is normally distributed with a mean of 3005.5 pages and a standard deviation of 50 pages when using plain text. Carlin would like to claim that 95% of the cartridges will print at least k pages.

- d. i. Find the value of k correct to one decimal place. 1 mark

- ii. A company buys five cartridges. What is the probability that more than 3 of them will print at least 3030 pages? Give your answer correct to four decimal places. 2 marks

SECTION B - Question 3 - continued

Carlin keeps two boxes of chocolates in the staffroom for the staff. The blue box contains 20 white Ferrero chocolates and 15 dark Ferrero chocolates. The green box contains p white Ferrero chocolates and $p^2 + 1$ dark Ferrero chocolates where $p \in \mathbb{Z}^+$. The probability Carlin selects the blue box is $\frac{1}{3}$. Carlin selects a box and then randomly selects a chocolate for his morning tea. It is a white chocolate.

- e. i. What is the probability it was from the green box? Give your answer in terms of p . 2 marks

- ii. What is the maximum value the probability in **part e.i.** can have? 1 mark


Whether a chocolate is classified as white or dark depends on the amount of cocoa it contains. The amount of cocoa in the white Ferrero chocolates is normally distributed with a mean of 12 g and a standard deviation of 1 g. The amount of cocoa in the dark Ferrero chocolates is normally distributed with a mean of 15 g and a standard deviation of 2 g. A chocolate is classified as white if it has less than b g of cocoa and dark otherwise. The probability that a white chocolate is misclassified is the same as a dark chocolate being misclassified.

- f. If the white chocolate Carlin is eating has been misclassified, what is the smallest amount of cocoa it could contain? 2 marks

SECTION B - continued
TURN OVER

Question 4 (15 marks)

The graph of $y = f(x) = (x+1)(x^3 + x^2 - x + 1)$ is shown below.



- a. Label the stationary points with their coordinates on the graph above.

1 mark

Now consider the graphs of $y = f(x) = (x+1)(x^3 + mx^2 - x + 1)$ where m is a real constant.

- b. Find the x coordinate of each of the stationary points, giving answers in terms of m where appropriate.

2 marks


- c. Hence, explain why there will always be three stationary points for $m \in \mathbb{R} \setminus \{1\}$ and only two when $m = 1$.

2 marks

- d. For what values of m will there be two negative x coordinates of the turning points? 1 mark

- e. State the nature of the stationary point at $(0, 1)$ for the different values of m . 2 marks

The graph of $g: \left[-\frac{3}{2}, 0\right] \rightarrow \mathbb{R}, g(x) = (x+1)(x^3 + x^2 - x + 1)$ is shown below.



- f. Sketch the graph of the inverse function, $y = g^{-1}(x)$ on the set of axes above, labelling the endpoints with their coordinates. 1 mark

The graph of g^{-1} is translated left 1 unit and up 1 unit.

- g. Find the area bounded by the graph of g and the translated graph of g^{-1} . 2 marks

SECTION B - Question 4 - continued
TURN OVER

- h.** Find the equation of the tangent line to the graph of g where the gradient is a maximum. 2 marks

- i.** State a sequence of transformations that will transform the tangent line found in **part h.** to the line that passes through the endpoints of g . 2 marks

END OF QUESTION AND ANSWER BOOK