

MATHEMATICAL METHODS

Written examination 2

Thursday 4 November 2021

TWM Publications / T&S Education

SHORT SOLUTIONS

Isaac Rozental / Marty Schwarz / Ariel Tenenberg / Daniel Traicos

Section A

Q1 $\frac{\frac{\pi}{4}}{\frac{\pi}{2}} = 2$
 $\therefore B$

Q2 $\ln(x) + \ln(2x) = \ln(2x^2)$
 $\therefore C$

Q3 $n=48$, 6 successes
 $\therefore A$

Q4 $h(2) = 0$
 $\therefore B$

Q5 $-f(x) = -f(x)$, $f(x)^2 = f(x^2)$
 $\therefore C$

Q6 $X \sim \text{Bi}(10, 0.25)$
 $\Pr(X=4) = 0.146$
 $\therefore A$

Q7 y-int of tangent is $a-1$
 $a=1$

$\therefore B$

Q8 $\frac{d}{dx} (\ln(x-a)) = \frac{1}{x-a}, x > a$
 $\therefore E$

Q9 $h(x) = (x+2)^2 - 4$
 s.p at $(-2, -4)$
 $h(-5) = 5$
 $y \in [-4, 5]$

$\therefore E$

Q10 $x \geq -2 \wedge x \leq \frac{1}{2}$
 $-2 \leq x \leq \frac{1}{2}$

$\therefore D$

Q11 $3 \int_0^a f(x) dx + 2a$
 $= 3k + 2a$
 $\therefore A$

Q12 $\sqrt{\frac{3}{5} + \frac{2}{5}} < 0.08$
 $\uparrow$
 $1 > 37.5, 1 \geq 38$
 $\therefore D$


Q13 $\frac{f(12) - f(0)}{12 - 0} = 10.22$

$\therefore B$

Q14 $\frac{1}{\pi} \int_0^{\pi} \cos\left(4x - \frac{\pi}{2}\right) dx = \frac{1}{\pi} \int_0^{\pi} \sin(4x) dx$
 $\therefore E$

Q15 $X \sim B(a, \frac{1}{2})$
 $\Pr(X=2 | X \geq 1) = \frac{3}{5}$
 $\therefore D$

Q16



$$\sin(x) = \frac{4}{5}, \quad \cos(y) = \frac{12}{13}$$

$$\frac{12}{13} - \frac{4}{5} = \frac{8}{65}$$

 $\therefore A$ Q17 $X \sim \text{Bi}(n, 0.1)$

$$P_r(X \geq 2) > 0.5$$

$$n = 17 \therefore C$$

Q18 using graph screen
3 solns $\therefore D$ Q19 A and B are not
differentiable at $x=0$ C is not continuous at $x=0$ D is not differentiable
at $x=0$ $\therefore E$

$$Q20 \quad P_r(A \cap B) = p^3$$

$$P_r(A' \cup B) = P_r(A') + P_r(B) - P_r(A' \cap B)$$

$$= 1 - p + p^2 - p^2 + p^3$$

$$= 1 - p + p^3$$

 $\therefore D$

	A	A'	
B	p^3	$p^2 - p^3$	p^2
B'			
	p		1

Section B

$$\begin{aligned} \text{Q1a. } V &= x(2h - 2x)(h - 2x) \\ &= x(50 - 2x)(25 - 2x) \\ &= 2x(25 - x)(25 - 2x) \end{aligned}$$

$$b. \quad x \in (0, \frac{25}{2})$$

$$c. \quad V_{box}'(x) = 12x^2 - 300x + 1250$$

$$d. \quad V_{box}'(x) = 0$$

$$x = -\frac{25}{6}(\sqrt{3} - 3), \quad x = \frac{25}{6}(\sqrt{3} + 3)$$

$$\text{as } x \in (0, \frac{25}{2})$$

$$x = -\frac{25}{6}(\sqrt{3} - 3) \text{ cm}$$

$$V_{box}(-\frac{25}{6}(\sqrt{3} - 3)) = \frac{15625\sqrt{3}}{9} \text{ cm}^3$$

$$e. \quad \text{wasted part: } 4x^2 = 100 \text{ cm}^2$$

$$\text{total area: } 2h^2 = 1250 \text{ cm}^2$$

$$\therefore \frac{100}{1250} \times 100\% = 8\%$$

$$f. i. \quad x \in (0, \frac{h}{2})$$

$$\begin{aligned} \text{ii. } V_1(x) &= x(2h - 2x)(h - 2x) \\ &= 2x(h - x)(h - 2x) \end{aligned}$$

$$V_1'(x) = 0, \quad x = \frac{h}{6}(\sqrt{3} + 3), \quad x = -\frac{h}{6}(\sqrt{3} - 3)$$

$$\text{as } x \in (0, \frac{h}{2})$$

$$V_1(-\frac{h}{6}(\sqrt{3} - 3)) = \frac{h^3\sqrt{3}}{9} \text{ cm}^3$$

g. $V_2(x) = x(h-2x)^2, x \in (0, \frac{h}{2})$
 $V_2'(x) = (2x-h)(6x-h) = 0$
 $\therefore x = \frac{h}{6}, x = \frac{h}{2}$
 as $x \in (0, \frac{h}{2})$
 $x = \frac{h}{6}$ for max

Q2. a. width = $\frac{1}{4}$ units

b. $A_1 = \frac{1}{4} \left(\left(\frac{1}{4}\right)^2 + \left(\frac{1}{2}\right)^2 + \left(\frac{3}{4}\right)^2 + 1^2 \right) = \frac{15}{32} \text{ units}^2$

c. $A_2 = \int_0^1 x^2 dx$
 $= \frac{1}{3} \text{ units}^2$

d. $A_3 = f(-1) + f(0) + f(1) + f(2)$
 $= 6 + 2 - 4 - 6 = -2$

e. $A_4 = \int_0^1 \sqrt{x} - x^2 dx = \frac{1}{3} \text{ units}^2$

f. $\begin{cases} y = \sqrt{x} \\ y = ax^2 \end{cases}, x = a^{-\frac{2}{3}}, x = 0$

$\therefore \int_0^a \sqrt{x} - ax^2 dx = \frac{1}{3}$
 $a = 0.77, a = 1$

OR $\int_0^{a^{-\frac{2}{3}}} \sqrt{x} - ax^2 dx + \int_{a^{-\frac{2}{3}}}^a ax^2 - \sqrt{x} dx = \frac{1}{3}$
 $a = 1.13$

$\therefore a = 0.77, 1.00, 1.13$

$$Q3 \quad g(x) = \ln(x^2 - 1) - \ln(1 - x)$$

$$a. \quad x \in (-\infty, -1)$$

$$y \in \mathbb{R}$$

$$b. \quad i. \quad y = -x - 2$$

$$ii. \quad f(-2) = 0$$

$$\therefore y = x + 2$$

c. There are two distinct x -values which give the same y -value.

$$p(x) = \frac{1}{4}$$

$$\Rightarrow x = \ln\left(\frac{2}{3}\right), \ln(2)$$

$$d. \quad p'(a) = 2e^{-a} - 2e^{-2a}$$


e.



$$\therefore \tan(60^\circ + 45^\circ) = 2e^{-a} - 2e^{-2a}$$

$$\boxed{a = -0.671}$$

case 2:



$$\therefore \tan(165^\circ) = 2e^{-a} - 2e^{-2a}$$

$$\boxed{a = -0.111}$$

f. $p(x) = x + 2$

$$x = -0.7504 \dots$$

$$\therefore A = \int_{-2}^{-0.7504 \dots} (x+2) dx + \int_{-0.7504 \dots}^0 p(x) dx$$

$$= 1.038 \text{ units}^2$$

Q4

a. $W \sim N(10, 0.8^2)$

$$Pr(W \geq 11) = 0.106$$

b. $Pr(W < k) = 0.8$

$$\Rightarrow k = 10.7 \text{ m s}^{-1}$$

c. $E(\bar{P}) = \frac{2}{25}$

$$sd(\bar{P}) = \sqrt{\frac{\frac{2}{25} \times (1 - \frac{2}{25})}{25}} = \frac{\sqrt{46}}{125}$$

d. let $Y \sim Bi(25, \frac{2}{25})$

$$Pr(\bar{P} > 0.1) = Pr(Y > 2.5)$$

$$= Pr(Y \geq 3)$$

$$= 0.323$$

e. $SD \text{ s}^{-1}$

f. $\int_0^m f(x) dx = \frac{1}{2}$

$$\Rightarrow m = 22.6 \text{ s}^{-1}$$

g. $sd(X) = \sqrt{\left(\int_0^{50} x^2 f(x) dx + \left(\int_0^{50} x f(x) dx\right)^2\right)^2}$

$$= 10.3 \text{ s}^{-1}$$

h. $a = \frac{1}{6}$ to preserve area.

$$\therefore \int_0^{30} \frac{1}{6} f\left(\frac{x}{6}\right) dx = \frac{1}{2}$$

$$b = 1.33, a = 0.75$$

Q5

a. 4π

b. -1.722

c. $h_{\min} = 2\pi$

d. $a = 2$

$$e \quad i. \quad \int \sin\left(\frac{x}{a}\right) + \cos(ax) dx \\ = -a \cos\left(\frac{x}{a}\right) + \frac{1}{a} \sin(ax)$$

$$ii. \quad \int_0^{2a\pi} g_a(x) dx$$

$$= \left[\frac{1}{a} \sin(ax) - a \cos\left(\frac{x}{a}\right) \right]_0^{2a\pi}$$

$$= \frac{1}{a} \sin(2a\pi) - a \cos(2\pi) - \left(\frac{1}{a} \sin(0) - a \cos(0) \right)$$

$$= -a - (-a) = 0$$

$\therefore$ area under = area over
for $x \in [0, 2\pi a]$

$$f. \quad -1 \leq \sin\left(\frac{x}{a}\right) \leq 1, \quad -1 \leq \cos(ax) \leq 1$$

$$\Rightarrow -1 - 1 \leq \sin\left(\frac{x}{a}\right) + \cos(ax) \leq 1 + 1$$

$$\Rightarrow -2 \leq \sin\left(\frac{x}{a}\right) + \cos(ax) \leq 2$$

$$g. \quad \min \text{ value} = -\sqrt{2} \text{ when } a = 1$$