



Student Name.....SOLUTIONS.....

Teacher (circle one).....DKI JOR VNA

Homegroup .....

## MATHEMATICAL METHODS (CAS) UNIT 1

### EXAMINATION 2

| Section | # of questions | # of questions to be answered | Number of marks |
|---------|----------------|-------------------------------|-----------------|
| 1       | 16             | 16                            | 16              |
| 2       | 9              | 9                             | 62              |
|         |                | Total                         | 78              |

**Monday November 7th 2016**

Reading Time: 11:45 – 12:00pm (15 minutes)

Writing time: 12:00 – 1:30pm (90 minutes)

#### Instructions to students

This exam consists of **17** questions.

All questions should be answered in the spaces provided.

There are **65** marks available in this examination.

A decimal approximation will not be accepted if an exact answer is required.

Where more than one mark is allocated to a question working must be shown.

Students **may not** bring any notes or any calculators into this exam.

Students may bring one bound reference book into the exam.

Students may bring a CAS and a Scientific calculator into the exam.

No paper or electronic dictionaries may be used.

Diagrams in this exam are not to scale except where otherwise stated.

### FORMULAS

$$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(A \cap B)$$

$$\Pr(A \cap B) = \Pr(A) \cdot \Pr(B)$$

$$\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)}$$

$${}^nC_r = \frac{n!}{(n-r)!r!}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Newton's Iterative formula for approximating roots of a polynomial:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

**Section A: Multiple Choice. Choose the best answer and RECORD ON ANSWER SHEET.**

1 If  $A = \begin{bmatrix} 2 & 7 \\ -5 & 4 \end{bmatrix}$  and  $B = \begin{bmatrix} 11 & 3 \\ -2 & 0 \end{bmatrix}$  then  $2A$

+  $3B$  is equal to

A  $\begin{bmatrix} -29 & 5 \\ -4 & 8 \end{bmatrix}$

B  $\begin{bmatrix} 13 & 10 \\ -7 & 4 \end{bmatrix}$

C  $\begin{bmatrix} 28 & 27 \\ -19 & 12 \end{bmatrix}$

D  $\begin{bmatrix} 37 & 23 \\ -4 & 8 \end{bmatrix}$

☒ E  $\begin{bmatrix} 37 & 23 \\ -16 & 8 \end{bmatrix}$

2 The determinant of the matrix  $\begin{bmatrix} 5 & 3 \\ 1 & 2 \end{bmatrix}$  is

A -7

B -1

C 1

☒ D 7

E 13

$10 - 3$



3 The matrix which determines the transformation, a dilation from the  $x$ -axis of factor 2 followed by reflection in the line  $y = x$  is

A  $\begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$

☒ B  $\begin{bmatrix} 0 & 2 \\ 1 & 0 \end{bmatrix}$

C  $\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$

D  $\begin{bmatrix} 2 & 1 \\ 0 & 1 \end{bmatrix}$

E  $\begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}$



4 The matrix which describes the composition of mappings

- dilation of factor 3 from the  $x$ -axis
- reflection in the line  $y = x$
- reflection in the  $x$ -axis

is

A  $\begin{bmatrix} 3 & 0 \\ -1 & 3 \end{bmatrix}$

B  $\begin{bmatrix} 0 & 0 \\ -1 & 3 \end{bmatrix}$

C  $\begin{bmatrix} 3 & 0 \\ -1 & 0 \end{bmatrix}$

D  $\begin{bmatrix} 0 & 0 \\ -3 & 1 \end{bmatrix}$

☒ E  $\begin{bmatrix} 0 & 3 \\ -1 & 0 \end{bmatrix}$

5 The exact value of  $\cos(-330^\circ)$  is:

A  $\frac{1}{2}$

B  $-\sqrt{3}$

C  $-\frac{\sqrt{3}}{2}$

☒ D  $\frac{\sqrt{3}}{2}$

E  $\sqrt{3}$

6 If  $\sin a = 0.2$ , then  $\cos a$  must equal:

A 0.80

B 0.96

☒ C 0.98

D  $\sqrt{0.96}$

E 0.2

$\cos(a) = \sqrt{1 - \sin^2(a)}$

7 The following limits  $\lim_{x \rightarrow -3} (3x^2 - 4)$  and

$\lim_{x \rightarrow -3} (4 - 3x^2)$  evaluate respectively to:

- A 23 and 9
- B ~~23 and 31~~
- C ~~-31 and 31~~
- D ~~23 and 23~~
- E 23 and -23

8 If  $y = -2x^3 + 6x^2 - 4x + 7$ , then  $\frac{dy}{dx}$  is equal to:

- A  $-2x^2 + 6x - 4$
- B  $-3x^2 + 2x - 4$
- C  $-5x^2 + 8x - 4$
- D  $4x^2 + 12x - 4$
- E  $-6x^2 + 12x - 4$

9 If  $y = x^3 - 2x^2 + 3x - 4$ , then the gradient of the *perpendicular* to this curve at  $x = -2$  is:

- A  $3x^2 - 4x + 3$
- B  $-\frac{1}{23}$
- C  $\frac{-1}{3x^2 - 4x + 3}$
- D  $\frac{1}{23}$
- E -1

10 If a curve passes through the point  $(2, -3)$  and at any point has a gradient given by  $3x^2 + 2x - 1$ , then the  $y$ -intercept of the curve must be:

- A -17
- B -13
- C -9
- D -5
- E -1

$$3x^2 - 4x + 3$$

$$3(-2)^2 + 8 + 3$$

$$23$$

$$x^3 + x^2 - x + c$$

$$2^3 + 2^2 - 2 + c = -3$$

$$10 + c = -3$$

$$c = -13$$

11 A colony of organisms multiplies at a rate according to  $N(t) = 4t^2 + 1$ . The average rate of increase between  $t = 3$  and  $t = 4$  is:

- A 28
- B 37
- C 65
- D 18
- E 38

$$t=3 \quad N(3) = 37$$

$$N(4) = 65$$

12 For the missile fired with displacement equation  $s(t) = 800 - 16t - 3t^2$ , the instantaneous rate of change (velocity) at  $t = 3$  is:

- A -16
- B -6
- C 725
- D -34
- E 34

$$s'(t) = -16 - 6t$$

13 The height,  $h$ , of a netball is given by the equation  $h(x) = -0.1x^2 + 0.9x + 1$  where  $x$  is the horizontal displacement of the ball. The horizontal displacement when the ball reaches its maximum height is:

- A 9
- B 4
- C 5
- D 4.7
- E 4.5


$$-0.2x + 0.9 = 0$$

$$-0.2x = -0.9$$

$$x = \frac{9}{2}$$

14 An arc subtends an angle of  $45^\circ$  at the centre of a circle of radius 8 cm. The length of the arc (in cm) is:

- A 360
- B 6.70
- C 0.68
- D 5.34
- E 53.61



$$\frac{\pi}{4} \times 8$$

15 The function with the rule  $f(x) = 4 \tan\left(\frac{\pi x}{2}\right)$  has period


- A 4
- B  $\frac{1}{2}$
- ☒ C 2
- D  $\frac{\pi}{2}$
- E  $\frac{2}{\pi}$

$$\frac{11}{\frac{11}{2}} \quad \frac{10}{1} \times \frac{2}{11}$$

16 A box contains six red blocks and four blue blocks. Three blocks are drawn from the box without replacement.

The probability that all of the blocks are red is

- A  $\frac{8}{125}$
- B  $\frac{3}{500}$
- C  $\frac{24}{29}$
- D  $\frac{27}{125}$
- ☒ E  $\frac{1}{6}$



$$\frac{6}{10} \times \frac{5}{9} \times \frac{4}{8} = \frac{120}{720}$$

**Section B: Short and Extended response questions. Answer in the space provided.**

**1. (4 marks)**

Evaluate the root of the equation  $x^3 = x + 8$  by carrying out two iterations of Newton's Method, starting at  $x_0 = 2$ . Give your answer to 4 decimal places.

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$f(x) = -x^3 + x + 8$$

$$f'(x) = -3x^2 + 1$$

$$x_1 = 2 - \frac{-2^3 + 2 + 8}{-3 \times 2^2 + 1}$$

$$= 2 - \frac{2}{-11}$$

$$= \frac{24}{11}$$

$$x_2 = \frac{24}{11} - \frac{-\left(\frac{24}{11}\right)^3 + 2 + 8}{-3\left(\frac{24}{11}\right)^2 + 1}$$

$$= 2.1664$$

**2. (5 marks)**

The curve with equation  $y = \sin(x)$  is subject to the following transformations;

- a dilation in the  $y$ -direction (from the  $x$  axis) of 3 units
- a reflection in the  $x$  axis

a What is the equation of the transformed graph?

$$y = -3 \sin(x)$$

b i) What is the maximum value of  $y$  on the transformed curve?

$$\underline{3}$$

ii) When does this maximum occur (exact answer)?

$$\underline{\frac{3\pi}{2}}$$

c What is the exact value of  $y$  when  $x = \frac{\pi}{3}$ ? (Show working)

$$y = -3 \sin\left(\frac{\pi}{3}\right)$$

$$= -3 \times \frac{\sqrt{3}}{2}$$

$$= \underline{\underline{-\frac{3\sqrt{3}}{2}}}$$

1 + 1 + 1 + 2 = 5 marks


3. (8 marks)

Harry hits a ball; which was resting on flat ground, up into a nearby tree. The ball becomes stuck at a point 5 metres horizontally from where it was hit. The path of the ball is given by

$$y = -x^2 + 8x, \quad x \geq 0, y \geq 0$$

where  $x$  is the horizontal distance in metres from the point where the ball rested on the ground and  $y$  is the height in metres of the ball above the ground.

- a. Sketch a graph which shows this situation.



2 marks

- b. How high vertically above the ground is the ball when it becomes stuck? (label this on your graph)

~~16 m~~ 15 m

1 mark

- c. What is the maximum height that the ball reaches above the ground before it becomes stuck? (label this on your graph)

at  $x=4$ ,  $y=16$   
16 m

2 marks

- d. Had the ball not become stuck in the tree, how far horizontally from its starting position would it have landed?

8 m

1 mark

- e. During the ball's flight, over what horizontal distance was the ball 15 metres or higher above the ground?

2 m

at  $x=3$ ,  $y=15$   
so from 3 to 5

2 marks


4. (13 marks)

In a laboratory, the temperature in a controlled environment is given by

$$y(t) = a \cos(bt) + c$$

where  $a$ ,  $b$  and  $c$  are constants,  $y$  represents the temperature in degrees Celsius and  $t$  represents the time in hours where  $t=0$  corresponds to noon on Wednesday.

The graph of the function is shown below.



a. Explain why

a.  $a=10$

amplitude

b.  $b = \frac{\pi}{4}$

$P = \frac{2\pi}{b} \Rightarrow P = 8$  so  $8 = \frac{2\pi}{b}$ ,  $b = \frac{2\pi}{8} = \frac{\pi}{4}$

c.  $c=20$

moved -p 20

b. Write down:

1+1+1 = 3 marks

a. the minimum temperature in the controlled environment.

$10^{\circ}\text{C}$

b. the mean temperature in the controlled environment.

$20^{\circ}\text{C}$

1+1 = 2 marks

c. Write down the temperature in the controlled environment at 3pm on Wednesday. Express your answer in degrees Celsius correct to 2 decimal places.

$t=3$ ,  $y(3) = 10 \cos\left(\frac{\pi}{4} \cdot 3\right) + 20$   
 $= 12.93^{\circ}\text{C}$

1 mark

- d. At what time did the temperature in the controlled environment first reach  $25^{\circ}\text{C}$ ?

$$t = 1\frac{1}{3} \text{ hours} \Rightarrow 1:20 \text{ pm Wednesday}$$

2 marks

- e. Hence find for what period of time the temperature is greater than or equal to  $25^{\circ}\text{C}$  between noon and midnight on Wednesday?

$$\text{first } 1\frac{1}{3} \text{ hrs, at } 6\frac{2}{3} \text{ till } 9\frac{1}{3} \\ \Rightarrow 2\frac{2}{3}$$

so for 4 hours

3 marks

In a second controlled environment the temperature is given by  $p(t) = 8 \cos\left(\frac{\pi t}{4}\right) + 20$ ,  $t \geq 0$  where  $p$  represents the temperature in degrees Celsius and  $t$  represents the time in hours where  $t=0$  corresponds to noon on Wednesday.

- f. When is the temperature in the original controlled environment first one degree higher than in the second controlled environment?

Explain how you found your answer.

$$10 \cos\left(\frac{\pi t}{4}\right) + 20 = 8 \cos\left(\frac{\pi t}{4}\right) + 21$$

intersection at

$$1\frac{1}{3} \text{ so } 1:20 \text{ pm}$$

2 marks

5. (3 marks)

If  $\sin \theta = 0.3$ , write down the value of:

|   |                      |   |                      |   |                 |
|---|----------------------|---|----------------------|---|-----------------|
| a | $\sin(\pi - \theta)$ | b | $\sin(\pi + \theta)$ | c | $\sin(-\theta)$ |
|   | 0.3                  |   | -0.3                 |   | -0.3            |

6. (9 marks)

A particle moves in a straight line and starts from a fixed point  $O$  on the line.

Its acceleration  $a \text{ m/s}^2$  is given by  $a = 24t - 72$  for any time  $t$  seconds,  $t \geq 0$ .

a What is its acceleration at  $t = 5$ ?

$$a|_{t=5} = 24 \times 5 - 72 \\ = 48 \text{ m/s}^2$$

1 mark

b If its initial velocity is  $96 \text{ m/s}$ , what is its velocity for any time  $t$  seconds?

$$v(t) = 12t^2 - 72t + C \\ v(0) = 96 = C \\ \Rightarrow v(t) = 12t^2 - 72t + 96$$

2 marks

c When is its velocity zero?

$$0 = 12t^2 - 72t + 96 \\ t = 2 \text{ and } t = 4$$


1 mark

d What is its displacement from  $O$  at any time  $t$  seconds?

$$\text{at } t=0, x(0)=0 \\ \Rightarrow x(t) = 4t^3 - 36t^2 + 96t$$

1 mark

e Find the total distance covered by the particle in the first 6 seconds.



$$\begin{aligned} 2 \text{ seconds} &= 80 \text{ m} \\ \text{next } 2 \text{ seconds} &= 16 \text{ m} \\ \text{next } 2 \text{ seconds} &= 80 \text{ m} \\ \hline \text{total} &= 176 \text{ m} \end{aligned}$$

4 marks

7. (4 marks)

From a group of 12 female students, 2 female staff, 10 male students and 3 male staff, a committee of 6 is to be formed.

Find the number of different committees if:

- a. there are no restrictions.

$${}^{27}C_6 = 296010$$

- b. All committee members must be students

$${}^{22}C_6 = 74613$$

- c. There is an equal number of males and females on the committee

$${}^{14}C_3 \times {}^{13}C_3 = 104104$$

- d. The committee must comprise of 1 male staff member, 1 female staff member, 2 female students and 2 male students.

$${}^3C_1 \times {}^2C_1 \times {}^{12}C_2 \times {}^{10}C_2 = 17820$$

**8. (14 marks)**

In an animated computer game an adventurer can travel from his base to a treasure chest by three different routes. The route is chosen randomly.

The probability of the adventurer travelling on the desert route is  $\frac{1}{2}$ .


The probability of the adventurer travelling on the jungle route is  $\frac{1}{6}$ .

The probability of the adventurer travelling on the mountain route is  $\frac{1}{3}$ .

The probability that the adventurer obtains the treasure by a given route is given in the table below.

| Treasure given<br>desert route | Treasure given<br>jungle route | Treasure given<br>mountain route |
|--------------------------------|--------------------------------|----------------------------------|
| $\frac{1}{5}$                  | $\frac{2}{5}$                  | $\frac{1}{10}$                   |

a Complete the following tree diagram.



3 marks

b Find the probability that the adventurer follows the desert route and obtains the treasure.

$$\frac{1}{2} \times \frac{1}{5} = \frac{1}{10}$$

1 mark

c Find the following probabilities:

i the adventurer obtains the treasure

$$\frac{1}{10} + \frac{1}{2} \times \frac{2}{5} + \frac{1}{3} \times \frac{1}{10} = \frac{1}{5}$$

2 marks

ii the adventurer does not obtain the treasure

$$1 - \frac{1}{5} = \frac{4}{5}$$

2 marks

d Find the probability that, given the adventurer obtains the treasure, he has travelled:

i the desert route

$$\begin{aligned} \Pr(\text{desert} | \text{treasure}) &= \frac{\Pr(\text{desert and treasure})}{\Pr(\text{treasure})} \\ &= \frac{1}{10} \times \frac{5}{1} = \frac{1}{2} \end{aligned}$$

2 marks

ii the jungle route

$$\begin{aligned} \Pr(\text{jungle} + \text{treasure}) &= \frac{1}{6} \times \frac{2}{5} = \frac{1}{15} \\ \Pr(\text{jungle} | \text{treasure}) &= \frac{1}{15} \times \frac{5}{1} = \frac{1}{3} \end{aligned}$$

2 marks

iii the mountain route.

$$\begin{aligned} \Pr(\text{mountain} + \text{treasure}) &= \frac{1}{3} \times \frac{1}{10} = \frac{1}{30} \\ \Pr(\text{mountain} | \text{treasure}) &= \frac{1}{30} \times \frac{5}{1} = \frac{1}{6} \end{aligned}$$

2 marks

9. (2 marks)

Find all possible rational  $x$  intercepts of  $3x^3 + 4x^2 - 7x + 6$

factors of 3 are  $\pm 1, \pm 3$

factors of 6 are  $\pm 1, \pm 2, \pm 3, \pm 6$

Possible roots are  $\pm 1, \pm \frac{1}{2}, \pm \frac{1}{3}, \pm \frac{1}{6}, \pm 3, \pm \frac{3}{2},$

**END OF EXAMINATION**