

Year 2004

VCE

Specialist Mathematics

Trial Examination 1

Suggested Solutions

© Kilbaha Pty Ltd 2004



Kilbaha Multimedia Publishing ABN 47 065 111 373
PO Box 2227
Kew Vic 3101
Australia
Tel: 03 9817 5374
Fax: 03 9817 4334
chemas@chemas.com
www.chemas.com

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Pty Ltd.
 - The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
 - For authorised copying within Australia please check that your institution has a licence from Copyright Agency Limited. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.
 - Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.
 - Schools which purchase a licence to use this material may distribute this electronic file to the students at the school for their exclusive use. This distribution can be done either on an Intranet Server or on media for the use on stand-alone computers.
 - Schools which purchase a licence to use this material may distribute this printed file to the students at the school for their exclusive use.
-
- **The Word file (if supplied) is for use ONLY within the school**
 - **It may be modified to suit the school syllabus and for teaching purposes.**
 - **All modified versions of the file must carry this copyright notice**
 - **Commercial used of this material is expressly prohibited**

Question 1

$$y = \frac{x^3 + b^3}{x^2} = \frac{x^3}{x^2} + \frac{b^3}{x^2} = x + \frac{b^3}{x^2}$$

The denominator is zero when $x = 0$, so $x = 0$ is a vertical asymptote.

As $x \rightarrow \pm \infty$ $y \rightarrow x$, so $y = x$ is an oblique asymptote.

Both $x = 0$ and $y = x$ are asymptotes.

Answer: D.

Question 2

The ellipse has centre $(-3, 1)$ semi-major axis is 3, semi-minor is 1, the general form is


$$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$$

with $h = -3$ $k = 1$ $a = 3$ and $b = 1$ so the equation is $\frac{(x+3)^2}{9} + \frac{(y-1)^2}{1} = 1$.


Answer: E.

Question 3

Given $f(x) = a(x - \alpha)(x + \beta)$ where $a > 0$ this graph crosses the x -axis at $x = \alpha$ and $x = -\beta$.



the graph of $\frac{1}{f(x)}$



has vertical asymptotes at $x = \alpha$ and $x = -\beta$. So alternatives A. and B. are incorrect.

$$f(x) = a(x^2 + (\beta - \alpha)x + \alpha\beta)$$

$$f'(x) = a(2x + (\beta - \alpha)) = 0 \text{ when}$$

$$2x + (\beta - \alpha) = 0$$

$$2x = (\alpha - \beta)$$

$$x = \frac{\alpha - \beta}{2}$$

So $\frac{1}{f(x)}$ has a local maximum at $x = \frac{\alpha - \beta}{2}$.

Answer: C.

Question 4

The range of $y = \cos^{-1}x$ is $[0, \pi]$

The range of $f(x) = a + b \cos^{-1}(cx)$ is dilated by b and translated by a ,

its range is $[a, a + b\pi]$

(the value of c has no effect on the range).

Answer: E.

Question 5

Given $|a| = 7$ $|b| = 3$ $ab = 2$, now

$$\begin{aligned} |a - b|^2 &= (a - b) \cdot (a - b) \\ &= a \cdot a - b \cdot a - a \cdot b + b \cdot b \\ &= |a|^2 - 2ab + |b|^2 \\ &= 49 - 4 + 9 \end{aligned}$$

$$|a - b|^2 = 54$$

so $|a - b| = \sqrt{54}$

Answer: C.

Question 6

Given $P(2, -8, 10)$ $Q(x, 4, -5)$ then

$$\overrightarrow{OP} = 2\mathbf{i} - 8\mathbf{j} + 10\mathbf{k} \quad \overrightarrow{OQ} = x\mathbf{i} + 4\mathbf{j} - 5\mathbf{k}$$

$$\overrightarrow{PQ} = \overrightarrow{OQ} - \overrightarrow{OP} = (x - 2)\mathbf{i} + 12\mathbf{j} - 15\mathbf{k}$$

Checking each alternative:

A When $x = 2$ $\overrightarrow{PQ} = 12\mathbf{j} - 15\mathbf{k}$ so it is parallel to the YZ plane.

Alternative A is correct.

B When $x = -1$, $\overrightarrow{OP} = -2\overrightarrow{OQ}$, so $\overrightarrow{OP}$ and $\overrightarrow{OQ}$ are linearly dependent

Alternative B is incorrect.

C When $x = -3$, $|\overrightarrow{OQ}| = \sqrt{9 + 16 + 25} = \sqrt{50} = 5\sqrt{2}$

D When $x = 3$, $|\overrightarrow{OQ}| = \sqrt{9 + 16 + 25} = \sqrt{50} = 5\sqrt{2}$. So alternatives C and D are correct.

E If $\overrightarrow{OP}$ is perpendicular to $\overrightarrow{OQ}$,

$$\overrightarrow{OP} \cdot \overrightarrow{OQ} = 0$$


$$\therefore 2x - 32 - 50 = 0$$

So alternative E is correct.

$$\therefore 2x = 82 \Rightarrow x = 41$$

Answer: B.

Question 7



$$\vec{OA} = 100 \sin 30^\circ \vec{i} - 100 \cos 30^\circ \vec{j} = 50\vec{i} - 50\sqrt{3}\vec{j}$$


$$\vec{AB} = 15\vec{k}$$

$$\therefore \vec{OB} = \vec{OA} + \vec{AB} = 50\vec{i} - 50\sqrt{3}\vec{j} + 15\vec{k}$$

Answer: B.

Question 8

If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ the vectors $\vec{a}$, $\vec{b}$ and $\vec{c}$ form the sides of a triangle.



The angles given are the exterior angles.

So the interior angles are $\pi - \alpha$, $\pi - \beta$ and $\pi - \gamma$

$$\Rightarrow \pi - \alpha + \pi - \beta + \pi - \gamma = \pi$$


$$\Rightarrow 3\pi - \pi = \alpha + \beta + \gamma$$

$$\Rightarrow \alpha + \beta + \gamma = 2\pi$$

Answer: E.

Question 9

Given $\sec(x) = -5$ so $\sec(x) = \frac{1}{\cos(x)} = -5$ so $\cos(x) = -\frac{1}{5}$, but $\frac{\pi}{2} < x < \pi$ so x is the 2nd quadrant.



by Pythagoras' theorem: $1^2 + b^2 = 5^2$

$$b^2 = 25 - 1 = 24$$

$$b = \sqrt{24} = \sqrt{4 \times 6} = 2\sqrt{6}$$

$$\text{Now } \operatorname{cosec}(x) = \frac{1}{\sin(x)} = \frac{1}{\frac{2\sqrt{6}}{5}} = \frac{5}{2\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}} = \frac{5\sqrt{6}}{12}$$

since $\sin(x)$ is positive in the 2nd quadrant.

Answer: E.

Question 10

Let $\underline{a} = 2\underline{i} - 3\underline{j} + 4\underline{k}$, the scalar resolute of $\underline{a}$ in the direction of $\underline{b}$ is $\underline{a} \cdot \hat{\underline{b}} = 2\sqrt{6}$.

The vector resolute of $\underline{a}$ perpendicular to $\underline{b}$ is $\underline{a} - (\underline{a} \cdot \hat{\underline{b}})\hat{\underline{b}} = \underline{j} + 2\underline{k}$.

$$2\underline{i} - 3\underline{j} + 4\underline{k} - 2\sqrt{6} \hat{\underline{b}} = \underline{j} + 2\underline{k}$$

$$2\sqrt{6} \hat{\underline{b}} = (2\underline{i} - 3\underline{j} + 4\underline{k}) - (\underline{j} + 2\underline{k})$$

$$2\sqrt{6} \hat{\underline{b}} = 2\underline{i} - 4\underline{j} + 2\underline{k}$$

$$\text{so } \hat{\underline{b}} = \frac{1}{\sqrt{6}}(\underline{i} - 2\underline{j} + \underline{k})$$

Answer: C.

Question 11

$P(z) = z^3 + az^2 + bz + c$ given that $P(\alpha - i\beta) = 0$ and a, b and c are real then

- A $P(\alpha + i\beta) = 0$ is correct by the conjugate root theorem.
 B $P(z)$ has three roots (1 pair of complex conjugates and one root). This is correct.
 C This is correct.

D Now $u = \alpha + i\beta$ $v = \alpha - i\beta$

$$u + v = 2\alpha \quad uv = \alpha^2 - i^2\beta^2 = \alpha^2 + \beta^2 \text{ so}$$

$$z^2 - (u + v)z + uv = 0$$

$$z^2 - (\text{sum of the roots})z + (\text{product of the roots}) = 0$$

$$\text{so} \quad z^2 - 2\alpha z + (\alpha^2 + \beta^2) = 0 \text{ is a factor of } P(z).$$

Option D is INCORRECT.

E is correct since

$$z^3 + az^2 + bz + c = 0$$

$$= (z^2 - 2\alpha z + (\alpha^2 + \beta^2))(z + d) = 0$$

$$\text{so} \quad d(\alpha^2 + \beta^2) = c$$

$$d = \frac{c}{\alpha^2 + \beta^2} \text{ and } P(-d) = 0$$

Answer: D.

Question 12

The shaded region is the intersection of the inside of a circle with centre at the origin and radius 2, with the region above the line $y = -x$.

The circle is $z\bar{z} \leq r^2 = 4$ (where r is the radius).

since if $z = x + iy$ $\bar{z} = x - iy$ so

$$z\bar{z} = x^2 - i^2y^2 = x^2 + y^2 = r^2$$


The line is $y \geq -x$, if $z = x + iy$

$\text{Re}(z) = x$ and $\text{Im}(z) = y$

so $\text{Im}(y) + \text{Re}(z) \geq 0$

Answer: A.

Question 13



Now $z = -a - bi$

a and b are real positive

$$-\pi < \text{Arg}(z) \leq \pi$$

$$\tan \theta = \frac{b}{a} > 0 \quad \theta = \text{Tan}^{-1}\left(\frac{b}{a}\right)$$

$$\text{Arg}(z) = -\pi + \theta$$

$$= -\pi + \text{Tan}^{-1}\left(\frac{b}{a}\right)$$

Answer: D.

Question 14

Solving $z^3 + a^3i = 0$ since $i^2 = -1$

$$z^3 = -a^3i = a^3i^3, \text{ one answer is:}$$

$$z = ai \text{ where } a > 0$$

There are three answers one of which is D are all equally spaced around the circle. The roots are D, H and L

Answer: B.

Question 15

$$\underline{r}(t) = \cos^2(t)\underline{i} + \cos(2t)\underline{j}$$

$$x = \cos^2(t)$$

$$y = \cos(2t) = 2\cos^2(t) - 1$$

$$y = 2x - 1, \text{ this is a straight line.}$$

Answer: A.

Question 16

$$y = \text{Cos}^{-1}\left(\frac{5x}{4}\right) = \text{Cos}^{-1}\left(\frac{u}{4}\right) \text{ where } u = 5x \text{ Chain Rule}$$

$$\frac{dy}{du} = \frac{-1}{\sqrt{16-u^2}} \quad \frac{du}{dx} = 5$$

$$\frac{dy}{dx} = \frac{-5}{\sqrt{16-25x^2}}, \text{ now at } x=0 \quad m_T = \left.\frac{dy}{dx}\right|_{x=0} = f'(0) = \frac{-5}{4}$$

$$\text{at } x=0 \quad y = \text{Cos}^{-1}(0) = \frac{\pi}{2} \text{ so } P\left(0, \frac{\pi}{2}\right) \text{ and the gradient of the normal is } \frac{4}{5},$$

$$\text{so the equation of the normal is: } y - \frac{\pi}{2} = \frac{4}{5}(x - 0)$$

$$\text{or } y = \frac{4x}{5} + \frac{\pi}{2}$$

Answer: A.

Question 17

$$y = \tan^{-1}\left(\frac{3}{4x}\right) = \tan^{-1} u \text{ where } u = \frac{3}{4x} = \frac{3}{4} x^{-1}$$

$$\frac{dy}{du} = \frac{1}{1+u^2} \quad \frac{du}{dx} = -\frac{3}{4} x^{-2} = \frac{-3}{4x^2}$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \text{ (chain rule)}$$

$$= -\frac{3}{4x^2} \left(\frac{1}{1 + \frac{9}{16x^2}} \right)$$

$$= -\frac{3}{4x^2} \left(\frac{1}{\frac{16x^2+9}{16x^2}} \right)$$

$$= -\frac{3}{4x^2} \left(\frac{16x^2}{16x^2+9} \right)$$

$$= \frac{-12}{9+16x^2}$$

Answer: A.

Question 18

$$\int_0^{\frac{\pi}{4}} \cos^3(2x) \sin(2x) dx$$

$$\text{let } u = \cos(2x) \quad \frac{du}{dx} = -2 \sin(2x)$$

changing terminals, when

$$x = \frac{\pi}{4} \quad u = \cos\left(\frac{\pi}{2}\right) = 0 \quad \text{and when } x = 0 \quad u = \cos(0) = 1$$

$$= \int_1^0 u^3 \times -\frac{1}{2} \frac{du}{dx} \cdot dx = -\frac{1}{2} \int_1^0 u^3 du = \frac{1}{2} \int_0^1 u^3 du$$

Answer: E.

Question 19

$$y = (A + Bx)e^{-2x} \text{ satisfies } \frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4y = 0$$

$$\frac{dy}{dx} = Be^{-2x} - 2(A + Bx)e^{-2x}$$

$$= ((B - 2A) - 2Bx)e^{-2x}$$

$$\frac{d^2y}{dx^2} = -2((B - 2A) - 2Bx)e^{-2x} - 2Be^{-2x}$$

$$= ((4A - 4B) + 4Bx)e^{-2x}$$

$$\text{so } \frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4y$$

$$= (4A - 4B + 4Bx + 4B - 8A - 8Bx + 4A + 4Bx)e^{-2x} = 0$$

If $B = 0$ and $A = 3$ or $A = 5$ so alternatives A. and B. both satisfy the differential equation.

If $A = 0$ and $B = 2$ or if $A = 4$ and $B = 2$ so alternatives C. and D. both satisfy the differential equation.

Alternative E. **does not** satisfy the differential equation.

Answer: E.

Question 20

$y_1 = \frac{\cos(2x)}{2x-1}$ using a graphic calculator, in the Radians mode we find the graph crosses the x -axis at $x \approx 2.36$, the area is:

$$\int_2^{2.36} \frac{\cos(2x)}{2x-1} dx + \int_{2.36}^2 \frac{\cos(2x)}{2x-1} dx$$
$$= A_1 + A_2$$

but $A_1 = -0.0376$ and $A_2 = 0.0796$ the shaded area is:

$$A = |A_1| + A_2$$
$$= 0.0376 + 0.0796$$
$$= 0.1173$$

Answer: D.

Question 21

Rearranging to $\frac{dy}{dx} = \frac{\cos(2x)}{2x-1}$ with the calculator in the radians mode, using a PRGM Euler

$$Y_1 = \frac{\cos(2x)}{2x-1} \quad x_0 = 0 \quad y_0 = -1 \quad h = 0.2$$

$$y(0.4) = -1.507$$

Answer: D.

Question 22

$$\frac{d}{dx} (x \log_e(4+x)) = \log_e(4+x) + \frac{x}{4+x}$$

so it follows that:

$$\int \left(\log_e(4+x) + \frac{2x}{4+x} \right) dx = x \log_e(4+x)$$

$$\begin{aligned} \int \log_e(4+x) dx &= x \log_e(4+x) - \int \frac{x}{4+x} dx \\ &= x \log_e(4+x) - \int \left(\frac{x+4-4}{4+x} \right) dx \\ &= x \log_e(4+x) - \int \left(1 - \frac{4}{4+x} \right) dx \\ &= x \log_e(4+x) - x + 4 \log_e(4+x) \\ &= (x+4) \log_e(x+4) - x \end{aligned}$$

Answer: C.

Question 23

Since for

$$x < 0 \quad \frac{dy}{dx} < 0 \quad \text{and} \quad 0 < x < 1 \quad \frac{dy}{dx} > 0$$


while at $x = 0$ $\frac{dy}{dx} = 0$ the graph has a local minimum at $x = 0$

and at $x = 4$ $\frac{d^2y}{dx^2} = 0$

The graph has a stationary point of inflexion at $x = 4$

Answer: B.

Question 24



$$m = 12 \text{ kg} \quad P = 11.65 \text{ N}$$

$$\mu = 0.1$$

$$N = mg$$

$$N = 12 \times 9.8$$

$$= 117.6 \text{ N}$$

$$\text{Now } \mu N = 0.1 \times 117.6$$


$$= 11.76 \text{ N}$$

$$\text{since } \mu N = 11.76 > P = 11.65$$

The suitcase remains at rest.

Answer: E.

Question 25



$$F_1 = -3\hat{i} + 3\hat{j} \quad |F_1| = 3\sqrt{2}$$

$$F_2 = -4\hat{i} - 4\hat{j} \quad |F_2| = 4\sqrt{2}$$

$$R = F_1 + F_2 = -7\hat{i} - \hat{j} = 2a$$

so $a = -\frac{1}{2}(7\hat{i} - \hat{j})$

$$|a| = \frac{1}{2}\sqrt{49+1} = \frac{\sqrt{50}}{2} = \frac{5\sqrt{2}}{2}$$

Answer: B.

Question 26

Resolving vertically gives: $T_1 \sin\beta + T_2 \sin\alpha - mg = 0$
so alternative B. is correct.

Resolving horizontally gives: $T_1 \cos\beta - T_2 \cos\alpha = 0$

so $T_1 \cos\beta = T_2 \cos\alpha$

$$\frac{T_1}{T_2} = \frac{\cos\alpha}{\cos\beta} = \frac{\frac{L_2}{a}}{\frac{L_1}{a}} = \frac{L_2}{L_1} \text{ so alternatives A. and C. are correct.}$$

The vector equation $T_1 + T_2 + mg = 0$ is true.

Alternative D. is correct. Alternative E. is INCORRECT.

The shorter string carries more tension.

Answer: E.

Question 27

Let $y_1 = 4e^{-\frac{x}{2}}$ and $y_2 = 3 \sin\left(\frac{x}{2}\right)$ the volume is

$$V = \pi \int_0^b (y_1^2 - y_2^2) dx = \pi \int_0^b \left(16e^{-x} - 9\sin^2\left(\frac{x}{2}\right)\right) dx$$

Answer: A.

Question 28

Since $v > 0$ for $0 \leq t < c$ and $v = 0$ for $t = c$
 $v < 0$ for $t > c$ Alternative B. is correct.

Answer: B.

Question 29

Given the initial conditions $t = 0$ $x = 0$ $v = v_0$

Now $\frac{dv}{dx} = 1$ so $v \frac{dv}{dx} = a = v$ Alternative A. is correct.

Since $\frac{dv}{dx} = 1$ integrating wrt x

$$v = \int 1 dx = x + c \text{ but } v = v_0 \text{ when } x = 0$$

so $c = v_0$ and $v = x + v_0$ Alternative B. is correct.

Since $a = \frac{dv}{dt} = v$ this has the solution $v = v_0 e^t$ Alternative C. is correct.

Alternative D. is the **constant** acceleration formula $v^2 = v_0^2 + 2vx$ is FALSE.

Alternative E. is correct since $v = \frac{dx}{dt} = v_0 e^t$ integrating wrt t

$$x = \int v_0 e^t dt = v_0 e^t + c$$

but when $t = 0$ $x = 0$ so $0 = v_0 + c$ so $c = -v_0$

$$\text{and } x = v_0 (e^t - 1)$$

Answer: D.

Question 30

Resolving perpendicular to the incline gives

$$(1) \quad N - mg \cos \theta = 0$$

resolving downwards parallel to the plane gives

$$(2) \quad mg \sin \theta - \mu N = ma$$

from (1) $N = mg \cos \theta$ into (2)

$$mg \sin \theta - \mu mg \cos \theta = ma$$

$$a = g(\sin \theta - \mu \cos \theta) \quad \text{Alternative A. and B. are correct.}$$

The acceleration is constant, using $u = 0 \quad t = T \quad s = D$

$$v^2 = u^2 + 2as \quad \text{gives}$$

$$v^2 = 0 + 2g(\sin \theta - \mu \cos \theta) D$$

$$\text{so} \quad v = \sqrt{2gD(\sin \theta - \mu \cos \theta)} \quad \text{Alternative C. is correct.}$$

$$\text{Using: } s = \left(\frac{u + v}{2} \right) t \quad \text{gives}$$

$$D = \left(\frac{0 + V}{2} \right) T = \frac{VT}{2}$$

$$\text{so} \quad T = \frac{2D}{V}$$

Alternative D. is correct.

Alternative E. is false, terminal or limiting velocity occurs in a situation when falling vertically with air resistance proportional to the velocity.

Answer: E.

Question 1

$$\text{Let } u = \sqrt{x} = x^{\frac{1}{2}} \quad \frac{du}{dx} = \frac{1}{2} x^{-\frac{1}{2}} = \frac{1}{2\sqrt{x}} \quad \text{so } 2 \, du = \frac{1}{\sqrt{x}} \, dx$$


$$\text{terminals } x = 4 \quad u = 2$$

$$x = 0 \quad u = 0$$

$$\begin{aligned} \int_0^4 \frac{\cos \sqrt{x}}{\sqrt{x}} \, dx &= 2 \int_0^2 \cos(u) \, du \\ &= 2 \sin u \Big|_0^2 = 2 \sin(2) - 2 \sin(0) \\ &= 2 \sin(2) \end{aligned}$$

Question 2

i. $x = 2t - \sin(2t) \quad y = 1 - \cos(2t) \quad 0 \leq t \leq 2\pi$



ii. $\underline{r}(t) = (2t - \sin(2t))\underline{i} + (1 - \cos(2t))\underline{j} \quad 0 \leq t \leq 2\pi$

$$\dot{\underline{r}}(t) = (2 - 2 \cos(2t))\underline{i} + 2 \sin(2t)\underline{j}$$

$$|\dot{\underline{r}}(t)| = \sqrt{(2 - 2 \cos 2t)^2 + (2 \sin 2t)^2}$$

$$= \sqrt{4 - 8 \cos(2t) + 4 \cos^2(2t) + 4 \sin^2(2t)}$$

$$= \sqrt{8 - 8 \cos(2t)}$$

$$= \sqrt{8(1 - \cos(2t))}$$

$$\text{since } \cos(2A) = 1 - 2 \sin^2(A)$$


$$= \sqrt{8(2 \sin^2(t))}$$

$$\text{and } 2 \sin^2(A) = 1 - \cos(2A)$$

$$= 4 \sin t$$

$$\text{so } c = 4$$

Question 3



Now $\overrightarrow{OA} = \overrightarrow{BO} = a$ and $\overrightarrow{OC} = \overrightarrow{DO} = c$

$$\overrightarrow{AC} = \overrightarrow{AO} + \overrightarrow{OC} = c - a$$

$$\overrightarrow{AD} = \overrightarrow{AO} + \overrightarrow{OD} = -a - c$$

$$\overrightarrow{CB} = \overrightarrow{CO} + \overrightarrow{OB} = -a - c \Rightarrow \overrightarrow{AD} = \overrightarrow{CB}$$

consider

$$\begin{aligned} \overrightarrow{AC} \cdot \overrightarrow{CB} &= (c - a) \cdot (-a - c) \\ &= -c \cdot a + a \cdot a - c \cdot c + a \cdot c \\ &= |a|^2 - |c|^2 \\ &= 0 \text{ since } |a| = |c| \text{ (both are radii of the circle)} \end{aligned}$$

so $\overrightarrow{AC}$ is perpendicular to $\overrightarrow{CB}$ and $\overrightarrow{AD} = \overrightarrow{CB}$

$ABCD$ is rectangle

Question 4

i. $u = 2\sqrt{2}(1 - i) \quad \text{Arg}(u) = -\frac{\pi}{4}$

so $u = 4 \text{cis}\left(-\frac{\pi}{4}\right)$

And $\arg(u^8) = 8 \times -\frac{\pi}{4} = -2\pi$

But $\text{Arg}(u^8) = -2\pi + 2\pi = 0$

ii. $u^2 = \left(4 \text{cis}\left(-\frac{\pi}{4}\right)\right)^2$

$= 16 \text{cis}\left(-\frac{\pi}{2}\right)$

$= -16i$

iii. $z^3 + 16iz = 0$


$z(z^2 + 16i) = 0$

$z = 0$ or $z^2 = -16i$ hence from i. and ii.

$z = 2\sqrt{2}(1 - i) = 4 \text{cis}\left(-\frac{\pi}{4}\right) = z_1$

$z = 2\sqrt{2}(-1 + i) = 4 \text{cis}\left(\frac{3\pi}{4}\right) = z_2$

note that $z_2 = -z_1$ these are **not** conjugates.



Question 5

a. $a = -2i + yj + 5k$

$$|a| = \sqrt{4 + y^2 + 25} = 6$$

$$\sqrt{29 + y^2} = 6$$

$$29 + y^2 = 36$$

$$y^2 = 7$$

$$y = \pm\sqrt{7} \quad \text{both } \pm \text{ are both acceptable answers.}$$

b. $\cos\beta = -\frac{\sqrt{7}}{6} = \frac{y}{|a|} = \frac{y}{\sqrt{29 + y^2}}$

$$6y = -\sqrt{7} (\sqrt{29 + y^2})$$

$$36y^2 = 7(29 + y^2)$$

$$= 7 \times 29 + 7 \times y^2$$

$$29y^2 = 7 \times 29$$

$$y = \pm\sqrt{7} \quad \text{but}$$

$$y = -\sqrt{7} \quad \text{since } y < 0 \quad \text{it is an obtuse angle, there is only one answer.}$$


Question 6

to find the first x intercept

$$\cos\left(\frac{2x}{3}\right) = 0$$

$$\frac{2x}{3} = \frac{\pi}{2}$$

$$x = \frac{3\pi}{4}$$




Now the volume

$$\begin{aligned}
 V_x &= \pi \int_a^b y^2 dx \\
 &= \pi \int_0^{\frac{3\pi}{4}} 16 \cos^2\left(\frac{2x}{3}\right) dx \\
 &= 8\pi \int_0^{\frac{3\pi}{4}} \left(1 + \cos\left(\frac{4x}{3}\right)\right) dx \\
 &= 8\pi \left[x + \frac{3}{4} \sin\left(\frac{4x}{3}\right) \right]_0^{\frac{3\pi}{4}} \\
 &= 8\pi \left[\frac{3\pi}{4} + \frac{3}{4} \sin(\pi) - \left(0 + \frac{3}{4} \sin 0\right) \right] \\
 &= 6\pi^2
 \end{aligned}$$

Question 7

i.



ii. isolating the particles and resolving around each individually,

resolving perpendicular to the plane around the mass m_1

$$N_1 - m_1 g \cos \alpha = 0 \quad \text{so that (1)} \quad N_1 = m_1 g \cos \alpha$$

resolving perpendicular to the plane around the mass m_2

$$N_2 - m_2 g \cos \alpha = 0 \quad \text{so that (2)} \quad N_2 = m_2 g \cos \alpha$$

resolving parallel and up the plane around the mass m_1 gives

$$T + \mu_1 N_1 - m_1 g \sin \alpha = 0 \quad \text{using (1) gives (3)} \quad T = m_1 g (\sin \alpha - \mu_1 \cos \alpha)$$

resolving parallel and up the plane around the mass m_2 gives

$$\mu_2 N_2 - T - m_2 g \sin \alpha = 0 \quad \text{using (2) gives (4)} \quad T = m_2 g (\mu_2 \cos \alpha - \sin \alpha)$$

Equating (3) and (4)

$$m_1 (\sin \alpha - \mu_1 \cos \alpha) = m_2 (\mu_2 \cos \alpha - \sin \alpha)$$

$$(m_1 + m_2) \sin \alpha = (m_1 \mu_1 + m_2 \mu_2) \cos \alpha$$

$$\tan \alpha = \frac{m_1 \mu_1 + m_2 \mu_2}{m_1 + m_2} \quad \text{shown.}$$

END OF SUGGESTED SOLUTIONS
2004 Specialist Mathematics Trial Examination 1

KILBAHA MULTIMEDIA PUBLISHING
PO BOX 2227
KEW VIC 3101
AUSTRALIA

TEL: (03) 9817 5374
FAX: (03) 9817 4334
chemas@chemas.com
www.chemas.com