

Year 2005

VCE

Specialist Mathematics

Trial Examination 2



KILBAHA MULTIMEDIA PUBLISHING
PO BOX 2227
KEW VIC 3101
AUSTRALIA

TEL: (03) 9817 5374
FAX: (03) 9817 4334
chemas@chemas.com
www.chemas.com

© Kilbaha Multimedia Publishing 2005

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Multimedia Publishing.
 - The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
 - For authorised copying within Australia please check that your institution has a licence from Copyright Agency Limited. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.
 - Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.
 - Schools which purchase a licence to use this material may distribute this electronic file to the students at the school for their exclusive use. This distribution can be done either on an Intranet Server or on media for the use on stand-alone computers.
 - Schools which purchase a licence to use this material may distribute this printed file to the students at the school for their exclusive use.
-
- **The Word file (if supplied) is for use ONLY within the school**
 - **It may be modified to suit the school syllabus and for teaching purposes.**
 - **All modified versions of the file must carry this copyright notice**
 - **Commercial use of this material is expressly prohibited**

STUDENT NUMBER

Figures

Words

VICTORIAN CERTIFICATE OF EDUCATION

2005

SPECIALIST MATHEMATICS

Trial Written Examination 2 (Analysis Task)

Reading time: 15 minutes

Total writing time: 1 hour 30 minutes

QUESTION AND ANSWER BOOK

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
5	5	60

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers, a protractor, set-squares, aids for curve sketching, up to four pages (two A4 sheets) of pre-written notes (typed or handwritten) and an approved scientific and/or graphics calculator (memory may be retained).
- Students are NOT permitted to bring into the examination room: blank sheets of paper and/or whiteout liquid/tape.

Materials supplied

- Question and answer book of 20 pages with a detachable sheet of miscellaneous formulas in the centrefold.
- Working space is provided throughout the book.

Instructions

- Detach the formula sheet from the centre of this book during reading time.
- Write your **student number** in the space provided above on this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other electronic communication devices into the examination room.

SPECIALIST MATHEMATICS

Written examinations 1 and 2

FORMULA SHEET

Directions to students

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

Specialist Mathematics Formulae

Mensuration

area of a trapezium: $\frac{1}{2}(a+b)h$

curved surface area of a cylinder: $2\pi rh$

volume of a cylinder: $\pi r^2 h$

volume of a cone: $\frac{1}{3}\pi r^2 h$

volume of a pyramid: $\frac{1}{3}Ah$

volume of a sphere: $\frac{4}{3}\pi r^3$

area of triangle: $\frac{1}{2}bc \sin(A)$

sine rule: $\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$

cosine rule: $c^2 = a^2 + b^2 - 2ab \cos(C)$

Coordinate geometry

ellipse: $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$

hyperbola: $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$

Circular (trigonometric) functions

$$\cos^2(x) + \sin^2(x) = 1$$

$$1 + \tan^2(x) = \sec^2(x)$$

$$\sin(x+y) = \sin(x)\cos(y) + \cos(x)\sin(y)$$

$$\sin(x-y) = \sin(x)\cos(y) - \cos(x)\sin(y)$$

$$\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$$

$$\cos(x-y) = \cos(x)\cos(y) + \sin(x)\sin(y)$$

$$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$$

$$\tan(x+y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$$

$$\tan(x-y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$$

$$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$$

$$\sin(2x) = 2\sin(x)\cos(x)$$

$$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$$

function	Sin^{-1}	Cos^{-1}	Tan^{-1}
domain	$[-1, 1]$	$[-1, 1]$	R
range	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$[0, \pi]$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Algebra (Complex Numbers)

$$z = x + yi = r(\cos \theta + i \sin \theta) = r \text{ cis } \theta$$

$$|z| = \sqrt{x^2 + y^2} = r$$

$$-\pi < \text{Arg } z \leq \pi$$

$$z_1 z_2 = r_1 r_2 \text{ cis } (\theta_1 + \theta_2)$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} \text{ cis } (\theta_1 - \theta_2)$$

$$z^n = r^n \text{ cis } (n\theta) \text{ (de Moivre's theorem)}$$

Vectors in two and three dimensions

$$\underline{r} = x\underline{i} + y\underline{j} + z\underline{k}$$

$$|\underline{r}| = \sqrt{x^2 + y^2 + z^2} = r$$

$$\underline{r}_1 \cdot \underline{r}_2 = r_1 r_2 \cos \theta = x_1 x_2 + y_1 y_2 + z_1 z_2$$

$$\dot{\underline{r}} = \frac{d\underline{r}}{dt} = \frac{dx}{dt} \underline{i} + \frac{dy}{dt} \underline{j} + \frac{dz}{dt} \underline{k}$$

Mechanics

momentum: $\underline{p} = m\underline{v}$

equation of motion: $\underline{R} = m\underline{a}$

sliding friction: $F \leq \mu N$

constant (uniform) acceleration:

$$v = u + at$$

$$s = ut + \frac{1}{2}at^2$$

$$v^2 = u^2 + 2as$$

$$s = \frac{1}{2}(u + v)t$$

acceleration:

$$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx} \left(\frac{1}{2}v^2 \right)$$

Calculus

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\frac{d}{dx}(e^{ax}) = ae^{ax}$$

$$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$$

$$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$$

$$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$$

$$\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$$

$$\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\cos^{-1}(x)) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{1+x^2}$$

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, \quad n \neq -1$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$$

$$\int \frac{1}{x} dx = \log_e(x) + c, \text{ for } x > 0$$

$$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$$

$$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$$

$$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1}\left(\frac{x}{a}\right) + c, \quad a > 0$$

$$\int \frac{-1}{\sqrt{a^2 - x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, \quad a > 0$$

$$\int \frac{a}{a^2 + x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$$

product rule: $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$

quotient rule: $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

chain rule: $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

mid-point rule: $\int_a^b f(x) dx \approx (b-a) f\left(\frac{a+b}{2}\right)$

trapezoidal rule: $\int_a^b f(x) dx \approx \frac{1}{2}(b-a)(f(a) + f(b))$

Euler's method

If $\frac{dy}{dx} = f(x)$, $x_0 = a$ and $y_0 = b$, then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + hf(x_n)$

Take the acceleration due to gravity to have magnitude $g \text{ m/s}^2$, where $g = 9.8$

Question 1

- a.** Jared sets out on a walk from a point O and walks 30 metres east to a point A and then $10\sqrt{10}$ metres on a bearing north α east where $\alpha = \tan^{-1}\left(\frac{1}{3}\right)$ to a point B , all the time on horizontal ground.
- i.** Taking $\underline{i}$ as a unit vector in the east direction, $\underline{j}$ as a unit vector in the north direction, find the vector $\overrightarrow{OB}$ in terms of $\underline{i}$ and $\underline{j}$.


1 mark

- ii.** Using a suitable scalar product, find the bearing of B from O in degrees and minutes.

2 marks

Question 1 (continued)

- b. $OABC$ is a trapezium and $OARS$ is a square as shown. Both $\overrightarrow{AB}$ and $\overrightarrow{OC}$ are inclined at an angle of α to the vertical where $\alpha = \tan^{-1}\left(\frac{1}{3}\right)$



Let $\overrightarrow{OA} = \underline{a}$ and $\overrightarrow{OC} = \underline{c}$.

- i. Show that $\overrightarrow{OB} = \frac{1}{3}(5\underline{a} + 3\underline{c})$.

1 mark

Question 1 (continued)

b.

- ii. Let M be the mid-point of $\overrightarrow{OA}$ and P be a point on MC such that the ratio of the distances $MP : MC = \frac{3}{13}$. Show that $\overrightarrow{MP} = \frac{3}{26}(2\mathbf{c} - \mathbf{a})$.

2 marks

- iii. Hence find the vector $\overrightarrow{OP}$ and show that O, P and B are collinear.
What is the ratio of the distances $OP : OB$?

3 marks


Question 1 (continued)

- c. A drinking trough has a length of 150 cm and its cross-sectional face is in the shape of a trapezium, with a height of 30 cm and with lengths 30 and 50 cm.

Both sloping edges are at an angle of α to the vertical, where $\alpha = \tan^{-1}\left(\frac{1}{3}\right)$ as

shown in the diagram below. The trough contains water to a height of h cm.

The water, however, leaks out through a crack along the base of the trough at a rate equal to $80\sqrt{h}$ cm³/min.



- i. Show that the differential equation for the height of water h cm in the trough,

where $0 \leq h \leq 30$, at a time t minutes, is given by $\frac{dh}{dt} = -\frac{4\sqrt{h}}{5(45+h)}$

2 marks

Question 1 (continued)

c.

ii. When the trough is filled with water to a height of 25 cm, the water starts to leak out through the crack along the base of the trough. Hence, find using calculus, the exact time in hours before the trough is empty.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

3 marks

Total 14 marks

Question 2

Let $\alpha = \tan^{-1}\left(\frac{1}{3}\right)$, $\beta = \tan^{-1}\left(\frac{1}{2}\right)$, $u = 3+i$ and $v = 2+i$

- a.** Find the exact value of $\cos(\alpha + \beta)$.

2 marks

- b.** Show that $\sin(2\beta) = \frac{\text{Im}(v^2)}{|v^2|}$.

2 marks

Question 2 (continued)

- c. Find $\text{Arg}(uv)$ and explain how this shows that $\alpha + \beta = \frac{\pi}{4}$.

2 marks

- d. $P(z) = z^3 + az^2 + bz + 20 = 0$, where a and b are real numbers and $P(u) = 0$.
Find the values of a and b and state all the roots of $P(z) = 0$.

2 marks

Question 2 (continued)

- e. If $Q(z) = z^3 - (2+i)z^2 + 5z - 10 - 5i = 0$, show that $Q(v) = 0$ and, hence, find all the values of z where $Q(z) = 0$.

2 marks

Total 10 marks

Question 3

- a. Ashley is driving his sports car along a straight road at a speed of 10 m/s when he brakes. Assuming a constant retardation of 2.5 m/s^2 , find how long in seconds and the distance travelled in metres, before the speed of the car is reduced to 5 m/s.

2 marks

- b. Ashley's sports car has a mass 800 kg and on another day he is moving along a level section of a street at a speed of 10 m/s when he brakes. The total resistance forces are $40v^2$ newtons, where $v \text{ m/s}$ is the speed of the sports car at a time t seconds, and x is its distance from the point where Ashley applied the brakes.
- i. By choosing an appropriate form for the acceleration, show that a differential equation relating v to x is $\frac{dv}{dx} = -\frac{v}{20}$.

1 mark

Question 3 (continued)

b.

- ii.** In this situation, express x in terms of v and find the exact distance travelled in metres, before the speed of the car is reduced to 5 m/s.

[illegible]

3 marks

Question 3 (continued)

- b.**
iii. Find the exact time in seconds to reduce the speed of the car from 10 m/s to 5 m/s.

3 marks

Question 3 (continued)**b.**

- iv.** Express the velocity v in terms of the time t and sketch the velocity time graph on the axes below, marking in suitable scales.



2 marks

Total 11 marks

Question 4

- a. The position vector of a particle at a time t seconds is given by

$$\underline{r}(t) = 2 \cot(t) \underline{i} + (1 - \cos(2t)) \underline{j}, \text{ for } t \geq 0.$$

Find the Cartesian equation of the curve.

2 marks


Question 4 (continued)

- b.** Given the function $f: R \rightarrow R$ where $f(x) = \frac{8}{x^2 + 4}$,
- i.** Find the exact coordinates of the turning point and the inflexion points for the function.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

4 marks

- ii. Sketch the graph of the function $f(x) = \frac{8}{x^2 + 4}$ on the axes below, marking in suitable scales.



1 mark


Question 4 (continued)**b.**

- iii.** An arched doorway at the entrance to a building has the shape given by the area under the curve $y = \frac{8}{x^2 + 4}$, between the x -axis and the lines $x = \pm \frac{2\sqrt{3}}{3}$. Find, using calculus, the exact area of the doorway.

3 marks

Question 4 (continued)

- c. Given the function $g(x) = \frac{8}{x^2 - 4}$,
- i. State the maximal domain and range of the function $g(x)$ and the equations of all the asymptotes. Sketch the graph of $g(x) = \frac{8}{x^2 - 4}$ on the axes below, marking in a suitable scale.



2 marks

Question 4 (continued)

c.

- ii. A section of piping is formed when the area bounded by the curve $y = \frac{8}{x^2 - 4}$, the x -axis and the lines $x = 6$ and $x = 8$ is rotated 360° about the x -axis. Find, using calculus, the exact volume of this section of piping.

[illegible]

4 marks


Total 16 marks

Question 5

Lilly is waiting for the delivery of some white goods, which were recently purchased.

- a.** A refrigerator has a mass of 102 kg and is on a loading bay from a delivery station. The loading bay is inclined at an angle of 40° to the horizontal as shown in the diagram below. When a delivery man exerts a force of P newtons on a rope up and parallel to the loading bay, the refrigerator is just prevented from sliding down the loading bay.
- i.** On the diagram below mark in all the forces acting on the refrigerator.


1 mark



- ii.** If the coefficient of friction between the refrigerator and the loading bay is 0.25, find the value of P correct to two decimal places.

3 marks

- b. A dishwasher has a mass 52 kg and is now on the loading bay from the delivery station. The loading bay is still inclined at an angle of 40° to the horizontal as shown in the diagram below. When the delivery man now exerts an upwards force of 300 newtons on a rope inclined at an angle of 35° to the loading bay, the dishwasher is moving down the loading bay with an acceleration of 0.5 m/s^2 .
- i. On the diagram below mark in all the forces acting on the dishwasher.



4 marks
Total 9 marks

WORKING SPACE

