



Victorian Certificate of Education 2006

SUPERVISOR TO ATTACH PROCESSING LABEL HERE

STUDENT NUMBER

Figures

Words

Letter

--

SPECIALIST MATHEMATICS

Written examination 1

Friday 27 October 2006

Reading time: 3.00 pm to 3.15 pm (15 minutes)

Writing time: 3.15 pm to 4.15 pm (1 hour)

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
9	9	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers.
- Students are not permitted to bring into the examination room: notes of any kind, a calculator of any type, blank sheets of paper and/or white out liquid/tape.

Materials supplied

- Question and answer book of 10 pages with a detachable sheet of miscellaneous formulas in the centrefold.
- Working space is provided throughout the book.

Instructions

- Detach the formula sheet from the centre of this book during reading time.
- Write your **student number** in the space provided above on this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

This page is blank

Instructions

Answer **all** questions in the spaces provided.

A decimal approximation will not be accepted if an **exact** answer is required to a question.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Take the **acceleration due to gravity** to have magnitude $g \text{ m/s}^2$, where $g = 9.8$.

Question 1

Consider the relation $2xy - 9y^2 + 9 = 0$.

- a. Find an expression for $\frac{dy}{dx}$ in terms of x and y .

2 marks


- b. Hence find the exact value of $\frac{dy}{dx}$ when $y = 1$.

2 marks

TURN OVER

Question 3

- a. Sketch the graph with equation $y = \frac{36}{2x^2 - 18}$, clearly indicating the location of any asymptotes and intercepts with the axes.



3 marks

- b. Find the exact area bounded by $y = \frac{36}{2x^2 - 18}$, the x-axis and the lines $x = -2$ and $x = 2$.


4 marks

TURN OVER

Question 4

A model glider with a mass of 4 kg is suspended from the roof of a wind tunnel by a thin wire. A light wind exerts a horizontal force of magnitude $\frac{g}{2}$ newtons on the model glider.

- a. Let T newtons be the magnitude of the tension in the suspending wire. Clearly label all forces acting on the model glider on the following diagram.



1 mark

- b. Calculate the value of T , giving your answer in the form $\frac{g\sqrt{a}}{b}$ where a and b are positive integers.

2 marks

Question 5

- a.** Show that $\tan\left(\frac{\pi}{8}\right) = \sqrt{2} - 1$.

4 marks

- b.** If $y = \tan^{-1}(x-1) + a \tan\left(\frac{\pi}{8}\right)$, where a is a real constant, find the minimum value of a for which $y > 0$ for all x .

2 marks

Question 6

The region in the first quadrant enclosed by the coordinate axes, the graph with equation $y = e^{-x}$ and the straight line $x = a$ where $a > 0$, is rotated about the x -axis to form a solid of revolution.

- a.** Express the volume of the solid of revolution as a definite integral.

1 mark

- b.** Calculate the volume of the solid of revolution, in terms of a .

1 mark

- c.** Find the exact value of a if the volume is $\frac{5\pi}{18}$ cubic units.

2 marks


Question 7

The position vector of a moving particle is given by $\underline{r}(t) = \sqrt{t-2} \underline{i} + 2t \underline{j}$ for $2 \leq t \leq 6$.

- a. Find the cartesian equation of the path followed by the particle.

2 marks

- b. Sketch the path of the particle on the axes provided.



2 marks

Question 8

Find an antiderivative of $\frac{2+6x}{\sqrt{4-x^2}}$.

4 marks

Question 9

a. Express $1 + \sqrt{3}i$ in polar form.

1 mark

b. Solve the quadratic equation $z^2 + 2z - \sqrt{3}i = 0$, expressing your answers in exact cartesian form.

3 marks

SPECIALIST MATHEMATICS

Written examinations 1 and 2

FORMULA SHEET

Directions to students

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

Specialist Mathematics Formulas

Mensuration

area of a trapezium: $\frac{1}{2}(a+b)h$

curved surface area of a cylinder: $2\pi rh$

volume of a cylinder: $\pi r^2 h$

volume of a cone: $\frac{1}{3}\pi r^2 h$

volume of a pyramid: $\frac{1}{3}Ah$

volume of a sphere: $\frac{4}{3}\pi r^3$

area of a triangle: $\frac{1}{2}bc \sin A$

sine rule: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

cosine rule: $c^2 = a^2 + b^2 - 2ab \cos C$

Coordinate geometry

ellipse: $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$

hyperbola: $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$

Circular (trigonometric) functions

$\cos^2(x) + \sin^2(x) = 1$

$1 + \tan^2(x) = \sec^2(x)$

$\sin(x+y) = \sin(x) \cos(y) + \cos(x) \sin(y)$

$\cos(x+y) = \cos(x) \cos(y) - \sin(x) \sin(y)$

$\tan(x+y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x) \tan(y)}$

$\cos(2x) = \cos^2(x) - \sin^2(x) = 2 \cos^2(x) - 1 = 1 - 2 \sin^2(x)$

$\sin(2x) = 2 \sin(x) \cos(x)$

$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$

$\sin(x-y) = \sin(x) \cos(y) - \cos(x) \sin(y)$

$\cos(x-y) = \cos(x) \cos(y) + \sin(x) \sin(y)$

$\tan(x-y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x) \tan(y)}$

$\tan(2x) = \frac{2 \tan(x)}{1 - \tan^2(x)}$

function	$\sin^{-1}$	$\cos^{-1}$	$\tan^{-1}$
domain	$[-1, 1]$	$[-1, 1]$	R
range	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$[0, \pi]$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Algebra (complex numbers)

$$z = x + yi = r(\cos \theta + i \sin \theta) = r \operatorname{cis} \theta$$

$$|z| = \sqrt{x^2 + y^2} = r$$

$$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$$

$$z^n = r^n \operatorname{cis}(n\theta) \text{ (de Moivre's theorem)}$$

$$-\pi < \operatorname{Arg} z \leq \pi$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$$

Calculus

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$$

$$\frac{d}{dx}(e^{ax}) = ae^{ax}$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$$

$$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$$

$$\int \frac{1}{x} dx = \log_e |x| + c$$

$$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$$

$$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$$

$$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$$

$$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$$

$$\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$$

$$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$$

$$\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$$

$$\frac{d}{dx}(\cos^{-1}(x)) = \frac{-1}{\sqrt{1-x^2}}$$

$$\int \frac{-1}{\sqrt{a^2 - x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$$

$$\frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{1+x^2}$$

$$\int \frac{a}{a^2 + x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$$

product rule:

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

quotient rule:

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

chain rule:

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Euler's method:

$$\text{If } \frac{dy}{dx} = f(x), x_0 = a \text{ and } y_0 = b, \text{ then } x_{n+1} = x_n + h \text{ and } y_{n+1} = y_n + hf(x_n)$$

acceleration:

$$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$$

$$\text{constant (uniform) acceleration: } v = u + at \quad s = ut + \frac{1}{2}at^2 \quad v^2 = u^2 + 2as \quad s = \frac{1}{2}(u+v)t$$

TURN OVER

Vectors in two and three dimensions

$$\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$$

$$|\vec{r}| = \sqrt{x^2 + y^2 + z^2} = r$$

$$\vec{r}_1 \cdot \vec{r}_2 = r_1 r_2 \cos \theta = x_1 x_2 + y_1 y_2 + z_1 z_2$$

$$\dot{\vec{r}} = \frac{d\vec{r}}{dt} = \frac{dx}{dt}\vec{i} + \frac{dy}{dt}\vec{j} + \frac{dz}{dt}\vec{k}$$

Mechanics

momentum:

$$\vec{p} = m\vec{v}$$

equation of motion:

$$\vec{R} = m\vec{a}$$

friction:

$$F \leq \mu N$$