

**THE
HEFFERNAN
GROUP**

P.O. Box 1180
Surrey Hills North VIC 3127
Phone 03 9836 5021
Fax 03 9836 5025
info@theheffernangroup.com.au
www.theheffernangroup.com.au

Students Name:.....

SPECIALIST MATHEMATICS

TRIAL EXAMINATION 1

2009

Reading Time: 15 minutes

Writing time: 1 hour

Instructions to students

This exam consists of 10 questions.
All questions should be answered.
There is a total of 40 marks available.
The marks allocated to each of the ten questions are indicated throughout.
Students may not bring any notes or calculators into the exam.
Where more than one mark is allocated to a question, appropriate working must be shown.
Where an exact answer is required to a question, a decimal approximation will not be accepted.
Unless otherwise indicated, diagrams in this exam are not drawn to scale.
The acceleration due to gravity should be taken to have magnitude $g \text{ m/s}^2$ where $g = 9.8$
Formula sheets can be found on pages 13-15 of this exam.

This paper has been prepared independently of the Victorian Curriculum and Assessment Authority to provide additional exam preparation for students. Although references have been reproduced with permission of the Victorian Curriculum and Assessment Authority the publication is in no way connected with or endorsed by the Victorian Curriculum and Assessment Authority.

© THE HEFFERNAN GROUP 2009

This Trial Exam is licensed on a non transferable basis to the purchasing school. It may be copied by the school which has purchased it. This license does not permit distribution or copying of this Trial Exam by any other party.

Question 1

Consider the function f with rule $f(x) = \cos^{-1}(3x - 2)$.

- a.** Write down the maximal domain of f .

1 mark

- b.** Find $f'(x)$.

1 mark


Question 2

Find the gradient of the relation $6x^2 - 3y^3 + 2x^2y^2 = 5$ at the point in the second quadrant where $y = 1$.

3 marks

Question 3

Sketch the graph of the relation $\frac{(x-1)^2}{4} - \frac{y^2}{25} = 1$ on the axes below. State the equations of all asymptotes and label clearly any intercepts.



4 marks

Question 4

In the right angled triangle ABC , the hypotenuse is $\vec{AC} = -4\hat{i} + \hat{j}$.


If $\vec{BC}$ is parallel to the vector $-3\hat{i} + 2\hat{j}$, find $\vec{AB}$.

3 marks

Question 5

A container of mass 20kg rests on a rough surface inclined at an angle of 30° to the horizontal.

- a. Clearly label the 3 forces, including the normal force N and the friction force F acting on the container, on the diagram below.



1 mark

A worker applies a pushing force of 200 newtons up the slope to the container which causes it to accelerate up the slope at 0.1m/s^2 .

- b. What is the coefficient of friction between the container and the surface?

2 marks

- c. What minimum force P must the worker exceed in order to cause the container to move up the slope?

2 marks

Question 6

- a. Find an antiderivative of $\sin^2\left(\frac{3x}{2}\right)$.

1 mark

- b. Evaluate $\int_0^1 \frac{5x}{\sqrt{1+x^2}} dx$.

3 marks

Question 7

Find the volume generated when the region enclosed by the curve with equation $y = \log_e(3x - 2)$, the y -axis and the lines $y = 0$ and $y = 1$ is rotated about the y -axis to form a solid of revolution.

3 marks

Question 8

Let $u = \sqrt{2}\text{cis}\left(\frac{\pi}{4}\right)$ and $w = 2\text{cis}\left(\frac{\pi}{6}\right)$.

- a.** Express u in Cartesian form.

1 mark

- b.** Express $\frac{u}{w}$ in polar form.

1 mark

- c.** Hence evaluate $\sin\left(\frac{\pi}{12}\right)$.

2 marks

- d. Given that u is a root of the equation $z^3 + z^2 - 4z + 6 = 0$, $z \in \mathbb{C}$, find the other roots of the equation.

2 marks

Question 9

A particle moves in a straight line from rest. At time t seconds its displacement is x metres from a fixed point, its velocity is v m/s and its acceleration is a m/s² where

$$a = v^2 + 3v + 2, \quad v \geq 0.$$

- a.** Find the acceleration when $t = 0$.

1 mark

- b.** Find an expression for v in terms of t .

4 marks

Question 10

- a. Show that $\cos\left(\frac{\pi}{8}\right) = \frac{\sqrt{2+\sqrt{2}}}{2}$.

3 marks

- b. Hence find the area enclosed by the graph of $y = \sin^{-1} x$, the y -axis and the line $y = \frac{\pi}{8}$.

2 marks

Specialist Mathematics Formulas

Mensuration

area of a trapezium:	$\frac{1}{2}(a+b)h$
curved surface area of a cylinder:	$2\pi rh$
volume of a cylinder:	$\pi r^2 h$
volume of a cone:	$\frac{1}{3}\pi r^2 h$
volume of a pyramid:	$\frac{1}{3}Ah$
volume of a sphere:	$\frac{4}{3}\pi r^3$
area of a triangle:	$\frac{1}{2}bc \sin A$
sine rule:	$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
cosine rule:	$c^2 = a^2 + b^2 - 2ab \cos C$

Coordinate geometry

ellipse: $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$ hyperbola: $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$

Circular (trigonometric) functions

$$\cos^2(x) + \sin^2(x) = 1$$

$$1 + \tan^2(x) = \sec^2(x)$$

$$\sin(x+y) = \sin(x)\cos(y) + \cos(x)\sin(y)$$

$$\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$$

$$\tan(x+y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$$

$$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$$

$$\sin(2x) = 2\sin(x)\cos(x)$$

$$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$$

$$\sin(x-y) = \sin(x)\cos(y) - \cos(x)\sin(y)$$

$$\cos(x-y) = \cos(x)\cos(y) + \sin(x)\sin(y)$$

$$\tan(x-y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$$

$$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$$

function	$\sin^{-1}$	$\cos^{-1}$	$\tan^{-1}$
domain	$[-1, 1]$	$[-1, 1]$	R
range	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$[0, \pi]$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Algebra (Complex numbers)

$$z = x + yi = r(\cos \theta + i \sin \theta) = r \operatorname{cis} \theta$$

$$|z| = \sqrt{x^2 + y^2} = r$$

$$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$$

$$z^n = r^n \operatorname{cis}(n\theta) \quad (\text{de Moivre's theorem})$$

$$-\pi < \operatorname{Arg} z \leq \pi$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$$

Reproduced with permission of the Victorian Curriculum and Assessment Authority, Victoria, Australia.

*This formula sheet has been copied in 2009 from the VCAA website www.vcaa.vic.edu.au
The VCAA publish an exam issue supplement to the VCAA bulletin.*

Calculus

$$\frac{d}{dx} x^n = nx^{n-1}$$

$$\frac{d}{dx} e^{ax} = ae^{ax}$$

$$\frac{d}{dx} \log_e(x) = \frac{1}{x}$$

$$\frac{d}{dx} \sin(ax) = a \cos(ax)$$

$$\frac{d}{dx} \cos(ax) = -a \sin(ax)$$

$$\frac{d}{dx} \tan(ax) = a \sec^2(ax)$$

$$\frac{d}{dx} \sin^{-1}(x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \cos^{-1}(x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \tan^{-1}(x) = \frac{1}{1+x^2}$$

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$$

$$\int \frac{1}{x} dx = \log_e |x| + c$$

$$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$$

$$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$$

$$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$$

$$\int \frac{-1}{\sqrt{a^2 - x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$$

$$\int \frac{a}{a^2 + x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$$

product rule:

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

quotient rule:

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

chain rule:

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Euler's method:

$$\text{If } \frac{dy}{dx} = f(x), x_0 = a \text{ and } y_0 = b,$$

$$\text{then } x_{n+1} = x_n + h \text{ and } y_{n+1} = y_n + hf(x_n)$$

acceleration:

$$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$$

$$\text{constant (uniform) acceleration: } v = u + at \quad s = ut + \frac{1}{2}at^2 \quad v^2 = u^2 + 2as \quad s = \frac{1}{2}(u+v)t$$

Reproduced with permission of the Victorian Curriculum and Assessment Authority, Victoria, Australia.

This formula sheet has been copied in 2009 from the VCAA website www.vcaa.vic.edu.au
The VCAA publish an exam issue supplement to the VCAA bulletin.

Vectors in two and three dimensions

$$\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$$

$$|\vec{r}| = \sqrt{x^2 + y^2 + z^2} = r \qquad \vec{r}_1 \cdot \vec{r}_2 = r_1 r_2 \cos \theta = x_1 x_2 + y_1 y_2 + z_1 z_2$$

$$\dot{\vec{r}} = \frac{d\vec{r}}{dt} = \frac{dx}{dt}\vec{i} + \frac{dy}{dt}\vec{j} + \frac{dz}{dt}\vec{k}$$

Mechanics

$$\text{momentum:} \qquad \vec{p} = m \vec{v}$$

$$\text{equation of motion:} \qquad \vec{R} = m \vec{a}$$

$$\text{friction:} \qquad F \leq \mu N$$

Reproduced with permission of the Victorian Curriculum and Assessment Authority, Victoria, Australia.

*This formula sheet has been copied in 2009 from the VCAA website www.vcaa.vic.edu.au
The VCAA publish an exam issue supplement to the VCAA bulletin.*