


2012 Specialist Maths Trial Exam 2 Solutions
Free download from www.itute.com © Copyright itute.com

Section 1

1	2	3	4	5	6	7	8	9	10	11
A	C	E	D	E	E	A	E	C	A	B

12	13	14	15	16	17	18	19	20	21	22
D	B	B	B	B	A	C	C	B	D	E


Q1 A possible complex number is $z = 1 - i$, $|z| = \sqrt{2}$ A



Q2 $z = -icis(2)$ is the clockwise rotation of $cis(2)$ about the origin O by $\frac{\pi}{2}$. $\therefore Arg(z) = 2 - \frac{\pi}{2}$ C

Q3 Let $z = x + yi$.
 $az^2 + b|z|^2 - c = 0$, $a(x^2 - y^2 + 2xyi) + b(x^2 + y^2) - c = 0$
 $\therefore [(a+b)x^2 + (b-a)y^2 - c] + 2axyi = 0$
 $\therefore (a+b)x^2 + (b-a)y^2 - c = 0$ and $xy = 0$
 $\therefore$ either $x = 0$ and $y = \pm \sqrt{\frac{c}{b-a}}$ if $b > a$
 or $y = 0$ and $x = \pm \sqrt{\frac{c}{b+a}}$ E

Q4 $0 < Arg[(1+i)z] < \frac{\pi}{2}$, $0 < Arg(1+i) + Arg(z) < \frac{\pi}{2}$,
 $0 < \frac{\pi}{4} + Arg(z) < \frac{\pi}{2}$, $-\frac{\pi}{4} < Arg(z) < \frac{\pi}{4}$



$$\frac{(x-h)^2}{9} + (y-k)^2 = 1 \quad \text{or} \quad (x-h)^2 + \frac{(y-k)^2}{9} = 1$$
 E

Q6 $y = \frac{1}{a+bx+ax^2}$ has a vertical and a horizontal asymptotes when $a+bx+ax^2$ is a perfect square, i.e. $\Delta = b^2 - 4a^2 = 0$
 $\therefore \frac{a}{b} = \pm \frac{1}{2}$ E

Q7 Asymptotes: $y = 2 \pm \left(\frac{1}{2}x + 1\right)$, $y - 2 = \pm \frac{1}{2}(x + 2)$
 $\therefore$ the two asymptotes intersect at $(-2, 2)$ which is the centre of the hyperbola, also $\frac{b}{a} = \frac{1}{2}$.


A possible equation of the hyperbola is $\frac{(x+2)^2}{2^2} - \frac{(y-2)^2}{1^2} = 1$,
 i.e. $(x+2)^2 - 4(y-2)^2 - 4 = 0$ A

Q8 $\sec(a) + \operatorname{cosec}(b) = 0$, $\sec(a) = -\operatorname{cosec}(b)$
 $\therefore \cos(a) = -\sin(b)$, $\cos(a) = -\cos\left(\frac{\pi}{2} - b\right)$
 Since $\pi < a < \frac{3\pi}{2}$ and $0 < b < \frac{\pi}{2}$, $\therefore a = \pi + \left(\frac{\pi}{2} - b\right) = \frac{3\pi}{2} - b$
 $\therefore a + b = \frac{3\pi}{2}$ E

Q9 The domain of the inverse of f is the range of f .
 As $x \rightarrow 1^+$, $y \rightarrow \sec \pi + a\pi = a\pi - 1$
 At $x = \frac{5}{3}$, $y = \sec \frac{4\pi}{3} + a\pi = a\pi - 2$
 $\therefore$ the domain of the inverse of f is $[a\pi - 2, a\pi - 1)$. C

Q10 $\tan^{-1}(x-a+b) - \tan^{-1}(x-a) = \pi$
 $\tan^{-1}(x-a+b)$ and $\tan^{-1}(x-a)$ have the same range
 $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, $\therefore \tan^{-1}(x-a+b) - \tan^{-1}(x-a) \neq \pi$ for $x \in R$ A

Q11 Since $\tilde{a} - \tilde{b} + \tilde{c} = \tilde{0}$, $\therefore$ unit vectors $\tilde{a}$, $\tilde{b}$ and $\tilde{c}$ form an equilateral triangle as shown below:



Q12 Let $\tilde{j} + c\tilde{k} = m(\tilde{k} + a\tilde{i}) + n(\tilde{i} + b\tilde{j})$ where m and n are non-zero real numbers.
 $\tilde{j} + c\tilde{k} = (ma+n)\tilde{i} + nb\tilde{j} + m\tilde{k}$
 $\therefore ma+n=0$, $nb=1$ and $m=c$
 $\therefore ca+n=0$, $n=-ca$
 $\therefore abc = -1$ D

Q13 $\tilde{p} = \pm(\tilde{i} + \tilde{j} + \tilde{k})$ are vectors which make equal angle θ with each of the orthogonal unit vectors $\tilde{i}$, $\tilde{j}$ and $\tilde{k}$.
 $\cos \theta = \pm \frac{1}{\sqrt{1+1+1}}$, $\therefore \theta \approx 55^\circ$ and 125° B

Q14 Given $f(0)=1.2$, by CAS $f(0.2)=\int_0^{0.2} \frac{1}{\cos^4 x} dx + 1.2 \approx 1.4$,

$f(0.4)=\int_0^{0.4} \frac{1}{\cos^4 x} dx + 1.2 \approx 1.6$, $f(0.6)=\int_0^{0.6} \frac{1}{\cos^4 x} dx + 1.2 \approx 2$

$f(0.8)=\int_0^{0.8} \frac{1}{\cos^4 x} dx + 1.2 \approx 2.6$, $f(1)=\int_0^1 \frac{1}{\cos^4 x} dx + 1.2 \approx 4$


Average value $\approx \frac{1.2+1.4+1.6+2+2.6+4}{6} \approx 2$

B

Q15 At a point of inflection the gradient of the curve is a local maximum/minimum.

$y = 3x^5 + 5x^4 - 10x^3 - 30x^2 + 5x - 10$

$\frac{dy}{dx} = 15x^4 + 20x^3 - 30x^2 - 60x + 5$



$\frac{dy}{dx}$ has a local minimum at $x = 1$.

B

Q16 When $y = 2$ and $0 \leq x \leq \frac{3}{2}$, $\tan^2 x = 4$, $x \approx 1.10715$

By CAS, $\int_0^{1.10715} \tan^2 x dx \approx 0.8929$

Area of the required region $\approx 4 \times 1.10715 - 0.8929 \approx 3.5$

B

Q17 The quadratic equation has exactly one solution, $\therefore \Delta = 0$

$k^2 - 4(1)(1) = 0$, $\therefore k = 2$


$\left(\frac{dy}{dx} + 1\right)^2 = 0$, $\therefore \frac{dy}{dx} = -1$, $y = -x + c$

$(1,1)$, $\therefore c = 2$, $\therefore x + y = 2$

Q18 $a = +2$, $u = -10$, $s = +16 - +5 = +11$, $v = ?$

$v^2 = u^2 + 2as$, $v = +12$

$v = 2t - 10$



Total distance $= 25 + 36 = 61$ m

C

Q19 Given $v = 2 - e^x$, when $a = v \frac{dv}{dx} = (2 - e^x)(-e^x) = 0$,

$e^x = 2$, $x = \log_e 2$ and $v = 0$. $\therefore$ the particle is moving in the same direction from $x = 0$ to $x = \log_e 2$.

Distance $= \log_e 2 \approx 0.7$ m

C

Q20 $a = +9.8 \sin 30^\circ = +4.9$, $s = +1.3$, $u = 0$, $v = ?$

$v^2 = u^2 + 2as$, $v \approx +3.57$

Momentum $= mv \approx 1.3 \times 3.57 \approx +4.6$ kg m s⁻¹

B

Q21 The particle exerts a downward force on the inclined plane equal in magnitude to the weight of the particle, $\therefore$ the upward reaction force of the inclined plane on the particle is also equal in magnitude to the weight of the particle according to Newton's third law, $mg = 2.04 \times 9.8 \approx 20$ N upward

D

Q22 Acceleration of the crate = acceleration of the chain
Consider the forces on the chain:

1000 N $\leftarrow$ 10 kg $\rightarrow$ 1010 N

$a = \frac{R}{m} = \frac{1010 - 1000}{10} = 1$ m s⁻²

E

Section 2

Q1a $x^2 + 4y^2 = 4 \xrightarrow{(1)} x^2 + 4\left(y - \frac{1}{2}\right)^2 = 4$

$\xrightarrow{(2)} x^2 + 4\left(\frac{1}{2}y - \frac{1}{2}\right)^2 = 4$, and in simplified form

$x^2 + (y - 1)^2 = 4$.

Q1b Let $y = 0$, $x^2 = 3$, $x = \pm\sqrt{3}$

The x -intercepts are $(-\sqrt{3}, 0)$ and $(\sqrt{3}, 0)$.


A

Q1ci $x^2 + (y - 1)^2 = 4$, $(y - 1)^2 = 4 - x^2$, $y - 1 = \pm\sqrt{4 - x^2}$

$\therefore g(x) = 1 - \sqrt{4 - x^2}$,

$\therefore f(x) = |g(x)| = \sqrt{4 - x^2} - 1$ for $-\sqrt{3} < x < \sqrt{3}$

Q1cii



Q1d Volume V_1 of solid by formed by rotating $f(x)$:

$$y = \sqrt{4-x^2} - 1, \sqrt{4-x^2} = y+1, x^2 = 4-(y+1)^2$$


$$V_1 = \int_0^1 \pi x^2 dy = \pi \int_0^1 (4-(y+1)^2) dy = \frac{5\pi}{3}$$

$$\text{Volume of the large sphere } V_2 = \frac{4}{3} \pi (2^3) = \frac{32\pi}{3}$$

$\therefore$ Volume of revolution of the required region

$$= \frac{32\pi}{3} - 2 \times \frac{5\pi}{3} = \frac{22\pi}{3}$$

Q2a



When $x = 2$, $y \approx 1.7$

Q2b

$$x = 0.5 \quad y = 0.5$$

$$\frac{dy}{dx} \approx 1.1765$$

$$x = 1.0 \quad y = 0.5 + 0.5 \times 1.1765 \approx 1.0882$$

$$\frac{dy}{dx} = 1$$

$$x = 1.5 \quad y = 1.0882 + 0.5 \times 1 \approx 1.5882$$

$$\frac{dy}{dx} \approx 0.5361$$

$$x = 2.0 \quad y \approx 1.5882 + 0.5 \times 0.5361 \approx 1.9$$

$$\text{Q2c } y = \int_{0.5}^2 \frac{1+x^2}{1+x^4} dx + 0.5 \approx 1.7 \text{ by CAS}$$

$$\text{Q2di } \frac{\frac{1+x^2}{x^2}}{\frac{1+x^4}{x^2}} = \frac{\frac{1}{x^2} + 1}{\frac{1}{x^2} + x^2}$$

$$\text{Q2dii } x^2 + \frac{1}{x^2} = \left(x + \frac{b}{x}\right)^2 + c = x^2 + 2b + \frac{b^2}{x^2} + c,$$

$$\therefore b^2 = 1, 2b + c = 0 \text{ and } c > 0$$

$$\therefore b = -1 \text{ and } c = 2$$

$$\therefore x^2 + \frac{1}{x^2} = \left(x - \frac{1}{x}\right)^2 + 2$$

$$\text{Q2diii } y = \int \frac{1+x^2}{1+x^4} dx = \int \frac{\frac{1}{x^2} + 1}{\left(x - \frac{1}{x}\right)^2 + 2} dx$$

$$= \int \frac{\frac{du}{dx}}{u^2 + 2} dx = \int \frac{1}{2+u^2} du$$

$$= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{u}{\sqrt{2}}\right) + c$$

$$= \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{1}{\sqrt{2}}\left(x - \frac{1}{x}\right)\right) + c$$

$$\text{Let } u = x - \frac{1}{x}$$

$$\frac{du}{dx} = 1 + \frac{1}{x^2}$$

Q2div When $x = 0.5$, $y = 0.5$

$$0.5 = \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{1}{\sqrt{2}}\left(0.5 - \frac{1}{0.5}\right)\right) + c, c = 1.0762$$

$$\therefore y = \frac{1}{\sqrt{2}} \tan^{-1}\left(\frac{1}{\sqrt{2}}\left(x - \frac{1}{x}\right)\right) + 1.0762$$

When $x = 2$, $y \approx 1.7$

$$\text{Q3a } \overrightarrow{P_1P_2} = \tilde{b} - \tilde{a}, |\overrightarrow{P_1P_2}|^2 = \overrightarrow{P_1P_2} \cdot \overrightarrow{P_1P_2} = (\tilde{b} - \tilde{a}) \cdot (\tilde{b} - \tilde{a})$$

$$= \tilde{a} \cdot \tilde{a} + \tilde{b} \cdot \tilde{b} - 2\tilde{a} \cdot \tilde{b} = |\tilde{a}|^2 + |\tilde{b}|^2 - 2|\tilde{a}||\tilde{b}|\cos\alpha$$

Q3b Since the length of arc P_1P_2 equals the length of arc P_2P_3 ,

$$\therefore \angle P_2OP_3 = \angle P_1OP_2 = \alpha$$

$$|\overrightarrow{P_2P_3}|^2 = |\tilde{b}|^2 + |\tilde{c}|^2 - 2|\tilde{b}||\tilde{c}|\cos\alpha$$

Q3c Since the length of arc P_1P_2 equals the length of arc P_2P_3 ,

$$\therefore |\overrightarrow{P_1P_2}|^2 = |\overrightarrow{P_2P_3}|^2$$

$$\therefore |\tilde{a}|^2 + |\tilde{b}|^2 - 2|\tilde{a}||\tilde{b}|\cos\alpha = |\tilde{b}|^2 + |\tilde{c}|^2 - 2|\tilde{b}||\tilde{c}|\cos\alpha$$

$$\therefore |\tilde{a}|^2 - |\tilde{c}|^2 = 2|\tilde{b}|(|\tilde{a}| - |\tilde{c}|)\cos\alpha$$

$$\therefore \cos\alpha = \frac{|\tilde{a}|^2 - |\tilde{c}|^2}{2|\tilde{b}|(|\tilde{a}| - |\tilde{c}|)} = \frac{|\tilde{a}| + |\tilde{c}|}{2|\tilde{b}|}$$

Q3d Since OP_1 is a diameter, $\therefore \angle OP_2P_1$ and $\angle OP_3P_1$ are right angles.

$\therefore \tilde{b}$ is the vector resolute of $\tilde{a}$ in the direction of $\overrightarrow{OP_2}$ and $\tilde{c}$

is the vector resolute of $\tilde{a}$ in the direction of $\overrightarrow{OP_3}$.

Q3e Given the length of $\overrightarrow{OP_1} = |\tilde{a}| = 1$, $\therefore |\tilde{b}| = 1 \cos\alpha = \cos\alpha$

and $|\tilde{c}| = 1 \cos 2\alpha = \cos 2\alpha$.

$$\text{Q3f } \cos\alpha = \frac{|\tilde{a}| + |\tilde{c}|}{2|\tilde{b}|}, \cos\alpha = \frac{1 + \cos 2\alpha}{2 \cos\alpha}, \therefore 1 + \cos 2\alpha = 2 \cos^2\alpha$$

$$\text{Q3g } \cos\alpha = \frac{|\tilde{b}| + |\tilde{d}|}{2|\tilde{c}|} \text{ and } |\tilde{d}| = \cos 3\alpha \text{ by similar methods,}$$

$$\therefore \cos\alpha = \frac{\cos\alpha + \cos 3\alpha}{2 \cos 2\alpha},$$

$$\therefore \cos\alpha + \cos 3\alpha = 2 \cos\alpha \cos 2\alpha$$

4