
Year 2013

VCE

Specialist Mathematics

Trial Examination 1

Solutions



KILBAHA MULTIMEDIA PUBLISHING
PO BOX 2227
KEW VIC 3101
AUSTRALIA

TEL: (03) 9018 5376
FAX: (03) 9817 4334
kilbaha@gmail.com
<http://kilbaha.com.au>

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Multimedia Publishing.
- The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
- For authorised copying within Australia please check that your institution has a licence from **Copyright Agency Limited**. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.

Reproduction and communication for educational purposes The Australian Copyright Act 1968 (the Act) allows a maximum of one chapter or 10% of the pages of this work, to be reproduced and/or communicated by any educational institution for its educational purposes provided that educational institution (or the body that administers it) has given a remuneration notice to Copyright Agency Limited (CAL) under the Act.

For details of the CAL licence for educational institutions contact
CAL, Level 15, 233 Castlereagh Street, Sydney, NSW, 2000
Tel: (02) 9394 7600
Fax: (02) 9394 7601
Email: info@copyright.com.au
Web: <http://www.copyright.com.au>

- Teachers and students are reminded that for the purposes of school requirements and external assessments, students must submit work that is clearly their own.

Question 1

$$\int \frac{3x-5}{\sqrt{25-9x^2}} dx \quad \text{separate out into two integrals}$$

$$= 3 \int \frac{x}{\sqrt{25-9x^2}} dx - 5 \int \frac{1}{\sqrt{25-9x^2}} dx \quad \text{M1}$$

$$\text{let } u = 25-9x^2 \quad \text{let } v = 3x$$

$$\frac{du}{dx} = -18x \quad \frac{dv}{dx} = 3 \quad \text{M1}$$

$$= -\frac{3}{18} \int u^{-\frac{1}{2}} du - \frac{5}{3} \int \frac{1}{\sqrt{25-v^2}} dv \quad \text{A1}$$

$$= -\frac{1}{6} \left(2u^{\frac{1}{2}} \right) + \frac{5}{3} \cos^{-1} \left(\frac{v}{5} \right) + c$$

$$= -\frac{1}{3} \sqrt{25-9x^2} + \frac{5}{3} \cos^{-1} \left(\frac{3x}{5} \right) + c \quad \text{for } |x| < \frac{5}{3} \quad b = \frac{5}{3} \quad \text{A1}$$

$$\text{alternatively } = -\frac{1}{3} \sqrt{25-9x^2} - \frac{5}{3} \sin^{-1} \left(\frac{3x}{5} \right) + c \quad \text{for } |x| < \frac{5}{3} \quad b = \frac{5}{3}$$

Question 2

$$y = Ax^2 e^{-3x} \quad \text{differentiating using the product rule} \quad \text{M1}$$

$$\frac{dy}{dx} = A(2xe^{-3x} - 3x^2 e^{-3x}) = Ae^{-3x}(2x - 3x^2) \quad \text{differentiating using the product rule again}$$

$$\frac{d^2 y}{dx^2} = -3Ae^{-3x}(2x - 3x^2) + Ae^{-3x}(2 - 6x) = Ae^{-3x}[-3(2x - 3x^2) + (2 - 6x)]$$

$$\frac{d^2 y}{dx^2} = Ae^{-3x}(9x^2 - 12x + 2) \quad \text{A1}$$

$$\text{substituting into } \frac{d^2 y}{dx^2} + 6 \frac{dy}{dx} + 9y = 8e^{-3x} \quad \text{M1}$$


$$Ae^{-3x}[(9x^2 - 12x + 2) + 6(2x - 3x^2) + 9x^2] = 8e^{-3x}$$

$$Ae^{-3x}[9x^2 - 12x + 2 + 12x - 18x^2 + 9x^2] = 8e^{-3x}$$

$$2Ae^{-3x} = 8e^{-3x}$$

$$2A = 8$$

$$A = 4 \quad \text{A1}$$

Question 3

all the forces are newtons

resolving parallel to the plane (1) $Fg \cos(30^\circ) - \frac{\sqrt{3}}{3}N = 5a$

resolving perpendicular to the plane (2) $Fg \sin(30^\circ) + N - 5g = 0$

(2) $\Rightarrow N = 5g - Fg \sin(30^\circ) = 5g - \frac{Fg}{2} = \frac{g}{2}(10 - F)$ A1

so if $F > 10 \Rightarrow N < 0$, so that the box leaves the table. A1

(1) $\Rightarrow Fg \frac{\sqrt{3}}{3} - \frac{\sqrt{3}}{3} \left(\frac{g}{2}(10 - F) \right) = 5a$

$\frac{g\sqrt{3}}{3} \left(F - \frac{1}{2}(10 - F) \right) = 5a$ M1

$\frac{g\sqrt{3}}{6}(3F - 10) = 5a$

$\frac{10}{3} < F < 10 \Rightarrow a > 0$ the box moves with constant acceleration. A1

Question 4

$z = 2\sqrt{2}\text{cis}\left(\frac{3\pi}{4}\right) = 2\sqrt{2}\left(\cos\left(\frac{3\pi}{4}\right) + i\sin\left(\frac{3\pi}{4}\right)\right) = 2\sqrt{2}\left(-\frac{1}{\sqrt{2}} + i\frac{1}{\sqrt{2}}\right) = -2 + 2i$ A1

The conjugate $\bar{z} = -2 - 2i$ is also a root, by the conjugate root theorem,

the sum of the roots $z + \bar{z} = -4$, the product of the roots $z\bar{z} = 4 - 4i^2 = 8$

since b is real $(z^2 + 4z + 8)$ is a factor M1

$P(z) = z^3 + z^2 + bz - 24 = (z^2 + 4z + 8)(z - 3)$

expanding $z: b = 8 - 12 = -4$

$b = -4$, and all the roots are $z = -2 \pm 2i$, $z = 3$ A1

Question 5

$$\cos(\theta) = \frac{2}{3} \text{ draw a triangle}$$

by Pythagoras

$$x = \sqrt{3^2 - 2^2} = \sqrt{5}$$

$$\sin(\theta) = \frac{\sqrt{5}}{3}$$

$$z = 3 \operatorname{cis}(\theta)$$

$$z^2 = 9 \operatorname{cis}(2\theta) = 9(\cos(2\theta) + i \sin(2\theta))$$



$$z^2 = 9(\cos^2(\theta) - \sin^2(\theta)) + 18 \sin(\theta) \cos(\theta) i$$

$$z^2 = 9\left(\frac{4}{9} - \frac{5}{9}\right) + 18 \times \frac{\sqrt{5}}{3} \times \frac{2}{3} i$$

$$z^2 = -1 + 4\sqrt{5}i$$

M1

A1

**Question 6**

$OABC$ is a parallelogram, $\overrightarrow{OA} = \overrightarrow{CB}$ and $\overrightarrow{OC} = \overrightarrow{AB}$

since P is the midpoint of OA , $\overrightarrow{OP} = \overrightarrow{PA} = \frac{1}{2}\overrightarrow{OA}$, given that $\overrightarrow{PD} = \frac{1}{3}\overrightarrow{PC}$

A1

consider $\overrightarrow{OD} = \overrightarrow{OP} + \overrightarrow{PD}$

$$= \frac{1}{2}\overrightarrow{OA} + \frac{1}{3}\overrightarrow{PC} = \frac{1}{2}\overrightarrow{OA} + \frac{1}{3}(\overrightarrow{PO} + \overrightarrow{OC})$$

M1

$$= \frac{1}{2}\overrightarrow{OA} + \frac{1}{3}\left(\overrightarrow{OC} - \frac{1}{2}\overrightarrow{OA}\right)$$

$$= \left(\frac{1}{2} - \frac{1}{6}\right)\overrightarrow{OA} + \frac{1}{3}\overrightarrow{OC} \quad \text{since } \overrightarrow{OC} = \overrightarrow{AB}$$

M1

$$= \frac{1}{3}(\overrightarrow{OA} + \overrightarrow{AB})$$

So that $\overrightarrow{OD} = \frac{1}{3}\overrightarrow{OB}$ this shows that O , D and B are collinear.


A1

Question 7

a. $\cos(3A) = \cos(2A + A)$
 $= \cos(2A)\cos(A) - \sin(2A)\sin(A)$ M1
 $= (2\cos^2(A) - 1)\cos(A) - 2\sin^2(A)\cos(A)$
 $= (2\cos^2(A) - 1)\cos(A) - 2(1 - \cos^2(A))\cos(A)$
 $= 2\cos^3(A) - \cos(A) - 2\cos(A) + 2\cos^3(A)$
 $\cos(3A) = 4\cos^3(A) - 3\cos(A)$

b.i. $\underline{r}(t) = 2\cos(t)\underline{i} + 4\cos(3t)\underline{j}$ for $t \geq 0$
 $x = 2\cos(t)$ $y = 4\cos(3t)$
 $\cos(t) = \frac{x}{2}$, $\cos(3t) = \frac{y}{4}$
 $\cos(3t) = 4\cos^3(t) - 3\cos(t)$
 $\frac{y}{4} = 4\left(\frac{x}{2}\right)^3 - 3\left(\frac{x}{2}\right)$
 $y = 2x^3 - 6x$ A1
 but since $t \geq 0$, $x = 2\cos(t) \Rightarrow x \in [-2, 2]$
 $y = 4\cos(3t) \Rightarrow y \in [-4, 4]$ A1

ii. the particle moves on part of a cubic,
 $y = 2x^3 - 6x = 2x(x^2 - 3) = 2x(x + \sqrt{3})(x - \sqrt{3})$
 crosses the x-axis when $y = 0$ at $(\sqrt{3}, 0)$, $(0, 0)$, $(-\sqrt{3}, 0)$
 $\frac{dy}{dx} = 6x^2 - 6 = 6(x^2 - 1) = 6(x + 1)(x - 1)$
 when $\frac{dy}{dx} = 0$ there are turning points at $(-1, 4)$ and $(1, -4)$ A1
 correct graph, on restricted domain, with key features. G1

**Question 8**

$$v = \frac{dx}{dt} = \frac{5}{100 - 9t^2}$$

distance travelled in two seconds $D = \int_0^2 \frac{5}{100 - 9t^2} dt$ A1

by partial fractions

$$\frac{5}{100 - 9t^2} = \frac{A}{10 - 3t} + \frac{B}{10 + 3t} = \frac{A(10 + 3t) + B(10 - 3t)}{(10 - 3t)(10 + 3t)} = \frac{10(A + B) + 3t(A - B)}{100 - 9t^2}$$
 M1

$$(1) A + B = \frac{1}{2} \quad (2) A - B = 0 \quad \text{solving} \quad A = B = \frac{1}{4}$$

$$D = \frac{1}{4} \int_0^2 \left(\frac{1}{10 - 3t} + \frac{1}{10 + 3t} \right) dt$$

$$D = \frac{1}{4} \left[-\frac{1}{3} \log_e(10 - 3t) + \frac{1}{3} \log_e(10 + 3t) \right]_0^2$$
 M1

$$D = \frac{1}{12} \left[\log_e \left(\frac{10 + 3t}{10 - 3t} \right) \right]_0^2 = \frac{1}{12} \left(\log_e \left(\frac{16}{4} \right) - \log_e(1) \right)$$

$$D = \frac{1}{12} \log_e(4) \text{ metres}$$
 A1

Question 9

a. $x^2 + 4x - 4y^2 = 0$

$$x^2 + 4x + 4 - 4y^2 = 4 \quad \text{completing the square}$$

$$(x+2)^2 - 4y^2 = 4$$

$$\frac{(x+2)^2}{4} - y^2 = 1, \text{ hyperbola centre } (-2, 0)$$

A1

crosses the x-axis, when


$$y = 0, (x+2)^2 = 4, x+2 = \pm 2, x = 0, -4 \quad (0, 0), (-4, 0),$$

$$\text{asymptotes } y = \pm \frac{1}{2}(x+2) \quad y = \frac{x}{2} + 1 \text{ and } y = -\frac{1}{2}x - 1$$

A1

correct graph, with asymptotes

A1



b. $x^2 + 4x - 4y^2 = 0$ using implicit differentiation

$$2x + 4 - 8y \frac{dy}{dx} = 0$$

$$8y \frac{dy}{dx} = 2x + 4 = 2(x+2)$$

$$\frac{dy}{dx} = \frac{x+2}{4y}$$

A1

Question 10

i. $y = \tan(2x)\sec(2x) = \frac{\sin(2x)}{\cos^2(2x)}$

asymptotes when denominator is zero, when $\cos(2x) = 0 \Rightarrow 2x = (2n+1)\frac{\pi}{2}$

$$x = (2n+1)\frac{\pi}{4} \quad \text{where } n \in \mathbb{Z} \quad \text{A1}$$

ii. area $A = \int_0^{\frac{\pi}{6}} \tan(2x)\sec(2x) dx = \int_0^{\frac{\pi}{6}} \frac{\sin(2x)}{\cos^2(2x)} dx$

let $u = \cos(2x) \quad \frac{du}{dx} = -2\sin(2x)$ M1

terminals when $x = \frac{\pi}{6} \quad u = \cos\left(\frac{\pi}{3}\right) = \frac{1}{2}$ and when $x = 0 \quad u = \cos(0) = 1$

$$A = -\frac{1}{2} \int_1^{\frac{1}{2}} \frac{1}{u^2} du = -\frac{1}{2} \int_1^{\frac{1}{2}} u^{-2} du \quad \text{M1}$$

$$A = \frac{1}{2} \left[\frac{1}{u} \right]_1^{\frac{1}{2}} = \frac{1}{2} (2-1)$$

$$A = \frac{1}{2} \quad \text{A1}$$

iii. volume $V = \pi \int_a^b y^2 dx$

$$V = \pi \int_0^{\frac{\pi}{6}} \tan^2(2x)\sec^2(2x) dx$$

let $u = \tan(2x) \quad \frac{du}{dx} = 2\sec^2(2x)$ M1

terminals when $x = \frac{\pi}{6} \quad u = \tan\left(\frac{\pi}{3}\right) = \sqrt{3}$ and when $x = 0 \quad u = \tan(0) = 0$

$$V = \frac{\pi}{2} \int_0^{\sqrt{3}} u^2 du$$

$$= \frac{\pi}{2} \left[\frac{1}{3} u^3 \right]_0^{\sqrt{3}} = \frac{\pi}{6} \left((\sqrt{3})^3 - 0 \right)$$

$$V = \frac{\pi\sqrt{3}}{2} \quad \text{A1}$$

END OF SUGGESTED SOLUTIONS