



Online & home tutors Registered business name: itute ABN: 96 297 924 083

Specialist Mathematics

2014

Trial Examination 1

Instructions

Answer **all** questions. Do **not** use calculators.

A decimal approximation will not be accepted if an **exact** answer is required to a question.

In questions where more than one mark is available, appropriate working or explanation **must** be shown.

Unless otherwise indicated, the diagrams in this exam are **not** drawn to scale.

Question 1

Consider function f with the rule $f(x) = \left(\frac{1}{\sqrt{x}} - \sqrt{x}\right)^2 + 2$.

a. Simplify the rule of f .


1 mark

b. Find the range of f .

1 mark

c. Sketch the graph of f . Show and label the turning point(s) and the asymptote(s) of f .

3 marks




Question 2

Consider $g : R \rightarrow R$, $g(x) = \tan^{-1}(3x) + \tan^{-1}(2x) - \frac{\pi}{4}$.

a. Find the exact value(s) of x where $g(x) = 0$. 2 marks

b. Use the method of addition of ordinates to sketch the graph of g . Show and label the axis-intercept(s) and the asymptote(s) of g . 3 marks



Question 3

a. Solve $\sqrt{3}z - \sqrt{2}i = \sqrt{2}iz + \sqrt{3}$ for z . Express your answer in $x + yi$ form.

2 marks

b. $P(z)$ is a cubic polynomial in z with real coefficients.

Given $P(z) = (z - i)Q(z) + 1$ and $P(z) = (2z - 1)T(z) + 1$ where $Q(z)$ and $T(z)$ are polynomials in z , solve $P(z) = 0$ for z .

3 marks

Question 4


Consider $|z| - |3 - z| = 1$ where $z = x + yi$ and $x, y \in \mathbb{R}$.

a. Express $|z| - |3 - z| = 1$ in the form $\frac{(x-h)^2}{a} - \frac{(y-k)^2}{b} = 1$.

2 marks

b. Hence or otherwise sketch the graph of $|z| - |3 - z| \leq 1$.

2 marks



Question 5


Show that $\tilde{p} = \tilde{i} - \tilde{j}$, $\tilde{q} = 2\tilde{i} + \tilde{j}$, $\tilde{r} = \tilde{i} + 2\tilde{j}$ and $\tilde{s} = 3\tilde{i} - 2\tilde{j}$ are linearly *dependent*.

2 marks

Question 6

$ABCDEFGH$ is a cuboid. Use vector method to find the shortest distance in surd form from vertex G to line AH .

3 marks



Question 7

The solution curve to the differential equation $\frac{dy}{dx} + \frac{y}{x} = 0$ passes through $(1, 2)$.

- a. Use Euler's method (first order approximation) to estimate the value of y at $x = 2.5$.
Choose 0.5 as the step size.

2 marks

- b. Show that $xy = 2$ is the equation of the solution curve.

1 mark


- c. If both x and y are functions of λ and $\frac{dy}{d\lambda} = -1$, find $\frac{dx}{d\lambda}$ at $x = 1$.

1 mark

Question 8

Consider $\frac{dy}{dx} = f(x)$ with $y = 5$ when $x = 1$. The graph of $\frac{dy}{dx} = f(x)$ for $x \in [1, 6]$ is shown below.

The areas of the regions (shaded) bounded by the curve, the x -axis, $x = 1$ and $x = 6$ are indicated in the graph.



- a. Evaluate $\int_1^6 f(x) dx$.


1 mark

- b. Find y when $x = 6$.

1 mark

Question 9


A sign is erected in the street which runs in the north-south direction. The displacement (m) of a cyclist from the sign as a function of time (s) is shown in the following graph. A positive displacement indicates a displacement to the north.



- a. State the direction of motion of the cyclist at time $t = 120$ s. 1 mark

- b. Calculate the average speed (in m s^{-1}) of the pedestrian between $t = 0$ and $t = 160$ s. 1 mark

The velocity-time graph of a car is shown below. The car starts at 500 metres north of the street sign.




- c. State the direction of motion of the car at $t = 120$ s. 1 mark

- d. Calculate the average speed (in km h^{-1}) of the car between $t = 0$ and $t = 160$ s. 2 marks

- e. How many times does the car pass the cyclist between $t = 0$ and $t = 160$ s? 1 mark

Question 10

A 1.9 kg parcel is attached to a frictionless 0.1 kg pulley. A cable of negligible mass is fastened to a garage ceiling and wall. The pulley is allowed to run along the cable until it comes to a stop as shown in the drawing below. Take $g = 10 \text{ N kg}^{-1}$. Assume the pulley is a point mass.



- a. Calculate the exact value in newtons of the tension in the cable.

2 marks

Now a horizontal force of F newtons is used to pull the pulley to the left until the section of the cable on the right is horizontal.

- b. Calculate the exact value of F .

2 marks

End of Exam 1