

**‘2016 Examination Package’ -
Trial Examination 4 of 5**

STUDENT NUMBER

Figures

Words

Letter

--

SPECIALIST MATHEMATICS

Units 3 & 4 – Written examination 2

(TSSM’s 2014 trial exam updated for the current study design)

Reading time: 15 minutes

Writing time: 2 hours

QUESTION AND ANSWER BOOK

Structure of book

Section	Number of questions	Number of questions to be answered	Number of marks
1	22	22	22
2	5	5	58
			Total 80

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers, a protractor, set-squares, aids for curve sketching, one bound reference, one approved graphics calculator or approved CAS calculator and a scientific calculator.
- Students are NOT permitted to bring into the examination room: blank sheets of paper and/or white out liquid/tape.

Materials supplied

- Question and answer book of 23 pages.(including a multiple choice answer sheet)

Instructions

- Print your name in the space provided on the top of this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other electronic devices into the examination room.

SECTION 1

Instructions for Section 1

Answer **all** questions on the answer sheet provided for multiple-choice questions.

Choose the response that is **correct** for the question.

A correct answer scores 1, an incorrect answer scores 0.

Marks are **not** deducted for incorrect answers.

If more than 1 answer is completed for any question, no mark will be given.

Take the **acceleration due to gravity**, to have magnitude $g \text{ m/s}^2$, where $g = 9.8$.

Question 1

The graph of $y = \frac{-x^2 + 1}{2x}$ has

- A. no straight line asymptotes.
- B. $y = 2x$ as its only straight line asymptote.
- C. $x = 0$ as its only straight line asymptote.
- D. $y = 0$ and $y = -\frac{1}{2}x$ as its only straight line asymptotes.
- E. $x = 0$ and $y = -\frac{1}{2}x$ as its only straight line asymptotes.

Question 2

An antiderivative of $\frac{2}{\sqrt{4-x^2}}$ could be:

- A. $\cos^{-1}\left(\frac{x}{2}\right)$
- B. $2\cos^{-1}\left(\frac{x}{2}\right)$
- C. $\sin^{-1}\left(\frac{x}{2}\right)$
- D. $1 - 2\cos^{-1}\left(\frac{x}{2}\right)$
- E. $\frac{1}{2}\sin^{-1}\left(\frac{x}{2}\right)$

Question 3

If $z = 3 + 2i$ then $\frac{\bar{z}}{z}$ is equal to:

- A. 13
- B. $\frac{5-12i}{5}$
- C. $\frac{5-12i}{13}$
- D. $\frac{13-12i}{5}$
- E. $\frac{13-12i}{13}$

Question 4

The implied domain and range of $\sin^{-1}\left(\frac{x-1}{2}\right)$ respectively are:

- A. $\left[-\frac{1}{2}, \frac{1}{2}\right]$ and $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
- B. $[-2, 2]$ and $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$
- C. $[-2, 2]$ and $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
- D. $\left[\frac{1}{2}, \frac{3}{2}\right]$ and $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$
- E. $[-1, 3]$ and $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

SECTION 1 - continued
TURN OVER

Question 5

A unit vector perpendicular to the vector $\vec{i} + 2\vec{j} - 3\vec{k}$ is:

- A. $\vec{i} + \vec{j} + \vec{k}$
- B. $-\vec{i} - 2\vec{j} + 3\vec{k}$
- C. $\frac{1}{\sqrt{14}}(\vec{i} + 2\vec{j} - 3\vec{k})$
- D. $\frac{1}{\sqrt{6}}(\vec{i} - 2\vec{j} - \vec{k})$
- E. $\frac{1}{4}(-\vec{i} + 3\vec{k})$

Question 6

Using a suitable substitution, $\int_0^1 x^3 \sqrt{1-x^2} dx$ can be expressed as:

- A. $\int_0^1 \left(u^{\frac{1}{2}} - u^{\frac{3}{2}} \right) du$
- B. $\frac{1}{2} \int_0^1 \left(u^{\frac{1}{2}} - u^{\frac{3}{2}} \right) du$
- C. $\int_0^1 \left(u^{\frac{3}{2}} - u^{\frac{1}{2}} \right) du$
- D. $2 \int_0^1 \left(u^{\frac{3}{2}} - u^{\frac{1}{2}} \right) du$
- E. $2 \int_0^1 \left(u^{\frac{1}{2}} - u^{\frac{3}{2}} \right) du$

SECTION 1 - continued

Question 7

The gradient of the curve $x^2 + (y-1)^2 = 4$ at the point in the third quadrant where $x = -1$ is:

- A. $\frac{1}{\sqrt{3}}$
- B. $-\frac{1}{\sqrt{3}}$
- C. $\frac{1}{\sqrt{5}}$
- D. $-\frac{1}{\sqrt{5}}$
- E. $1 - \sqrt{3}$

Question 8

The rule of the relation determined by the parametric equations $x = 2\operatorname{cosec}(t) + 1$ and $y = 3\cot(t) - 1$

- A. $\frac{(x-1)^2}{4} - \frac{(y+1)^2}{9} = 1$
- B. $\frac{(y+1)^2}{9} - \frac{(x-1)^2}{4} = 1$
- C. $\frac{(x-1)^2}{4} + \frac{(y+1)^2}{9} = 1$
- D. $\frac{(y+1)^2}{3} - \frac{(x-1)^2}{2} = 1$
- E. $\frac{(x-1)^2}{2} - \frac{(y+1)^2}{3} = 1$

SECTION 1 - continued
TURN OVER

Question 9

The region in the complex plane that is **outside** the circle of radius b centred at the origin is given by the set of points z , where $z \in C$, such that

- A. $|z| < b$
- B. $|z| > b$
- C. $|z| < b^2$
- D. $|z| > b^2$
- E. $|z| = b$


Question 10

The solutions of the equation $z^2 = -2 - 2\sqrt{3}i$ in polar form are:

- A. $4 \operatorname{cis}\left(-\frac{\pi}{3}\right), 4 \operatorname{cis}\left(\frac{2\pi}{3}\right)$
- B. $4 \operatorname{cis}\left(-\frac{\pi}{3}\right), 4 \operatorname{cis}\left(\frac{\pi}{3}\right)$
- C. $4 \operatorname{cis}\left(-\frac{2\pi}{3}\right), 4 \operatorname{cis}\left(\frac{\pi}{3}\right)$
- D. $2 \operatorname{cis}\left(-\frac{2\pi}{3}\right), 2 \operatorname{cis}\left(\frac{\pi}{3}\right)$
- E. $2 \operatorname{cis}\left(\frac{2\pi}{3}\right), 2 \operatorname{cis}\left(-\frac{\pi}{3}\right)$

SECTION 1 - continued

Question 11




The shaded region, with boundaries not included, of the complex plane shown above is best described by:

- A. $\left\{ z : \text{Arg}(z) > \frac{\pi}{3} \right\}$
- B. $\left\{ z : \text{Arg}(z) > \frac{\pi}{3} \right\} \cup \left\{ z : \text{Arg}(z) < \frac{\pi}{2} \right\}$
- C. $\left\{ z : \text{Arg}(z) > \frac{\pi}{3} \right\} \cap \left\{ z : \text{Arg}(z) < \frac{\pi}{2} \right\}$
- D. $\left\{ z : \text{Arg}(z) > \frac{\pi}{2} \right\} \cup \left\{ z : \text{Arg}(z) < \frac{\pi}{3} \right\}$
- E. $\left\{ z : \text{Arg}(z) > \frac{\pi}{2} \right\} \cap \left\{ z : \text{Arg}(z) < \frac{\pi}{3} \right\}$

SECTION 1 - continued
TURN OVER

Question 12

The right-angled triangle shown below has sides represented by vectors $\vec{a}$, $\vec{b}$ and $\vec{c}$.



Which one of the following statements is **not** true?

- A. $|\vec{a}|^2 = |\vec{b}|^2 + |\vec{c}|^2$
- B. $\vec{b} \cdot (\vec{a} - \vec{c}) = |\vec{b}|^2$
- C. $\vec{b} \cdot (\vec{a} - \vec{b}) = |\vec{b}| |\vec{c}|$
- D. $\vec{b} \cdot \vec{a} = |\vec{b}| |\vec{a}| \cos(\theta)$
- E. $\vec{c} \cdot \vec{a} = |\vec{c}| |\vec{a}| \sin(\theta)$

Question 13

A body of mass 4 kg slides from rest down a sloping plane of length 3 m. If it takes 2 seconds to slide down the plane, the body's momentum at the bottom of the plane, in kg m/s, is:

- A. 8
- B. 12
- C. 24
- D. 36
- E. 48

SECTION 1 - continued

Question 14

Two forces $\vec{P} = 2\vec{i} + 3\vec{j}$ and $\vec{Q} = 3\vec{i} - 4\vec{j}$ act on a particle, of mass 2kg, at rest. The magnitude of the acceleration of the particle, in m/s^2 , is:

- A. 2.5
- B. 4.9
- C. 5
- D. 5.1
- E. 6.5

Question 15

Euler's method, with a step size of 0.2, is used to solve the differential equation $\frac{dy}{dx} = e^{\sqrt{\frac{x}{2}}}$, with initial condition $y = 2$, when $x = 0$. The approximation for y when $x = 0.4$ is given by:


- A. $2 + 0.4e^{\sqrt{0.2}}$
- B. $2 + 0.4e^{\sqrt{0.1}}$
- C. $2.2 + 0.2e^{\sqrt{0.1}}$
- D. $2.2 + 0.2e^{\sqrt{0.2}}$
- E. $2 + 0.2e^{\sqrt{0.1}} + 0.2e^{\sqrt{0.1}}$

SECTION 1 - continued
TURN OVER

Question 16

A 50L tank initially contains an acid solution of concentration 50%. Pure water flows into the tank at 2L/min. The solution is kept uniform by stirring, and flows out through a hole at the bottom of the tank at 2L/min. A differential equation for the mount of pure acid A litres in the tank after t minutes is:

- A. $\frac{dA}{dt} = \frac{25}{A}$
- B. $\frac{dA}{dt} = -\frac{A}{25}$
- C. $\frac{dA}{dt} = 2 - \frac{25}{A}$
- D. $\frac{dA}{dt} = 2 + \frac{25}{A}$
- E. $\frac{dA}{dt} = \frac{25}{A} - 4$

Question 17

The differential equation which best represents the above slope field could be:

- A. $\frac{dy}{dx} = \frac{1}{2x}$
- B. $\frac{dy}{dx} = \frac{1}{x^2}$
- C. $\frac{dy}{dx} = x^3$
- D. $\frac{dy}{dx} = x^2$
- E. $\frac{dy}{dx} = \frac{1}{x^3}$

SECTION 1 - continued

Question 18

A body of mass 5.0 kg is sliding down a plane of inclination 15° . If the inclination is increased to 30° , by how much would the acceleration increase?

- A. $\frac{\sqrt{6}-\sqrt{2}-2}{4}g$
 B. $\frac{5(\sqrt{6}-\sqrt{2}-2)}{4}g$
 C. $\frac{\sqrt{6}-2\sqrt{3}+\sqrt{2}}{4}g$
 D. $\frac{5(\sqrt{6}-2\sqrt{3}+\sqrt{2})}{4}g$
 E. $\frac{\sqrt{6}-\sqrt{2}}{4}g$

Question 19

The region bounded by the lines $x = 0$, $y = 3$ and the graph of $y = x^{\frac{4}{3}}$ where $x \geq 0$ is rotated about the **y-axis** to form a solid of revolution.

The volume of this solid is:

- A. $\frac{81\pi \times \sqrt[3]{9}}{11}$
 B. $\frac{12\pi \times \sqrt[4]{27}}{7}$
 C. $\frac{18\pi \times \sqrt{3}}{5}$
 D. $\frac{27\pi \times \sqrt[3]{3}}{11}$
 E. $\frac{81\pi}{4}$

SECTION 1 - continued
TURN OVER

Question 20

The velocity of a particle moving in a straight line is given by $v = \sqrt{9 - x^2}$, $x > 0$. The acceleration, in m/s^2 , of the particle when velocity is $\sqrt{5} \text{ m/s}$ is:

- A. -2
- B. 2
- C. 4
- D. -4
- E. 8

Question 21

A particle of mass 2 kg moves in a straight line with an initial velocity of 20 m/s. A constant force opposing the direction of the motion acts on the particle so that after 4 seconds its velocity is 2 m/s.

The magnitude of the force, in newtons, is:

- A. 4.5
- B. 6
- C. 9
- D. 18
- E. 36

Question 22

A mass of 60kg sits on the floor of a lift moving down, but decelerating at $\frac{g}{4} \text{ m/s}^2$, where g is the acceleration due to gravity. The reaction of the lift floor on the body, in Newtons, is:

- A. 45
- B. 60
- C. 75
- D. $45g$
- E. $75g$

END OF SECTION 1

SECTION 2

Instructions for Section 2

Answer **all** questions.

A decimal approximation will not be accepted if the question specifically asks for an **exact** answer.

For questions worth more than one mark, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams are **not** drawn to scale.

Take the **acceleration due to gravity**, to have magnitude $g \text{ m/s}^2$, where $g = 9.8$.

Question 1 (11 marks)

A curve is defined by the parametric equations

$$x = -3 - 2 \cos\left(\frac{t}{2}\right) \quad \text{and} \quad y = 4 + 3 \sin\left(\frac{t}{2}\right) \quad \text{for } t \in [0, 4\pi]$$

- a.** Find the Cartesian equation of the curve.

2 marks

- b.** Find the values of t for which the gradient of the curve is $-\frac{3\sqrt{3}}{2}$.


4 marks

SECTION 2 - Question1 - continued
TURN OVER

- c. Sketch the graph of the curve whose parametric equations are given below. Label the axes intercepts.

$$x = 1 - 2\cos(t) \quad \text{and} \quad y = 3\sin(t)$$

2 marks



The region in the first quadrant bounded by the above graph, x -axis and lines $x = 1$ and $x = 2.5$ is rotated about the x -axis to form a solid of revolution.

- d. Write down an integral that will give the volume of this solid.

2 marks

- e. Find the volume of this solid of revolution.

1 mark

SECTION 2 - continued

Question 2 (10 marks)


Let the regions S_1 and S_2 be defined as follows on the complex plane.

$$S_1 : \{z : |z - 2| = 2, z \in \mathbb{C}\}$$

$$S_2 : \{z : \operatorname{Re}(z) + \operatorname{Im}(z) = 4, z \in \mathbb{C}\}$$

- a.** Sketch both the regions on the axes below. Shade the region bound by the two lines.

4 marks



- b.** Find the area of the shaded region.

2 marks

SECTION 2 - Question 2 - continued
TURN OVER

S_3 is the region defined as $S_3 : \{z : |z + 2 + 2i| = 2, z \in \mathbb{C}\}$

c. Sketch the region S_3 on the axes given in part a. 2 marks

d. Find the minimum value of $|z_1 - z_3|$ where $z_1 \in S_1$ and $z_3 \in S_3$ 2 marks

Question 3 (16 marks)

Two bodies of mass 3 kg and 7 kg rest on a smooth horizontal plane. The bodies are acted on by a force of F newtons, acting in the horizontal direction, which just brings the body to the point of moving.

a. Draw all the forces acting on the body in the diagram below. 4 marks



b. What is the acceleration of the two blocks in terms of F ? 2 marks

SECTION 2 - Question 3 - continued


- c. What is the magnitude of force applied by the 3kg block on the 7 kg block? 2 marks

- d. If the force applied on the block is 120N, find the acceleration of the two blocks.

1 mark

A block of mass 10kg rests on a smooth inclined plane making an angle of 30° with the horizontal.

- e. Mark all the forces acting on the body in the diagram below. 2 marks



SECTION 2 - Question 3 - continued
TURN OVER

f. Find the acceleration of the block.

3 marks

g. Find the force exerted by the block on the incline, correct to two decimal places.

2 marks

SECTION 2 - continued

Question 4 (12 marks)

The position vectors, at time t seconds, where $t \geq 0$, of two particles A and B are given respectively by

$$\mathbf{r}_{\sim A}(t) = (t^3 - 9t + 8)\mathbf{i} + t^2\mathbf{j}$$

$$\mathbf{r}_{\sim B}(t) = (2 - t^2)\mathbf{i} + (3t - 2)\mathbf{j}$$

- a.** Prove that the two particles collide. Find the time when they collide. 4 marks

- b.** Show that the particles are at the same speed at the time of collision. Hence, find the speed. 3 marks

- c.** Find $\dot{\mathbf{r}}_{\sim A}(t) \bullet \dot{\mathbf{r}}_{\sim B}(t)$ at the time of collision. 1 mark

SECTION 2 - Question 4 - continued
TURN OVER

d. Interpret the result obtained in part **c**.

2 marks

e. Find the acceleration of particle *B* at the time of collision.

2 marks

Question 5 (9 marks)

The number of insects, *N*, in a certain area after *t* years, may be modelled by

$$\log_e N = 6 - 3e^{-0.4t}, \quad t \geq 0$$

a. Verify that $\log_e N = 6 - 3e^{-0.4t}$ satisfies the differential equation

2 marks

$$\frac{1}{N} \frac{dN}{dt} + 0.4 \log_e N - 2.4 = 0$$

SECTION 2 - Question 5 - continued

- b.** Find the initial number of insects in the area. Give your answer to the nearest integer.

1 mark

- c.** Using this mathematical model, find the limiting number of insects that would eventually be present in this area.

2 marks

- d.** Find $\frac{d^2 N}{dt^2}$ in terms of N and $\log_e N$

2 marks

SECTION 2 - Question 5 - continued
TURN OVER

- e. The graph of N as a function of t has a point of inflection. Find the value of the coordinates of this point, correct to the nearest integer values. 2 marks

END OF QUESTION AND ANSWER BOOK

Multiple choice answer sheet

Instructions: Circle the correct response

1.	A	B	C	D	E
2.	A	B	C	D	E
3.	A	B	C	D	E
4.	A	B	C	D	E
5.	A	B	C	D	E
6.	A	B	C	D	E
7.	A	B	C	D	E
8.	A	B	C	D	E
9.	A	B	C	D	E
10.	A	B	C	D	E
11.	A	B	C	D	E
12.	A	B	C	D	E
13.	A	B	C	D	E
14.	A	B	C	D	E
15.	A	B	C	D	E
16.	A	B	C	D	E
17.	A	B	C	D	E
18.	A	B	C	D	E
19.	A	B	C	D	E
20.	A	B	C	D	E
21.	A	B	C	D	E
22.	A	B	C	D	E