

Victorian Certificate of Education
2016

SUPERVISOR TO ATTACH PROCESSING LABEL HERE

STUDENT NUMBER

									Letter
--	--	--	--	--	--	--	--	--	--------

SPECIALIST MATHEMATICS

Written examination 1

Friday 4 November 2016

Reading time: 9.00 am to 9.15 am (15 minutes)

Writing time: 9.15 am to 10.15 am (1 hour)

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
10	10	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or correction fluid/tape.

Materials supplied

- Question and answer book of 8 pages.
- Formula sheet.
- Working space is provided throughout the book.

Instructions

- Write your **student number** in the space provided above on this page.
- Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.
- All written responses must be in English.

At the end of the examination

- You may keep the formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

THIS PAGE IS BLANK

Instructions

Answer **all** questions in the spaces provided.

Unless otherwise specified, an **exact** answer is required to a question.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.


Take the **acceleration due to gravity** to have magnitude $g \text{ ms}^{-2}$, where $g = 9.8$

Question 1 (4 marks)

A taut rope of length $1\frac{2}{3} \text{ m}$ suspends a mass of 20 kg from a fixed point O . A horizontal force of P newtons displaces the mass by 1 m horizontally so that the taut rope is then at an angle of θ to the vertical.

- a. Show all the forces acting on the mass on the diagram below.

1 mark



- b. Show that $\sin(\theta) = \frac{3}{5}$.

1 mark

- c. Find the magnitude of the tension force in the rope in newtons.

2 marks

TURN OVER

Question 2 (3 marks)

A farmer grows peaches, which are sold at a local market. The mass, in grams, of peaches produced on this farm is known to be normally distributed with a variance of 16. A bag of 25 peaches is found to have a total mass of 2625 g.

Based on this sample of 25 peaches, calculate an approximate 95% confidence interval for the mean mass of all peaches produced on this farm. Use an integer multiple of the standard deviation in your calculations.

Question 3 (4 marks)

Find the equation of the line perpendicular to the graph of $\cos(y) + y \sin(x) = x^2$ at $\left(0, -\frac{\pi}{2}\right)$.

Question 4 (4 marks)

Chemicals are added to a container so that a particular crystal will grow in the shape of a cube. The side length of the crystal, x millimetres, t days after the chemicals were added to the container, is given by $x = \arctan(t)$.

Find the rate at which the surface area, A square millimetres, of the crystal is growing one day after the chemicals were added. Give your answer in square millimetres per day.

Question 5 (4 marks)

Consider the vectors $\underline{a} = 3\underline{i} + 5\underline{j} - 2\underline{k}$, $\underline{b} = \underline{i} - 2\underline{j} + 3\underline{k}$ and $\underline{c} = \underline{i} + d\underline{k}$, where d is a real constant.

- a. Find the vector resolute of $\underline{a}$ in the direction of $\underline{b}$. 2 marks

- b. Find the value of d if the vectors are **linearly dependent**. 2 marks

Question 6 (3 marks)

Write $\frac{(1-\sqrt{3}i)^4}{1+\sqrt{3}i}$ in the form $a + bi$, where a and b are real constants.

Question 7 (4 marks)

Find the arc length of the curve $y = \frac{1}{3}(x^2 + 2)^{\frac{3}{2}}$ from $x = 0$ to $x = 2$.

Question 8 (6 marks)

The position of a body with mass 3 kg from a fixed origin at time t seconds, $t \geq 0$, is given by

$\underline{r} = (3 \sin(2t) - 2)\underline{i} + (3 - 2 \cos(2t))\underline{j}$, where components are in metres.

- a.** Find an expression for the speed, in metres per second, of the body at time t . 2 marks

- b.** Find the speed of the body, in metres per second, when $t = \frac{\pi}{12}$. 1 mark

- c.** Find the maximum magnitude of the net force acting on the body in newtons. 3 marks

Question 9 (3 marks)

Given that $\cos(x - y) = \frac{3}{5}$ and $\tan(x) \tan(y) = 2$, find $\cos(x + y)$.

Question 10 (5 marks)

Solve the differential equation $\sqrt{2 - x^2} \frac{dy}{dx} = \frac{1}{2 - y}$, given that $y(1) = 0$. Express y as a function of x .

**Victorian Certificate of Education
2016**

SPECIALIST MATHEMATICS

Written examination 1

FORMULA SHEET

Instructions

This formula sheet is provided for your reference.
A question and answer book is provided with this formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Specialist Mathematics formulas

Mensuration

area of a trapezium	$\frac{1}{2}(a+b)h$
curved surface area of a cylinder	$2\pi rh$
volume of a cylinder	$\pi r^2 h$
volume of a cone	$\frac{1}{3}\pi r^2 h$
volume of a pyramid	$\frac{1}{3}Ah$
volume of a sphere	$\frac{4}{3}\pi r^3$
area of a triangle	$\frac{1}{2}bc \sin(A)$
sine rule	$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$
cosine rule	$c^2 = a^2 + b^2 - 2ab \cos(C)$

Circular functions

$\cos^2(x) + \sin^2(x) = 1$	
$1 + \tan^2(x) = \sec^2(x)$	$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$
$\sin(x+y) = \sin(x)\cos(y) + \cos(x)\sin(y)$	$\sin(x-y) = \sin(x)\cos(y) - \cos(x)\sin(y)$
$\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$	$\cos(x-y) = \cos(x)\cos(y) + \sin(x)\sin(y)$
$\tan(x+y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$	$\tan(x-y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$
$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$	
$\sin(2x) = 2\sin(x)\cos(x)$	$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$

Circular functions – continued

Function	$\sin^{-1}$ or arcsin	$\cos^{-1}$ or arccos	$\tan^{-1}$ or arctan
Domain	$[-1, 1]$	$[-1, 1]$	R
Range	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$[0, \pi]$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Algebra (complex numbers)

$z = x + iy = r(\cos(\theta) + i\sin(\theta)) = r \operatorname{cis}(\theta)$	
$ z = \sqrt{x^2 + y^2} = r$	$-\pi < \operatorname{Arg}(z) \leq \pi$
$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$	$\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$
$z^n = r^n \operatorname{cis}(n\theta)$ (de Moivre's theorem)	

Probability and statistics

for random variables X and Y	$E(aX + b) = aE(X) + b$ $E(aX + bY) = aE(X) + bE(Y)$ $\operatorname{var}(aX + b) = a^2 \operatorname{var}(X)$
for independent random variables X and Y	$\operatorname{var}(aX + bY) = a^2 \operatorname{var}(X) + b^2 \operatorname{var}(Y)$
approximate confidence interval for μ	$\left(\bar{x} - z \frac{s}{\sqrt{n}}, \bar{x} + z \frac{s}{\sqrt{n}} \right)$
distribution of sample mean $\bar{X}$	mean $E(\bar{X}) = \mu$ variance $\operatorname{var}(\bar{X}) = \frac{\sigma^2}{n}$

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e x + c$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$	$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$
$\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$	$\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\cos^{-1}(x)) = \frac{-1}{\sqrt{1-x^2}}$	$\int \frac{-1}{\sqrt{a^2-x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{1+x^2}$	$\int \frac{a}{a^2+x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$
	$\int (ax+b)^n dx = \frac{1}{a(n+1)} (ax+b)^{n+1} + c, n \neq -1$
	$\int (ax+b)^{-1} dx = \frac{1}{a} \log_e ax+b + c$
product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$
quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
Euler's method	If $\frac{dy}{dx} = f(x)$, $x_0 = a$ and $y_0 = b$, then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + hf(x_n)$
acceleration	$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$
arc length	$\int_{x_1}^{x_2} \sqrt{1+(f'(x))^2} dx$ or $\int_{t_1}^{t_2} \sqrt{(x'(t))^2 + (y'(t))^2} dt$

Vectors in two and three dimensions

$\underline{r} = x\hat{i} + y\hat{j} + z\hat{k}$
$ \underline{r} = \sqrt{x^2 + y^2 + z^2} = r$
$\dot{\underline{r}} = \frac{d\underline{r}}{dt} = \frac{dx}{dt}\hat{i} + \frac{dy}{dt}\hat{j} + \frac{dz}{dt}\hat{k}$
$\underline{r}_1 \cdot \underline{r}_2 = r_1 r_2 \cos(\theta) = x_1 x_2 + y_1 y_2 + z_1 z_2$

Mechanics

momentum	$\underline{p} = m\underline{v}$
equation of motion	$\underline{R} = m\underline{a}$