

Year 2017

VCE

Specialist Mathematics

Trial Examination 1

Solutions



KILBAHA MULTIMEDIA PUBLISHING
PO BOX 2227
KEW VIC 3101
AUSTRALIA

TEL: (03) 9018 5376
FAX: (03) 9817 4334
kilbaha@gmail.com
<http://kilbaha.com.au>

IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Multimedia Publishing.
- The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
- For authorised copying within Australia please check that your institution has a licence from **Copyright Agency Limited**. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.

Reproduction and communication for educational purposes The Australian Copyright Act 1968 (the Act) allows a maximum of one chapter or 10% of the pages of this work, to be reproduced and/or communicated by any educational institution for its educational purposes provided that educational institution (or the body that administers it) has given a remuneration notice to Copyright Agency Limited (CAL) under the Act.

For details of the CAL licence for educational institutions contact
CAL, Level 15, 233 Castlereagh Street, Sydney, NSW, 2000

Tel: (02) 9394 7600

Fax: (02) 9394 7601

Email: info@copyright.com.au

Web: <http://www.copyright.com.au>

- While every care has been taken, no guarantee is given that these answers are free from error. Please contact us if you believe you have found an error.

Question 1

a. $z = \sqrt{2} \operatorname{cis}\left(-\frac{2\pi}{3}\right)$

$$= \sqrt{2} \left(\cos\left(-\frac{2\pi}{3}\right) + i \sin\left(-\frac{2\pi}{3}\right) \right) \text{ since } \cos(-\theta) = \cos(\theta), \sin(-\theta) = -\sin(\theta)$$

$$= \sqrt{2} \left(\cos\left(\frac{2\pi}{3}\right) - i \sin\left(\frac{2\pi}{3}\right) \right)$$

$$= \sqrt{2} \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i \right) \quad \text{M1}$$

$$= \frac{\sqrt{2}}{2} (-1 - \sqrt{3}i)$$

by the conjugate root theorem, $\bar{z} = \frac{\sqrt{2}}{2} (-1 + \sqrt{3}i)$ is also a root

$$z + \bar{z} = -\sqrt{2}, \quad z \cdot \bar{z} = \frac{2}{4} (1 - 3i^2) = 2$$

$z^2 - (\text{sum of the roots})z + \text{product of the roots}$

$$= (z^2 + \sqrt{2}z + 2) \text{ is the quadratic factor} \quad \text{A1}$$

b. $f(z) = z^4 + \sqrt{2}z^3 + 5z^2 + 3\sqrt{2}z + 6 = 0$

$$= (z^2 + \sqrt{2}z + 2)(z^2 + 3) = 0$$

$$= \left(z - \frac{\sqrt{2}}{2}(-1 + \sqrt{3}i) \right) \left(z - \frac{\sqrt{2}}{2}(-1 - \sqrt{3}i) \right) (z - \sqrt{3}i)(z + \sqrt{3}i) = 0 \quad \text{M1}$$

all the roots are $z = \frac{\sqrt{2}}{2}(-1 \pm \sqrt{3}i), \pm \sqrt{3}i$ A1

Question 2


95% confidence interval is $\mu \pm 2 \times \frac{\mu}{2\sqrt{n}} = \left(\mu \left(1 - \frac{1}{\sqrt{n}} \right), \mu \left(1 + \frac{1}{\sqrt{n}} \right) \right)$

this has a width of $\frac{2\mu}{\sqrt{n}} \leq \frac{\mu}{10}$ M1

$$\Rightarrow \sqrt{n} \geq 20$$

the minimum sample size is 400 A1

Question 3



when the 4kg mass moves downwards,
with an acceleration of a , resolving

$$(1) \quad 4g - T_1 = 4a$$


$$(2) \quad T_1 - 2g = 2a$$

adding (1) + (2)

$$2g = 6a$$

$$a = \frac{g}{3}$$

A1



when the $2 + m$ kg moves downwards
with an acceleration of $\frac{g}{3}$, resolving

$$(3) \quad (2 + m)g - T_2 = (2 + m)\frac{g}{3}$$

$$(4) \quad T_2 - 4g = 4 \times \frac{g}{3}$$

adding (3) + (4)

$$(2 + m - 4)g = \frac{g}{3}(2 + m + 4)$$

$$3(m - 2) = m + 6$$

$$3m - 6 = m + 6$$

$$2m = 12$$

$$m = 6$$

M1

A1

Question 4

$$\log_e(2xy) + \frac{x}{y} = 8 \quad \text{by log laws}$$

$$\log_e(2) + \log_e(x) + \log_e(y) + \frac{x}{y} = 8 \quad \text{using implicit differentiation} \quad \text{M1}$$

$$\frac{1}{x} + \frac{1}{y} \frac{dy}{dx} + \frac{1}{y} - \frac{x}{y^2} \frac{dy}{dx} = 0$$

$$\frac{1}{x} + \frac{1}{y} = \frac{dy}{dx} \left(\frac{x}{y^2} - \frac{1}{y} \right) \quad \text{A1}$$

$$\frac{x+y}{xy} = \frac{dy}{dx} \left(\frac{x-y}{y^2} \right)$$

$$\frac{dy}{dx} = \frac{y(x+y)}{x(x-y)} \quad \text{A1}$$

$$\text{at the point } \left(2, \frac{1}{4} \right)$$

$$\left. \frac{dy}{dx} \right|_{\left(2, \frac{1}{4} \right)} = \frac{\frac{1}{4} \left(2 + \frac{1}{4} \right)}{2 \left(2 - \frac{1}{4} \right)} = \frac{1}{8} \times \frac{9}{7} = \frac{9}{56}$$

$$\text{gradient of the normal is } -\frac{56}{9} \quad \text{A1}$$

Question 5

$$\text{a.} \quad s = \int_{t_1}^{t_2} \sqrt{(x'(t))^2 + (y'(t))^2} dt$$

$$x = t^3 + 5, \quad y = 6t^2 - 1$$

$$\dot{x} = \frac{dx}{dt} = 3t^2 \quad \dot{y} = \frac{dy}{dt} = 12t \quad \text{M1}$$

$$\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 = 9t^4 + 144t^2 = 9t^2(t^2 + 16)$$

$$\sqrt{\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2} = 3t\sqrt{t^2 + 16} \quad \text{since } t \geq 0 \quad \text{A1}$$

$$s = \int_0^3 3t\sqrt{t^2 + 16} dt$$

b. $s = \int_0^3 3t\sqrt{t^2+16} dt$ let $u = t^2 + 16$ $\frac{du}{dt} = 2t \Rightarrow dt = \frac{1}{2t} du$

terminals when $t = 0 \Rightarrow u = 16$ when $t = 3 \Rightarrow u = 25$

$$s = \int_{16}^{25} 3t\sqrt{u} \times \frac{1}{2t} du \quad \text{M1}$$

$$s = \frac{3}{2} \int_{16}^{25} u^{\frac{1}{2}} du$$

$$= \frac{3}{2} \left[\frac{2}{\frac{3}{2}} u^{\frac{3}{2}} \right]_{16}^{25}$$

$$= 25^{\frac{3}{2}} - 16^{\frac{3}{2}}$$

$$= (\sqrt{25})^3 - (\sqrt{16})^3$$

$$= 5^3 - 4^3 = 125 - 64$$

$$= 61$$

A1

Question 6

$$\frac{dy}{dx} = \frac{\sqrt{4-y^2}}{4+x^2} \text{ and } y(0)=1 \text{ separating the variables}$$


$$\int \frac{1}{\sqrt{4-y^2}} dy = \int \frac{1}{4+x^2} dx \quad \text{M1}$$

$$\sin^{-1}\left(\frac{y}{2}\right) = \frac{1}{2} \tan^{-1}\left(\frac{x}{2}\right) + c \quad \text{A1}$$

$$\text{when } y=1, x=0 \quad \sin^{-1}\left(\frac{1}{2}\right) = c = \frac{\pi}{6}$$

$$\sin^{-1}\left(\frac{y}{2}\right) = \frac{\pi}{6} + \frac{1}{2} \tan^{-1}\left(\frac{x}{2}\right)$$

$$y = 2 \sin\left(\frac{\pi}{6} + \frac{1}{2} \tan^{-1}\left(\frac{x}{2}\right)\right) \quad \text{A1}$$

Question 7**a.**

A1

b.Let T be the tension in the strings, all the forces are in newtons.resolving horizontally (1) $F + T \cos(60^\circ) - T \cos(30^\circ) = 0$ resolving vertically (2) $T \sin(60^\circ) + T \sin(30^\circ) - mg = 0$

A1

$$(1) \quad F = T(\cos(30^\circ) - \cos(60^\circ)) \Rightarrow F = T\left(\frac{\sqrt{3}-1}{2}\right) \Rightarrow T = \frac{2F}{\sqrt{3}-1}$$

$$(2) \quad mg = T(\sin(60^\circ) + \sin(30^\circ)) \Rightarrow mg = T\left(\frac{\sqrt{3}+1}{2}\right) \Rightarrow T = \frac{2mg}{\sqrt{3}+1}$$

M1

eliminating T , gives $T = \frac{2F}{\sqrt{3}-1} = \frac{2mg}{\sqrt{3}+1} \Rightarrow F = \left(\frac{\sqrt{3}-1}{\sqrt{3}+1}\right)mg$

$$F = \left(\frac{\sqrt{3}-1}{\sqrt{3}+1}\right)mg \times \left(\frac{\sqrt{3}-1}{\sqrt{3}-1}\right) = \left(\frac{3-2\sqrt{3}+1}{3-1}\right)mg$$

$$F = (2-\sqrt{3})mg \quad a=2, b=3$$

A1

Question 8

a.i. Let $y = \sec(kx) = \frac{1}{\cos(kx)} = (\cos(kx))^{-1}$ using the chain rule

$$\frac{dy}{dx} = -1 \times -k \sin(kx) \times (\cos(kx))^{-2} = \frac{k \sin(kx)}{\cos^2(kx)} = \frac{k \sin(kx)}{\cos(kx)} \times \frac{1}{\cos(kx)}$$

A1

$$\frac{d}{dx} [\sec(kx)] = k \tan(kx) \sec(kx)$$

ii Let $y = \log_e(\tan(kx) + \sec(kx)) = \log_e(u)$ where $u = \tan(kx) + \sec(kx)$

$$\frac{dy}{du} = \frac{1}{u} \quad \text{using the chain rule}$$

$$\frac{du}{dx} = k \sec^2(kx) + k \sec(kx) \tan(kx) = k \sec(kx) (\tan(kx) + \sec(kx))$$

A1


$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = \frac{1}{\tan(kx) + \sec(kx)} \times k \sec(kx) (\sec(kx) + \tan(kx)) = k \sec(kx)$$

$$\text{so that } \frac{d}{dx} [\log_e(\tan(kx) + \sec(kx))] = k \sec(kx)$$

b. $y = 2 \sec\left(\frac{x}{2}\right) = \frac{2}{\cos\left(\frac{x}{2}\right)}$ when $x = 0$, $y = 2$, y intercept $(0, 2)$

The graph has vertical asymptotes when $\cos\left(\frac{x}{2}\right) = 0 \Rightarrow \frac{x}{2} = \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}$

so that $x = \pm\pi, \pm 3\pi$ are the vertical asymptotes. A1



c. $A = \int_0^{\frac{\pi}{2}} 2 \sec\left(\frac{x}{2}\right) dx$ using a. with $k = \frac{1}{2}$

$$A = \left[4 \log_e \left(\tan\left(\frac{x}{2}\right) + \sec\left(\frac{x}{2}\right) \right) \right]_0^{\frac{\pi}{2}}$$

A1

$$= 4 \log_e \left(\tan\left(\frac{\pi}{4}\right) + \sec\left(\frac{\pi}{4}\right) \right) - 4 \log_e (\tan(0) + \sec(0))$$

$$= 4 \log_e (1 + \sqrt{2}) \text{ units}^2$$

A1

Question 9

Let $\underline{a} = 2\underline{i} - 3\underline{j} + \underline{k}$, $\underline{b} = \underline{i} + 2\underline{j} + 4\underline{k}$, $\underline{c} = x\underline{i} + y\underline{j} + z\underline{k}$

Since $\underline{a}$ is perpendicular to $\underline{c}$ $\underline{a} \cdot \underline{c} = 0 \Rightarrow (1) \ 2x - 3y + z = 0$

Since $\underline{b}$ is perpendicular to $\underline{c}$ $\underline{b} \cdot \underline{c} = 0 \Rightarrow (2) \ x + 2y + 4z = 0$ A1

Since $\underline{c}$ is a unit vector $|\underline{c}| = 1 \Rightarrow x^2 + y^2 + z^2 = 1$

$$(1) \ 2x - 3y + z = 0$$

$$2 \times (2) \ 2x + 4y + 8z = 0 \quad \text{subtracting gives } 7y + 7z = 0 \Rightarrow z = -y$$

$$4 \times (1) \ 8x - 12y + 4z = 0$$

M1

$$(2) \ x + 2y + 4z = 0 \quad \text{subtracting gives } 7x - 14y = 0 \Rightarrow x = 2y$$

substituting into (3) $4y^2 + y^2 + y^2 = 6y^2 = 1 \Rightarrow y = \pm \frac{1}{\sqrt{6}}$

$$\hat{\underline{c}} = \frac{\pm 1}{\sqrt{6}} (2\underline{i} + \underline{j} - \underline{k})$$

A1

Question 10

a. $\left. \frac{dB}{dt} \right|_{B=60} = \frac{60}{600}(200-60) = 14 \text{ gm/month}$

$$\left. \frac{dB}{dt} \right|_{B=120} = \frac{120}{600}(200-120) = 16 \text{ gm/month}$$

Since when $B = 120$ the rate is larger,
the bird is gaining weight faster when $B = 120$

A1

b. $\frac{dB}{dt} = \frac{B}{600}(200-B)$ separating the variables

$$\int \frac{600}{B(200-B)} dB = t \quad \text{using partial fractions}$$

$$\frac{600}{B(200-B)} = \frac{A}{B} + \frac{D}{200-B} = \frac{A(200-B) + DB}{B(200-B)} = \frac{B(D-A) + 200A}{B(200-B)}$$

M1

$$\Rightarrow 200A = 600 \quad D - A = 0 \Rightarrow A = D = 3$$

$$3 \int \left(\frac{1}{B} + \frac{1}{200-B} \right) dB = t$$

$$\frac{t}{3} = \log_e(|B|) - \log_e(|200-B|) + c$$

A1

since $20 \leq B < 200$ the moduli are not needed

Now when $t = 0$, $B = 20$

$$0 = \log_e 20 - \log_e (200-20) + c \Rightarrow$$

$$c = \log_e (180) - \log_e (20) = \log_e \left(\frac{180}{20} \right) = \log_e (9)$$

A1

$$\frac{t}{3} = \log_e B - \log_e (200-B) + \log_e (9)$$

$$\frac{t}{3} = \log_e \left(\frac{9B}{200-B} \right)$$

$$e^{\frac{t}{3}} = \frac{9B}{200-B} \Rightarrow \frac{200-B}{9B} = e^{-\frac{t}{3}}$$

$$200-B = 9Be^{-\frac{t}{3}}$$

$$200 = B \left(1 + 9e^{-\frac{t}{3}} \right)$$

M1

$$B(t) = \frac{200}{1 + 9e^{-\frac{t}{3}}}$$

c. $\frac{dB}{dt} = \frac{1}{600}(200B - B^2)$
 $\frac{d^2B}{dt^2} = \frac{1}{600}(200 - 2B)\frac{dB}{dt} = \frac{1}{300}(100 - B)\frac{dB}{dt} = 0$ for inflexion points $\frac{d^2B}{dt^2} = 0$
 $\Rightarrow B = 100$

when $B(t) = 100 = \frac{200}{1 + 9e^{-\frac{t}{3}}} \Rightarrow 1 + 9e^{-\frac{t}{3}} = 2$ M1

$9e^{-\frac{t}{3}} = 1 \Rightarrow e^{\frac{t}{3}} = 9$


$t = 3\log_e(9)$

point of inflexion $(3\log_e(9), 100)$

the curve is concave up for $0 \leq t < 3\log_e(9)$ A1

d. Since $B(0) = 20$ and as $t \rightarrow \infty$, $B \rightarrow 200$, the horizontal asymptote is 200

G1



END OF SUGGESTED SOLUTIONS