

# **SPECIALIST MATHEMATICS**

**Units 3 & 4 – Written examination 2**



**2017 Trial Examination**

**SOLUTIONS**

**SECTION A: Multiple-choice questions (1 mark each)****Question 1****Answer: D***Explanation:*

$$x = \frac{1}{3} \cos(2t) = \frac{1}{3} (1 - 2 \sin^2(t)) = \frac{1}{3} (1 - 2(2y)^2) \Rightarrow 3x + 8y^2 = 1$$

**Question 2****Answer: B***Explanation*

$$\begin{aligned} -\frac{\pi}{2} &< \arctan(3x + 1) < \frac{\pi}{2} \\ -\frac{3\pi}{2} &< -3 \arctan(3x + 1) < \frac{3\pi}{2} \\ 2 \arctan(\sqrt{3}) - \frac{3\pi}{2} &< 2 \arctan(\sqrt{3}) - 3 \arctan(3x + 1) < 2 \arctan(\sqrt{3}) + \frac{3\pi}{2} \\ -\frac{5\pi}{6} = \frac{2\pi}{3} - \frac{3\pi}{2} &< 2 \arctan(\sqrt{3}) - 3 \arctan(3x + 1) < \frac{2\pi}{3} + \frac{3\pi}{2} = \frac{13\pi}{6} \end{aligned}$$

**Question 3****Answer: E***Explanation:*

Simplify by CAS

$$\text{expand} \left( \frac{2 \cdot x^3 + 5 \cdot x^2 + 2 \cdot x - 1}{x^2 - 1} \right) \qquad \frac{4}{x-1} + 2 \cdot x + 5$$

$f(x) = 2x + 5 + \frac{4}{x-1}$  has an oblique asymptote  $y = 2x + 5$   
 a vertical asymptote  $x = 1$

**Question 4****Answer: D***Explanation:*Let  $z_1 = 2 + 3i$ ,  $z_2 = 2 - 3i$  and  $z_3$  be the three roots of the polynomial.Then  $P(z) = (z - z_1)(z - z_2)(z - z_3)$ 

$$-z_1 z_2 z_3 = -39$$

$$z_3 = \frac{39}{z_1 z_2} = \frac{39}{13} = 3$$

Expand  $(z - z_1)(z - z_2)(z - z_3)$  by CAS

$$\text{expand}((z - (2 + 3 \cdot i)) \cdot (z - (2 - 3 \cdot i)) \cdot (z - 3)) \quad z^3 - 7 \cdot z^2 + 25 \cdot z - 39$$

$$b = -7, \quad c = 25$$

**Question 5****Answer: A***Explanation:*

Using CAS

Define  $\arg(a, b) = \text{angle}(a - 2 \cdot b \cdot i)$  Done

$$\arg(-6, -\sqrt{3}) \quad \frac{5 \cdot \pi}{6}$$

**Question 6****Answer: E***Explanation:*

$$\begin{aligned} z^4 + ri &= (z - z_1)(z - z_2)(z - z_3)(z - z_4) \\ &= z^4 - (z_1 + z_2 + z_3 + z_4)z^3 + (z_1 z_2 + z_2 z_3 + z_3 z_4 + z_4 z_1)z^2 \\ &\quad - (z_1 z_2 z_3 + z_2 z_3 z_4 + z_3 z_4 z_1 + z_4 z_1 z_2)z + z_1 z_2 z_3 z_4 \end{aligned}$$

Therefore

$$z_1 + z_2 + z_3 + z_4 = 0$$

$$z_1 z_2 + z_2 z_3 + z_3 z_4 + z_4 z_1 = 0$$

$$z_1 z_2 z_3 + z_2 z_3 z_4 + z_3 z_4 z_1 + z_4 z_1 z_2 = 0$$

$$z_1 z_2 z_3 z_4 = ri, \text{ hence it's impossible to have } z_1 = \overline{z_2}, \quad z_3 = \overline{z_4}.$$

OR

$$z^4 = -ri = r \operatorname{cis}\left(-\frac{\pi}{2}\right) \Rightarrow$$

$z_1, z_2, z_3$  and  $z_4$  have principal arguments  $-\frac{\pi}{8}, \frac{3\pi}{8}, \frac{7\pi}{8}$  and  $-\frac{5\pi}{8}$ . They are not in conjugate pairs.

**Question 7****Answer: C***Explanation:***Using CAS**Define  $x(t)=\tan(t)-2 \cdot t$  DoneDefine  $y(t)=2 \cdot \ln(\sec(t))$  Done  $\frac{d}{dt}(y(t))$   $2 \cdot \tan(t)$  $\frac{d}{dt}(x(t))$   $\frac{-\left(2 \cdot (\cos(t))^2 - 1\right)}{(\cos(t))^2}$   $\frac{\frac{2 \cdot \tan(t)}{-\left(2 \cdot (\cos(t))^2 - 1\right)}}{(\cos(t))^2}$   $\frac{-2 \cdot \sin(t) \cdot \cos(t)}{2 \cdot (\cos(t))^2 - 1}$  tCollect  $\left(\frac{-2 \cdot \sin(t) \cdot \cos(t)}{2 \cdot (\cos(t))^2 - 1}\right)$   $-\tan(2 \cdot t)$ **Question 8****Answer: D***Explanation:*Let  $u = x^3 - x^2 + 19x + 8.$ Then  $\frac{du}{dx} = 3x^2 - 2x + 19 \Rightarrow dx = \frac{1}{3x^2 - 2x + 19} du$ 

$$\int_0^1 ((3x^2 - 2x + 19)\sqrt[3]{x^3 - x^2 + 19x + 8})dx = \int_8^{27} u^{\frac{1}{3}} du$$

**Question 9**

**Answer: C**

*Explanation:*

From CAS we have

Define  $f(x) = \sin(x) + \cos(x)$

Done

$\text{euler}(f(x), x, y, \{0, 0.3\}, 1, 0.1)$

$\begin{bmatrix} 0. & 0.1 & 0.2 & 0.3 \\ 1. & 1.1 & 1.20948 & 1.32736 \end{bmatrix}$

Therefore


$$y_3 = y_2 + 0.1 \times f(0.2) = 1.21 + 0.1f(0.2)$$

**Question 10**

**Answer: E**

*Explanation:*

From CAS we have



**Question 11****Answer: D***Explanation:*

A useful mathematical model for setting up differential equations of dynamic systems

$$\frac{dQ}{dt} = R_{in} \times C_{in} - R_{out} \times C_{out}$$

where  $R_{in}$  and  $R_{out}$  are the flowing in and flowing out rate;  $C_{in}$  and  $C_{out}$  are the concentrations of the solutions which are flowing in and flowing out respectively.

Therefore

$$\frac{dQ}{dt} = 8 \times 200 - 6 \times \frac{Q}{300 + (8 - 6)t} = 1600 - \frac{3Q}{150 + t}$$

$$\frac{dQ}{dt} + \frac{3Q}{150 + t} = 1600$$

**Question 12****Answer: A***Explanation:*Solve  $\left| \tilde{u} - (\tilde{u} \cdot \tilde{v}) \tilde{v} \right| = \frac{\sqrt{10}}{2}$  for  $a$  in CASDefine  $u = \begin{bmatrix} -1 & a & 1 \end{bmatrix}$ 

Done

Define  $v = \begin{bmatrix} 1 & 3 & 2 \end{bmatrix}$ 

Done

Define  $vh = \text{unitV}(v)$ 

Done

$$\text{solve} \left( \text{norm} \left( u - \text{dotP}(u, vh) \cdot vh \right) = \frac{\sqrt{10}}{2}, a \right)$$

$$a = \frac{-4}{5} \text{ or } a = 2$$

**Question 13****Answer: C***Explanation:*

$$\text{Solve } \det \left( \begin{bmatrix} 3 & 2 & 4 \\ m & 1 & -2 \\ 1 & m & 2 \end{bmatrix} \right) \neq 0 \text{ for } m \text{ in CAS, } m \neq -1 \text{ and } m \neq \frac{1}{2}$$

**Question 14****Answer: E**

Explanation:

$$\vec{a} = \frac{1}{m}(\vec{F}_1 + \vec{F}_2) = \frac{1}{12}(-5\vec{i} + 2\vec{j}) \Rightarrow |\vec{a}| = \frac{1}{12}\sqrt{(-5)^2 + 2^2} = \frac{\sqrt{29}}{12}$$

**Question 15****Answer: A**

Explanation:

By Lammi's Theorem

$$\frac{F_1}{\sin(120^\circ)} = \frac{F_2}{\sin(90^\circ)} = \frac{F_3}{\sin(150^\circ)}$$

Therefore

$$\frac{2\sqrt{3}}{3}F_1 = F_2 = 2F_3$$

**Question 16****Answer: D**

Explanation:

The velocity  $\vec{v} = 30 \cos(15^\circ) \vec{i} + (30 \sin(15^\circ) + 9.8t) \vec{j}$ The vertical height travelled  $h = 30 \sin(15^\circ) t + \frac{1}{2} \times 9.8 t^2 = 2.5 \Rightarrow t \approx 0.274 \text{ s}$ 

Therefore, the travelled distance

$$L = \int_0^{0.2744} \left| 30 \cos(15^\circ) \vec{i} + (30 \sin(15^\circ) + 9.8t) \vec{j} \right| dt \approx 8.34 \text{ m}$$

$$\text{solve}\left(30 \cdot \sin(15^\circ) \cdot t + \frac{1}{2} \cdot 9.8 \cdot t^2 = 2.5, t\right) \quad t = -1.85905 \text{ or } t = 0.274443$$

$$\int_0^{0.274443} \text{norm}\left(\begin{bmatrix} 30 \cdot \cos(15^\circ) & 30 \cdot \sin(15^\circ) + 9.8 \cdot t \end{bmatrix}\right) dt \quad 8.33891$$

**Question 17****Answer: B***Explanation:*

$$v \frac{dv}{dx} = \frac{v(2x+1)}{(x+1)(3x+2)}$$

$$v(10) - v(2) = \int_2^{10} \frac{2x+1}{(3x+2)(x+1)} dx$$

$$v(10) = \int_2^{10} \left( \frac{-1}{3x+2} - \frac{-1}{x+1} \right) dx + v(2) = \int_2^{10} \left( \frac{1}{x+1} - \frac{1}{3x+2} \right) dx + 5$$

**Question 18****Answer:***Explanation:*

$$mean = 5 \times 430 + 10 \times 250 = 4650$$

$$std = \sqrt{5 \times 25^2 + 10 \times 10^2} = 5\sqrt{165}$$

**Question 19****Answer: E***Explanation:* $X \sim N(\mu, 5.2^2)$ ,  $n = 64$ ,  $\bar{x} = 20.5$ , using CAS

z Interval

$\sigma$ : 5.2

$\bar{x}$ : 20.5

n: 64

C Level: 0.95

OK Cancel

zInterval 5.2,20.5,64,0.95: *stat.results*

|               |              |
|---------------|--------------|
| "Title"       | "z Interval" |
| "CLower"      | 19.226       |
| "CUpper"      | 21.774       |
| " $\bar{x}$ " | 20.5         |
| "ME"          | 1.27398      |
| "n"           | 64.          |
| " $\sigma$ "  | 5.2          |

**Question 20****Answer: B***Explanation:*

Enter the statistics into CAS

z Test

$\mu_0$ : 200

$\sigma$ : 30

$\bar{x}$ : 211.6

n: 36

Alternate Hyp:  $H_a: \mu > \mu_0$

OK Cancel

zTest 200,30,211.6,36,1: *stat.results*

|                 |                   |
|-----------------|-------------------|
| "Title"         | "z Test"          |
| "Alternate Hyp" | " $\mu > \mu_0$ " |
| "z"             | 2.32              |
| "PVal"          | 0.01017           |
| " $\bar{x}$ "   | 211.6             |
| "n"             | 36.               |
| " $\sigma$ "    | 30.               |

$p \approx 0.0102$  is less than 0.05, hence enough evidence to reject  $H_0$ .

**SECTION B: Extended Response questions****Question 1 (10 marks)**

a.  $f'(x) = \frac{4x^3 - 16x^2 + 16x + 3}{(x-2)^2}$

Solve  $f'(x) = 0$  for  $x$ ,  $x \approx -0.1607$ .

$$f(-0.1607) \approx 2.440$$

The required stationary point is  $(-0.16, 2.44)$ .

1 mark

b. Solve  $f''(x) = 0$  for  $x$  by CAS,  $x \approx 3.1447$


1 mark

$$f(3.1447) \approx 18.1577$$

The required point of inflection is  $(3.14, 18.16)$

1 mark

c. Graph



Labelling the turning point and the point of inflection

1 mark

Labelling the end points

1 mark

Correct graph for  $x \in [-3, 2) \cup (2, 5]$

1 mark

d. i.  $L = \int_{-3}^1 \sqrt{1 + (f'(x))^2} dx = \int_{-3}^1 \sqrt{1 + \left(\frac{4x^3 - 16x^2 + 16x + 3}{(x-2)^2}\right)^2} dx$

1 mark

ii.  $\approx 21.44 \text{ units}$

1 mark

e.  $V = \pi \int_{-3}^1 (f(x))^2 dx$

1 mark

$$\approx 888 \text{ units}^3$$

1 mark

**Question 2 (11 marks)**

- a.** Solve  $\left| x + yi + \frac{9}{2} + \left( \frac{7\sqrt{3}}{2} - 1 \right) i \right| = |x + yi - (\sqrt{3} + 1)i|$  for  $y$  in CAS. 1 mark

$$\text{solve} \left( \left| x + y \cdot i + \frac{9}{2} + \left( \frac{7 \cdot \sqrt{3}}{2} - 1 \right) \cdot i \right| = |x + y \cdot i - (\sqrt{3} + 1) \cdot i|, y \right)$$

$$y = \frac{-\sqrt{3} \cdot (x - \sqrt{3} + 6)}{3}$$

$$\text{expand} \left( \frac{-\sqrt{3} \cdot (x - \sqrt{3} + 6)}{3} \right)$$

$$\frac{-\sqrt{3} \cdot x}{3} - 2 \cdot \sqrt{3} + 1$$

Therefore,  $y = \frac{-\sqrt{3}}{3}x - 2\sqrt{3} + 1$  1 mark


- b.** Solve  $\left| z + \frac{9}{2} + \left( \frac{7\sqrt{3}}{2} - 1 \right) i \right| = |z - (\sqrt{3} + 1)i|$  and  $|z + 1 - i| = 5$  simultaneously in CAS. 1 mark

$$\text{solve} \left( \left( \left| x + y \cdot i + \frac{9}{2} + \left( \frac{7 \cdot \sqrt{3}}{2} - 1 \right) \cdot i \right| = |x + y \cdot i - (\sqrt{3} + 1) \cdot i|, x, y \right), \left| x + y \cdot i + 1 - i \right| = 5 \right)$$

$$x = -6 \text{ and } y = 1 \text{ or } x = \frac{3}{2} \text{ and } y = \frac{-(5 \cdot \sqrt{3} - 2)}{2}$$

The intersections are  $B(-6, 1)$  and  $C(\frac{3}{2}, 1 - \frac{5\sqrt{3}}{2})$ . 1 mark

- c.** Graph



Correct graphs for the circle and the line  
 Labelling the intersections

1 mark  
 1 mark

d.  $\angle ABC = \arctan\left(\frac{\sqrt{3}}{3}\right) = 30^\circ \Rightarrow$   
 $\angle BAC = 180^\circ - 2 \times 30^\circ = 120^\circ$

1 mark  
 1 mark

e. The area of the minor sector  $A_1 = \frac{1}{3} \times \pi r^2 = \frac{25\pi}{3}$

1 mark

The area of the triangle ABC  $A_2 = \frac{1}{2} \times (-1 + 6) \times \left(1 - \left(1 - \frac{5\sqrt{3}}{2}\right)\right) = \frac{25\sqrt{3}}{4}$

1 mark

Alternative:  $A_2 = \frac{1}{2} AB \times AC \times \sin(120^\circ) = \frac{25\sqrt{3}}{4}$

Therefore, the area of the minor segment

$$A = \frac{25\pi}{3} - \frac{25\sqrt{3}}{4}$$

1 mark

### Question 3 (9 marks)

a. i. Solve  $\frac{dp}{dt} = r p \left(1 - \frac{p}{k}\right)$  by CAS,  
 1 mark

$$\text{deSolve}\left(p' = r \cdot p \cdot \left(1 - \frac{p}{k}\right), t, p\right) \quad p = \frac{k \cdot e^{r \cdot t}}{e^{r \cdot t} + k \cdot c1}$$

Therefore  $p = \frac{k e^{rt}}{e^{rt} + kc}$  or  $\frac{k e^{rt}}{e^{rt} + A}$ , where  $c$  and  $A$  are constants.

1 mark

ii.  $5000 = \lim_{t \rightarrow \infty} \frac{k e^{rt}}{e^{rt} + kc} = \lim_{t \rightarrow \infty} \frac{k}{1 + kc e^{-rt}} = k$   
 $k = 5000$

1 mark

- b. From Parts a i) and a ii),  $P(t) = \frac{5000 \times e^{rt}}{e^{rt} + 5000c}$   
Solve  $P(0) = 2500$  and  $P(10) = 2600$  simultaneously by CAS,

$$c = \frac{1}{5000}, \quad r = \frac{1}{10} \ln\left(\frac{13}{12}\right) \quad 1 \text{ mark}$$

Therefore

$$P(t) = \frac{5000 \times e^{\ln\left(\frac{13}{12}\right) \frac{t}{10}}}{e^{\ln\left(\frac{13}{12}\right) \frac{t}{10}} + 5000 \times \frac{1}{5000}} \quad 1 \text{ mark}$$

$$= \frac{5000 \times \left(\frac{13}{12}\right)^{\frac{t}{10}}}{1 + \left(\frac{13}{12}\right)^{\frac{t}{10}}} \quad 1 \text{ mark}$$

- c. Let  $u = \frac{dp}{dt} = \frac{1}{10} \ln\left(\frac{13}{12}\right) p \left(1 - \frac{p}{5000}\right)$

$$\frac{d^2p}{dt^2} = \frac{du}{dt} = \frac{du}{dp} \times \frac{dp}{dt} \quad 1 \text{ mark}$$

$$= \frac{1}{10} \ln\left(\frac{13}{12}\right) \left(1 - \frac{p}{2500}\right) \times \frac{1}{10} \ln\left(\frac{13}{12}\right) p \left(1 - \frac{p}{5000}\right) \quad 1 \text{ mark}$$

$$= \left(\frac{1}{10} \ln\left(\frac{13}{12}\right)\right)^2 p \left(1 - \frac{p}{2500}\right) \left(1 - \frac{p}{5000}\right) \quad 1 \text{ mark}$$

#### Question 4 (11 marks)

- a. Solve  $r_A(s) = r_B(t)$  for  $s, t$  by CAS,

At  $s \approx 0.37$  Particle A passes (3.72, 2.00) 2 mark

At  $t = 1.11$  Particle B passes (3.72, 2.00) 1 mark


Define  $ra(t) = \begin{bmatrix} 5 \cdot \cos(2 \cdot t) \\ 3 \cdot \sin(2 \cdot t) \end{bmatrix}$  Done

Define  $rb(t) = \begin{bmatrix} -3 + 3 \cdot \sec(t) \\ \tan(t) \end{bmatrix}$  Done

 solve( $ra(s) = rb(t), s, t$ ) |  $0 \leq s \leq \frac{5 \cdot \pi}{12}$  and  $0 \leq t \leq \frac{5 \cdot \pi}{12}$   $s = 0.365859$  and  $t = 1.10804$

$ra(0.365859)$   $\begin{bmatrix} 3.72014 \\ 2.00445 \end{bmatrix}$

b. Graph



Correct end points

1 mark

Correct directions

1 mark

Correct shapes

1 mark

c.  $\tilde{r}_A(t) = -10 \sin(2t) \tilde{i} + 6 \cos(2t) \tilde{j}$

$\tilde{r}_B(t) = 3 \sec(t) \tan(t) \tilde{i} + \sec^2(t) \tilde{j}$

1 mark

Let  $\theta$  be the angle between the moving directions when they cross the same point.

$$\cos(\theta) = \frac{\tilde{r}_A'(0.366) \cdot \tilde{r}_B'(1.108)}{|\tilde{r}_A'(0.366)| |\tilde{r}_B'(1.108)|} = -0.5855$$

$$\theta \approx 126^\circ$$

1 mark

d. i. Let  $h(t) = |\tilde{r}_A(t) - \tilde{r}_B(t)|$

1 mark

Solve  $\frac{d}{dt}(h(t)) = 0$  for  $t$ ,  $t \approx 0.703$  s

1 mark

ii.  $h(0.7035) \approx 2.11$  m

1 mark

**Question 5 (11marks)****a. Proof**

$$ma = -mg - 0.025mv^2$$

$$40a = -40g - v^2$$

1 mark

$$40 \frac{dv}{dt} = 40 v \frac{dv}{dx} = -40g - v^2$$

$$\frac{dv}{dx} = -\frac{40g+v^2}{40v}$$

1 mark

**b. Solve  $x' = -\frac{40v}{40g+v^2}$  in CAS,  $x = c - 20 \ln(v^2 + 40g)$** 

1 mark

$$\text{deSolve}\left(x' = \frac{-40 \cdot v}{40 \cdot g + v^2}, v, x\right)$$

$$x = c - 20 \cdot \ln(v^2 + 40 \cdot g)$$

Substitute  $x = 0, v = 80$ ,

$$0 = c - 20 \ln(80^2 + 40g) \Rightarrow c = 20 \ln(6400 + 40g)$$

$$x = 20 \ln\left(\frac{6400 + 40g}{v^2 + 40g}\right)$$

1 mark

**c. At the highest point  $v = 0$ .**

1 mark

$$x = 20 \ln\left(\frac{6400 + 40 \times 9.8}{0^2 + 40 \times 9.8}\right) \approx 57.04 \text{ m}$$

1mark

**d. Solve the differential equation  $y' = \frac{1000v}{1000 \times 9.8 - v^2}$  in CAS**

$$y = c - 500 \ln |v^2 - 9800|$$

1 mark

Substitute  $v = 0, y = 0$ ,

$$c = 500 \ln(9800)$$

1 mark

$$y = 500 \ln \left| \frac{9800}{9800 - v^2} \right|$$

$$\text{Solve } 120 = 500 \ln \left| \frac{9800}{9800 - v^2} \right| \quad \text{for } v = 45.728 \text{ ms}^{-1}$$

1 mark

$$\text{e.} \quad a = \frac{dv}{dt} = g - 0.001v^2$$

1 mark

$$t = \int_0^{45.728} \frac{1}{9.8 - 0.001v^2} dt \approx 5.05 \text{ s}$$

1 mark

**Question 6 (8 marks)**

- a. Let  $Y = X_1 + X_2 + \dots + X_{25}$ , where  $X_i \sim N(200, 5^2)$ .

Therefore  $E(Y) = 25 \times 200 = 5000$ ,  $Var(Y) = 25 \times 5^2 = 625$

1 mark

$$\Pr(Y > 5050) = 0.02275$$

1 mark

- b. Let  $\bar{X}$  be the sample mean of the amount of the liquid. Then  $\bar{X} \sim N(200, \left(\frac{5}{\sqrt{25}}\right)^2)$ .

1 mark

$$\Pr(\bar{X} > 201) = 0.1587$$

1 mark

- c. Find from CAS

zInterval 5,197.5,25,0.95: stat.results

|          |              |
|----------|--------------|
| "Title"  | "z Interval" |
| "CLower" | 195.54       |
| "CUpper" | 199.46       |
| "x"      | 197.5        |
| "ME"     | 1.95996      |
| "n"      | 25.          |
| "s"      | 5.           |

(195.54, 199.46)

- d.  $H_0: \mu = 200$ ,  $H_1: \mu < 200$

1 mark

- e.  $p = \Pr(\bar{X} < 197.5 | \mu = 200) = 0.00621$

zTest 200,5,197.5,25,-1: stat.results

|                 |                   |
|-----------------|-------------------|
| "Title"         | "z Test"          |
| "Alternate Hyp" | " $\mu < \mu_0$ " |
| "z"             | -2.5              |
| "PVal"          | 0.00621           |
| "x"             | 197.5             |
| "n"             | 25.               |
| "s"             | 5.                |

Conclusion: There is sufficient evidence to reject  $H_0$ .