

# **Year 2018**

## **VCE**

### **Specialist Mathematics**

### **Trial Examination 1**



KILBAHA MULTIMEDIA PUBLISHING  
PO BOX 2227  
KEW VIC 3101  
AUSTRALIA

TEL: (03) 9018 5376  
FAX: (03) 9817 4334  
kilbaha@gmail.com  
<http://kilbaha.com.au>

## IMPORTANT COPYRIGHT NOTICE

- This material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Multimedia Publishing.
- The contents of this work are copyrighted. Unauthorised copying of any part of this work is illegal and detrimental to the interests of the author.
- For authorised copying within Australia please check that your institution has a licence from **Copyright Agency Limited**. This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.

Reproduction and communication for educational purposes The Australian Copyright Act 1968 (the Act) allows a maximum of one chapter or 10% of the pages of this work, to be reproduced and/or communicated by any educational institution for its educational purposes provided that educational institution (or the body that administers it) has given a remuneration notice to Copyright Agency Limited (CAL) under the Act.

For details of the CAL licence for educational institutions contact  
CAL, Level 15, 233 Castlereagh Street, Sydney, NSW, 2000

Tel: (02) 9394 7600

Fax: (02) 9394 7601

Email: [info@copyright.com.au](mailto:info@copyright.com.au)

Web: <http://www.copyright.com.au>

- While every care has been taken, no guarantee is given that these questions are free from error. Please contact us if you believe you have found an error.

# Victorian Certificate of Education 2018

## STUDENT NUMBER

Figures  
Words

|  |  |
|--|--|
|  |  |
|  |  |

|  |  |  |  |  |  |
|--|--|--|--|--|--|
|  |  |  |  |  |  |
|  |  |  |  |  |  |

Letter

|  |
|--|
|  |
|--|

## SPECIALIST MATHEMATICS

### Trial Written Examination 1

Reading time: 15 minutes

Total writing time: 1 hour

## QUESTION AND ANSWER BOOK

### Structure of book

| <i>Number of questions</i> | <i>Number of questions to be answered</i> | <i>Number of marks</i> |
|----------------------------|-------------------------------------------|------------------------|
| 10                         | 10                                        | 40                     |

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or white out liquid/tape.

### Materials supplied

- Question and answer book of 20 pages with a detachable sheet of miscellaneous formulas at the end of this booklet.

### Instructions

- Detach the formula sheet from the end of this book during reading time.
- Write your **student number** in the space provided above on this page.
- All written responses must be in English.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.**



**Question 2** (4 marks)

Let  $f(x) = \sqrt{\arcsin\left(\frac{3x}{4}\right)}$

**a.** State the maximal domain and the range of the function  $f$ .

2 marks

---



---



---



---



---

**b.** Find  $f'(x)$  and hence evaluate  $\int_0^{\frac{4}{3}} \frac{1}{\sqrt{(16-9x^2)\sin^{-1}\left(\frac{3x}{4}\right)}} dx$ ,

giving your answer in the form  $\frac{\sqrt{b}\pi}{c}$  where  $b$  and  $c$  are positive integers.

2 marks

---



---



---



---



---



---



---



---



---



---



---



---



---



---



---




---



**Question 4** (6 marks)

The diagram shows the graph of the relation  $y^2 = x^3 + 4x^2$ .



- a. Write down a definite integral which gives the total area of the loop in the second and third quadrants.

1 mark

---

---

---

---







**Question 5** (4 marks)

The weights of medium sized dogs are normally distributed with a mean of 10 kg and a standard deviation of 2.5 kg. The weights of cats are normally distributed with a mean of 5 kg and a standard deviation of 1.5 kg. Assume that the weights of medium sized dogs and cats are independent random variables.

- a.** Find the mean and variance of a medium sized dog and two cats.

1 mark

---

---

---

---

- b.** A dog breeder is concerned that his medium sized dogs are over-weight. A sample of 25 dogs is found to have a mean weight of 11 kg. Assume that the population standard deviation for medium sized dogs is still 2.5 kg

- i.** State the appropriate null and alternative hypotheses for the weights of medium sized dogs in this situation.

1 mark

---

---

---

- ii.** The  $p$  value for this test is given by  $\Pr(Z \geq a)$ , where  $Z$  has the standard normal distribution. Find the value of  $a$  and hence determine whether the null hypothesis should be rejected at the 0.05 level of significance.

2 marks

---

---

---

---

---



**Question 7** (4 marks)

A particle moves so that its velocity vector is  $\underline{v}(t) = -2e^{-t}\underline{i} + 2e^{2t}\underline{j} \text{ ms}^{-1}$  at time  $t$  seconds, where  $t \geq 0$ .

- a. Given that  $\underline{r}(0) = 2\underline{i}$ , find the position vector  $\underline{r}(t)$  and hence show that the particle moves on the part of the curve  $y = \frac{4}{x^2} - 1$ , stating the domain and range.

3 marks

---



---



---



---



---



---



---



---




---



---


- b. On the diagram below, sketch the graph of the path of the particle, indicating the direction of motion.

1 mark



**Question 8** (3 marks)

The diagram shows part of the graph of  $y = 2\sec\left(\frac{x}{3}\right)$ .



The shaded region  $R$  is the area bounded by the graph of  $y = 2\sec\left(\frac{x}{3}\right)$ , the  $y$ -axis and the line  $y = 4$ . Find the volume generated when the region  $R$  is rotated about the  $x$ -axis.

---

---

---

---

---

---

---

---

---





[illegible]



# **SPECIALIST MATHEMATICS**

## **Written examination 1**

### **FORMULA SHEET**

#### **Directions to students**

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

## Specialist Mathematics formulas

### Mensuration

|                                   |                                                             |
|-----------------------------------|-------------------------------------------------------------|
| area of a trapezium               | $\frac{1}{2}(a+b)h$                                         |
| curved surface area of a cylinder | $2\pi rh$                                                   |
| volume of a cylinder              | $\pi r^2 h$                                                 |
| volume of a cone                  | $\frac{1}{3}\pi r^2 h$                                      |
| volume of a sphere                | $\frac{4}{3}\pi r^3$                                        |
| volume of a pyramid               | $\frac{1}{3}Ah$                                             |
| area of triangle                  | $\frac{1}{2}bc \sin(A)$                                     |
| sine rule                         | $\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$ |
| cosine rule                       | $c^2 = a^2 + b^2 - 2ab \cos(C)$                             |

### Circular (trigonometric) functions

|                                                                      |                                                            |
|----------------------------------------------------------------------|------------------------------------------------------------|
| $\cos^2(x) + \sin^2(x) = 1$                                          |                                                            |
| $1 + \tan^2(x) = \sec^2(x)$                                          | $\cot^2(x) + 1 = \operatorname{cosec}^2(x)$                |
| $\sin(x+y) = \sin(x)\cos(y) + \cos(x)\sin(y)$                        | $\sin(x-y) = \sin(x)\cos(y) - \cos(x)\sin(y)$              |
| $\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$                        | $\cos(x-y) = \cos(x)\cos(y) + \sin(x)\sin(y)$              |
| $\tan(x+y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$           | $\tan(x-y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$ |
| $\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$ |                                                            |
| $\sin(2x) = 2\sin(x)\cos(x)$                                         | $\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$                |

## Circular (trigonometric) functions - continued

| Function | $\sin^{-1}$ (arcsin)                         | $\cos^{-1}$ (arccos) | $\tan^{-1}$ (arctan)                         |
|----------|----------------------------------------------|----------------------|----------------------------------------------|
| Domain   | $[-1, 1]$                                    | $[-1, 1]$            | $R$                                          |
| Range    | $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ | $[0, \pi]$           | $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ |

## Algebra ( complex numbers )

|                                                                                |                                                                             |
|--------------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| $z = x + yi = r(\cos(\theta) + i \sin(\theta)) = r \operatorname{cis}(\theta)$ |                                                                             |
| $ z  = \sqrt{x^2 + y^2} = r$                                                   | $-\pi < \operatorname{Arg}(z) \leq \pi$                                     |
| $z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$                    | $\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$ |
| $z^n = r^n \operatorname{cis}(n\theta)$ (de Moivre's theorem)                  |                                                                             |

## Probability and statistics

|                                              |                                                                                                                     |
|----------------------------------------------|---------------------------------------------------------------------------------------------------------------------|
| for random variables $X$ and $Y$             | $E(aX + b) = aE(X) + b$<br>$E(aX + bY) = aE(X) + bE(Y)$<br>$\operatorname{Var}(aX + b) = a^2 \operatorname{Var}(X)$ |
| for independent random variables $X$ and $Y$ | $\operatorname{Var}(aX + bY) = a^2 \operatorname{Var}(X) + b^2 \operatorname{Var}(Y)$                               |
| approximate confidence interval for $\mu$    | $\left( \bar{x} - z \frac{s}{\sqrt{n}}, \bar{x} + z \frac{s}{\sqrt{n}} \right)$                                     |
| distribution of sample mean $\bar{X}$        | mean $E(\bar{X}) = \mu$ variance $\operatorname{Var}(\bar{X}) = \frac{\sigma^2}{n}$                                 |

## Vectors in two and three dimensions

|                                                                                                                        |
|------------------------------------------------------------------------------------------------------------------------|
| $\underline{r} = x\hat{i} + y\hat{j} + z\hat{k}$                                                                       |
| $ \underline{r}  = \sqrt{x^2 + y^2 + z^2} = r$                                                                         |
| $\dot{\underline{r}} = \frac{d\underline{r}}{dt} = \frac{dx}{dt}\hat{i} + \frac{dy}{dt}\hat{j} + \frac{dz}{dt}\hat{k}$ |
| $\underline{r}_1 \cdot \underline{r}_2 = r_1 r_2 \cos(\theta) = x_1 x_2 + y_1 y_2 + z_1 z_2$                           |

## Mechanics

|                    |                                  |
|--------------------|----------------------------------|
| momentum           | $\underline{p} = m\underline{v}$ |
| equation of motion | $\underline{R} = m\underline{a}$ |

## Calculus

|                                                        |                                                                                                                 |
|--------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|
| $\frac{d}{dx}(x^n) = nx^{n-1}$                         | $\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$                                                            |
| $\frac{d}{dx}(e^{ax}) = ae^{ax}$                       | $\int e^{ax} dx = \frac{1}{a} e^{ax} + c$                                                                       |
| $\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$                | $\int \frac{1}{x} dx = \log_e x  + c$                                                                           |
| $\frac{d}{dx}(\sin(ax)) = a \cos(ax)$                  | $\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$                                                                  |
| $\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$                 | $\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$                                                                   |
| $\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$                | $\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$                                                                 |
| $\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$  | $\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$                               |
| $\frac{d}{dx}(\cos^{-1}(x)) = \frac{-1}{\sqrt{1-x^2}}$ | $\int \frac{-1}{\sqrt{a^2-x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$                              |
| $\frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{1+x^2}$         | $\int \frac{a}{a^2+x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$                                             |
|                                                        | $\int (ax+b)^n dx = \frac{1}{a(n+1)} (ax+b)^{n+1} + c, n \neq -1$                                               |
|                                                        | $\int (ax+b)^{-1} dx = \frac{1}{a} \log_e ax+b  + c$                                                            |
| product rule                                           | $\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$                                                          |
| quotient rule                                          | $\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$                          |
| chain rule                                             | $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$                                                                   |
| Euler's method                                         | If $\frac{dy}{dx} = f(x)$ , $x_0 = a$ and $y_0 = b$ , then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + hf(x_n)$    |
| acceleration                                           | $a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$             |
| arc length                                             | $\int_{x_1}^{x_2} \sqrt{1+(f'(x))^2} dx \quad \text{or} \quad \int_{t_1}^{t_2} \sqrt{(x'(t))^2 + (y'(t))^2} dt$ |

**END OF FORMULA SHEET**