

Victorian Certificate of Education
2018

SUPERVISOR TO ATTACH PROCESSING LABEL HERE

STUDENT NUMBER

									Letter
--	--	--	--	--	--	--	--	--	--------

SPECIALIST MATHEMATICS
Written examination 1

Tuesday 5 June 2018

Reading time: 2.00 pm to 2.15 pm (15 minutes)

Writing time: 2.15 pm to 3.15 pm (1 hour)

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
9	9	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or correction fluid/tape.

Materials supplied

- Question and answer book of 11 pages
- Formula sheet
- Working space is provided throughout the book.

Instructions

- Write your **student number** in the space provided above on this page.
- Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.
- All written responses must be in English.

At the end of the examination

- You may keep the formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

THIS PAGE IS BLANK

DO NOT WRITE IN THIS AREA

Instructions

Answer **all** questions in the spaces provided.

Unless otherwise specified, an **exact** answer is required to a question.


In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Take the **acceleration due to gravity** to have magnitude $g \text{ ms}^{-2}$, where $g = 9.8$

Question 1 (3 marks)

A light inextensible string hangs over a frictionless pulley connecting masses of 3 kg and 7 kg, as shown below.



- a. Draw all of the forces acting on the two masses on the diagram above. 1 mark
- b. Calculate the tension in the string. 2 marks

Question 2 (3 marks)

Let $\underline{a} = 3\underline{i} - 2\underline{j} + m\underline{k}$ and $\underline{b} = 2\underline{i} - \underline{j} + 3\underline{k}$, where $m \in \mathbb{R}$.

Find the value(s) of m such that the magnitude of the vector resolute of $\underline{a}$ parallel to $\underline{b}$ is equal to $\sqrt{14}$.

Question 3 (3 marks)

Find $\sin(t)$ given that $t = \arccos\left(\frac{12}{13}\right) + \arctan\left(\frac{3}{4}\right)$.

Question 4 (4 marks)

Throughout this question, use an integer multiple of standard deviations in calculations.

The standard deviation of all scores on a particular test is 21.0

- a. From the results of a random sample of n students, a 95% confidence interval for the mean score for all students was calculated to be (44.7, 51.7).

Calculate the mean score and the size of this random sample.

2 marks

- b. Determine the size of another random sample for which the endpoints of the 95% confidence interval for the population mean of the particular test would be 1.0 either side of the sample mean.

2 marks

Question 5 (4 marks)

Evaluate $\int_1^{2\sqrt{3}-1} \left(\frac{1}{x^2 + 2x + 5} \right) dx$.

Question 6 (4 marks)

Given that $y = (x - 1)e^{2x}$ is a solution to the differential equation $a \frac{d^2 y}{dx^2} + b \frac{dy}{dx} = y$, find the values of a and b , where a and b are real constants.

Question 7 (4 marks)

a. Find $\frac{d}{dx} \left((1-x^2)^{\frac{1}{2}} \right)$.

2 marks

b. Hence, find the length of the curve specified by $y = \sqrt{1-x^2}$ from $x = \frac{1}{2}$ to $x = \frac{\sqrt{3}}{2}$. Give your answer in the form $k\pi$, $k \in R$.

2 marks


TURN OVER

Question 8 (6 marks)

A circle in the complex plane is given by the relation $|z - 1 - i| = 2$, $z \in C$.

- a. Sketch the circle on the Argand diagram below.

1 mark



- b. i. Write the equation of the circle in the form $(x - a)^2 + (y - b)^2 = c$ and show that the gradient of a tangent to the circle can be expressed as $\frac{dy}{dx} = \frac{1-x}{y-1}$.

2 marks

- ii. Find the gradient of the tangent to the circle where $x = 2$ in the first quadrant of the complex plane.

1 mark

- c. Find the equations of all rays that are perpendicular to the circle in the form $\text{Arg}(z) = \alpha$. 2 marks

DO NOT WRITE IN THIS AREA

TURN OVER

Question 9 (9 marks)

- a.** **i.** Given that $\cot(2\theta) = a$, show that $\tan^2(\theta) + 2a \tan(\theta) - 1 = 0$.

2 marks

- ii.** Show that $\tan(\theta) = -a \pm \sqrt{a^2 + 1}$.

1 mark

- iii.** Hence, show that $\tan\left(\frac{\pi}{12}\right) = 2 - \sqrt{3}$, given that $\cot(2\theta) = \sqrt{3}$, where $\theta \in (0, \pi)$.

1 mark

- b.** Find the gradient of the tangent to the curve $y = \tan(\theta)$ at $\theta = \frac{\pi}{12}$.

2 marks

- c. A solid of revolution is formed by rotating the region between the graph of $y = \tan(\theta)$, the horizontal axis, and the lines $\theta = \frac{\pi}{12}$ and $\theta = \frac{\pi}{3}$ about the horizontal axis.

Find the volume of the solid of revolution.

3 marks

**Victorian Certificate of Education
2018**

SPECIALIST MATHEMATICS

Written examination 1

FORMULA SHEET

Instructions

This formula sheet is provided for your reference.
A question and answer book is provided with this formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Specialist Mathematics formulas

Mensuration

area of a trapezium	$\frac{1}{2}(a+b)h$
curved surface area of a cylinder	$2\pi rh$
volume of a cylinder	$\pi r^2 h$
volume of a cone	$\frac{1}{3}\pi r^2 h$
volume of a pyramid	$\frac{1}{3}Ah$
volume of a sphere	$\frac{4}{3}\pi r^3$
area of a triangle	$\frac{1}{2}bc \sin(A)$
sine rule	$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$
cosine rule	$c^2 = a^2 + b^2 - 2ab \cos(C)$

Circular functions

$\cos^2(x) + \sin^2(x) = 1$	
$1 + \tan^2(x) = \sec^2(x)$	$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$
$\sin(x+y) = \sin(x)\cos(y) + \cos(x)\sin(y)$	$\sin(x-y) = \sin(x)\cos(y) - \cos(x)\sin(y)$
$\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$	$\cos(x-y) = \cos(x)\cos(y) + \sin(x)\sin(y)$
$\tan(x+y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$	$\tan(x-y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$
$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$	
$\sin(2x) = 2\sin(x)\cos(x)$	$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$

Circular functions – continued

Function	$\sin^{-1}$ or arcsin	$\cos^{-1}$ or arccos	$\tan^{-1}$ or arctan
Domain	$[-1, 1]$	$[-1, 1]$	R
Range	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$[0, \pi]$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Algebra (complex numbers)

$z = x + iy = r(\cos(\theta) + i\sin(\theta)) = r \operatorname{cis}(\theta)$	
$ z = \sqrt{x^2 + y^2} = r$	$-\pi < \operatorname{Arg}(z) \leq \pi$
$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$	$\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$
$z^n = r^n \operatorname{cis}(n\theta)$ (de Moivre's theorem)	

Probability and statistics

for random variables X and Y	$E(aX + b) = aE(X) + b$ $E(aX + bY) = aE(X) + bE(Y)$ $\operatorname{var}(aX + b) = a^2 \operatorname{var}(X)$
for independent random variables X and Y	$\operatorname{var}(aX + bY) = a^2 \operatorname{var}(X) + b^2 \operatorname{var}(Y)$
approximate confidence interval for μ	$\left(\bar{x} - z \frac{s}{\sqrt{n}}, \bar{x} + z \frac{s}{\sqrt{n}} \right)$
distribution of sample mean $\bar{X}$	mean $E(\bar{X}) = \mu$ variance $\operatorname{var}(\bar{X}) = \frac{\sigma^2}{n}$

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e x + c$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$	$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$
$\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$	$\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\cos^{-1}(x)) = \frac{-1}{\sqrt{1-x^2}}$	$\int \frac{-1}{\sqrt{a^2-x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{1+x^2}$	$\int \frac{a}{a^2+x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$
	$\int (ax+b)^n dx = \frac{1}{a(n+1)} (ax+b)^{n+1} + c, n \neq -1$
	$\int (ax+b)^{-1} dx = \frac{1}{a} \log_e ax+b + c$
product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$
quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
Euler's method	If $\frac{dy}{dx} = f(x)$, $x_0 = a$ and $y_0 = b$, then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + hf(x_n)$
acceleration	$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$
arc length	$\int_{x_1}^{x_2} \sqrt{1+(f'(x))^2} dx$ or $\int_{t_1}^{t_2} \sqrt{(x'(t))^2 + (y'(t))^2} dt$

Vectors in two and three dimensions

$\underline{r} = x\hat{i} + y\hat{j} + z\hat{k}$
$ \underline{r} = \sqrt{x^2 + y^2 + z^2} = r$
$\dot{\underline{r}} = \frac{d\underline{r}}{dt} = \frac{dx}{dt}\hat{i} + \frac{dy}{dt}\hat{j} + \frac{dz}{dt}\hat{k}$
$\underline{r}_1 \cdot \underline{r}_2 = r_1 r_2 \cos(\theta) = x_1 x_2 + y_1 y_2 + z_1 z_2$

Mechanics

momentum	$\underline{p} = m\underline{v}$
equation of motion	$\underline{R} = m\underline{a}$