

Trial Examination 2020

VCE Specialist Mathematics Units 3&4

Written Examination 1

Question and Answer Booklet

Reading time: 15 minutes

Writing time: 1 hour

Student's Name: _____

Teacher's Name: _____

Structure of booklet

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
10	10	40

Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers.

Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or correction fluid/tape.

Materials supplied

Question and answer booklet of 12 pages

Formula sheet

Working space is provided throughout the booklet.

Instructions

Write your **name** and your **teacher's name** in the space provided above on this page.

Unless otherwise indicated, the diagrams in this booklet are **not** drawn to scale.

All written responses must be in English.

At the end of the examination

You may keep the formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Students are advised that this is a trial examination only and cannot in any way guarantee the content or the format of the 2020 VCE Specialist Mathematics Units 3&4 Written Examination 1.

Neap Education (Neap) Trial Exams are licensed to be photocopied or placed on the school intranet and used only within the confines of the school purchasing them, for the purpose of examining that school's students only. They may not be otherwise reproduced or distributed. The copyright of Neap Trial Exams remains with Neap. No Neap Trial Exam or any part thereof is to be issued or passed on by any person to any party inclusive of other schools, non-practising teachers, coaching colleges, tutors, parents, students, publishing agencies or websites without the express written consent of Neap.

Instructions

Answer **all** questions in the spaces provided.

Unless otherwise specified, an **exact** answer is required to a question.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this booklet are **not** drawn to scale.

Take the **acceleration due to gravity** to have magnitude $g \text{ ms}^{-2}$, where $g = 9.8$.

Question 1 (2 marks)

Consider the function $f(x) = \arctan\left(\frac{2}{x}\right)$, where $x \neq 0$.

Find an expression for $f'(x)$. Give your answer in the form $\frac{a}{x^2 + b}$, where $a, b \in \mathbb{Z}$.

Question 2 (3 marks)

Consider the equation $z^2 - (1 - 2i)z + 1 + 5i = 0$. One of the roots of the equation is $-1 + i$.

- a. State why the conjugate root theorem **cannot** be applied to find the other root of the equation.


1 mark

- b. Find the other root of the equation, expressing your answer in cartesian form.

2 marks

Question 3 (4 marks)

Sketch the graph of $y = \frac{x^2 - x + 2}{x - 2}$ on the axes below. Clearly label any intercepts and stationary points with their the coordinates, and any asymptotes with their equations.



Question 4 (3 marks)

Find the set of values of x for which $|2x + 1| < |x|$.

Question 5 (5 marks)

Relative to a fixed origin O , the point A has position vector $\underline{a} = 8\underline{i} - 6\underline{j} + 5\underline{k}$ and the point B has position vector $\underline{b} = t\underline{i} + t\underline{j} + 2t\underline{k}$, where $t \in \mathbb{R}$, $t \geq 0$.

- a. Show that $|\overrightarrow{AB}| = \sqrt{6t^2 - 24t + 125}$. 2 marks

- b. Find the minimum distance between points A and B . 3 marks

Question 7 (4 marks)

A curve C has the equation $y^2 = 2x - (x + y)^2$.

Find the equations of the tangents to C that are parallel to the x -axis.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 9 (4 marks)

Solve the equation $\arctan(2x) + \arctan(x) = \arctan(3)$, $x \in R$.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 10 (7 marks)

A particle P moves such that at time t its position vector $\underline{r}(t)$ from a fixed origin is given by

$\underline{r}(t) = e^t \cos(t)\underline{i} + e^t \sin(t)\underline{j}$, $0 \leq t \leq \pi$. The velocity vector of particle P is denoted by $\dot{\underline{r}}(t)$.

- a. Determine the particle's initial velocity.

3 marks

- b. Show that $\underline{r}(t)$ always makes an angle of $\frac{\pi}{4}$ with $\dot{\underline{r}}(t)$.

4 marks

END OF QUESTION AND ANSWER BOOKLET

Trial Examination 2020

VCE Specialist Mathematics Units 3&4

Written Examinations 1 and 2

Formula Sheet

Instructions

This formula sheet is provided for your reference.
A question and answer booklet is provided with this formula sheet.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

SPECIALIST MATHEMATICS FORMULAS**Mensuration**

area of a trapezium	$\frac{1}{2}(a + b)h$
curved surface area of a cylinder	$2\pi rh$
volume of a cylinder	$\pi r^2 h$
volume of a cone	$\frac{1}{3}\pi r^2 h$
volume of a pyramid	$\frac{1}{3}Ah$
volume of a sphere	$\frac{4}{3}\pi r^3$
area of a triangle	$\frac{1}{2}bc\sin(A)$
sine rule	$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$
cosine rule	$c^2 = a^2 + b^2 - 2ab\cos(C)$

Circular functions

$\cos^2(x) + \sin^2(x) = 1$	
$1 + \tan^2(x) = \sec^2(x)$	$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$
$\sin(x + y) = \sin(x)\cos(y) + \cos(x)\sin(y)$	$\sin(x - y) = \sin(x)\cos(y) - \cos(x)\sin(y)$
$\cos(x + y) = \cos(x)\cos(y) - \sin(x)\sin(y)$	$\cos(x - y) = \cos(x)\cos(y) + \sin(x)\sin(y)$
$\tan(x + y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$	$\tan(x - y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$
$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$	
$\sin(2x) = 2\sin(x)\cos(x)$	$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$

Circular functions – continued

Function	$\sin^{-1}$ or arcsin	$\cos^{-1}$ or arccos	$\tan^{-1}$ or arctan
Domain	$[-1, 1]$	$[-1, 1]$	R
Range	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$	$[0, \pi]$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$

Algebra (complex numbers)

$z = x + iy = r(\cos(\theta) + i\sin(\theta)) = r\text{cis}(\theta)$	
$ z = \sqrt{x^2 + y^2} = r$	$-\pi < \text{Arg}(z) \leq \pi$
$z_1 z_2 = r_1 r_2 \text{cis}(\theta_1 + \theta_2)$	$\frac{z_1}{z_2} = \frac{r_1}{r_2} \text{cis}(\theta_1 - \theta_2)$
$z^n = r^n \text{cis}(n\theta)$ (de Moivre's theorem)	

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1}x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a}e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e x + c$
$\frac{d}{dx}(\sin(ax)) = a\cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a}\cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a\sin(ax)$	$\int \cos(ax) dx = \frac{1}{a}\sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = a\sec^2(ax)$	$\int \sec^2(ax) dx = \frac{1}{a}\tan(ax) + c$
$\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$	$\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\cos^{-1}(x)) = \frac{-1}{\sqrt{1-x^2}}$	$\int \frac{-1}{\sqrt{a^2-x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$

Calculus – continued

$\frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{1+x^2}$	$\int \frac{a}{a^2+x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$
	$\int (ax+b)^n dx = \frac{1}{a(n+1)}(ax+b)^{n+1} + c, n \neq -1$
	$\int (ax+b)^{-1} dx = \frac{1}{a} \log_e ax+b + c$
product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$
quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
Euler's method	If $\frac{dy}{dx} = f(x)$, $x_0 = a$ and $y_0 = b$, then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + hf(x_n)$
acceleration	$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$
arc length	$\int_{x_1}^{x_2} \sqrt{1+(f'(x))^2} dx$ or $\int_{t_1}^{t_2} \sqrt{(x'(t))^2 + (y'(t))^2} dt$

Vectors in two and three dimensions

$\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$
$ \mathbf{r} = \sqrt{x^2 + y^2 + z^2} = r$
$\dot{\mathbf{r}} = \frac{d\mathbf{r}}{dt} = \frac{dx}{dt}\mathbf{i} + \frac{dy}{dt}\mathbf{j} + \frac{dz}{dt}\mathbf{k}$
$\mathbf{r}_1 \cdot \mathbf{r}_2 = r_1 r_2 \cos(\theta) = x_1 x_2 + y_1 y_2 + z_1 z_2$

Mechanics

momentum	$\mathbf{p} = m\mathbf{v}$
equation of motion	$\mathbf{R} = m\mathbf{a}$

END OF FORMULA SHEET