



P.O. Box 1180
Surrey Hills North VIC 3127
Phone 03 9836 5021

info@theheffernangroup.com.au
www.theheffernangroup.com.au

Student Name:.....

SPECIALIST MATHEMATICS UNITS 3 & 4

TRIAL EXAMINATION 1

2023

Reading Time: 15 minutes

Writing time: 1 hour

Instructions to students

This exam consists of 10 questions.
All questions should be answered in the spaces provided.
There is a total of 40 marks available.
The marks allocated to each of the questions are indicated throughout.
Students may **not** bring any notes or calculators into the exam.
Where more than one mark is allocated to a question, appropriate working must be shown.
An exact answer is required to a question unless otherwise specified.
Unless otherwise indicated, diagrams in this exam are not drawn to scale.
Formula sheets can be found at the end of this exam.

This paper has been prepared independently of the Victorian Curriculum and Assessment Authority to provide additional exam preparation for students. Although references have been reproduced with permission of the Victorian Curriculum and Assessment Authority the publication is in no way connected with or endorsed by the Victorian Curriculum and Assessment Authority.

© THE HEFFERNAN GROUP 2023

This Trial Exam is licensed on a non transferable basis to the purchasing school. It may be copied by the school which has purchased it. This license does not permit distribution or copying of this Trial Exam by any other party.

Question 1 (3 marks)

Prove by mathematical induction that the number $6^n + 4$ is divisible by 5, for all $n \in \mathbb{N}$.

Question 2 (3 marks)

Find the solution to the differential equation $\frac{dy}{dx} = \frac{\sin^2(x)}{\cos(y)}$, given $y(0) = \frac{\pi}{6}$.

Give your answer as an expression for y in terms of x .

Question 3 (4 marks)

Wholegrain sourdough loaves of bread are baked for a large supermarket chain. The weight of these loaves is normally distributed with a mean of 850 g and a standard deviation of 10 g.

- a.** Find the probability that the **total** weight of four randomly selected loaves is less than 3360 g. Use $\Pr(-2 < Z < 2) = 0.95$ and give your answer correct to three decimal places. 2 marks

- b.** The loaves are placed on trays which hold 16 loaves. Find the probability that the mean weight of a loaf on a randomly selected tray is more than 855 g. Give your answer correct to three decimal places. 2 marks

Question 4 (3 marks)

Find the gradient of the curve $x \sec(2y) + \log_e(x) + \frac{6}{\pi} y^2 = 2 + \frac{\pi}{6}$ at the point $\left(1, \frac{\pi}{6}\right)$.

Express your answer in the form $\frac{a - 2a\sqrt{a}}{b}$, where $a, b \in \mathbb{N}$.

Question 5 (4 marks)

Let $z = \frac{(1+i)^6}{(\sqrt{3}-i)^4}$.

- a.** Write z in the form $a + bi$, where a and b are real constants.

3 marks

- b.** Let $z^n = d$, where $n \in \mathbb{Z}$ and $d \in \mathbb{R}$.
Find all possible values of n .

1 mark

Question 6 (5 marks)


A particle moves so that its velocity at time t is given by

$$\underline{v}(t) = 2\sin(4t)\underline{i} + 4\cos(4t)\underline{j} \quad \text{for } 0 \leq t \leq \frac{\pi}{4}.$$

- a.** Given that $\underline{r}(0) = -\frac{1}{2}\underline{i}$, find the position vector $\underline{r}(t)$ of the particle. 2 marks

- b.** Show that the Cartesian equation of the path followed by the particle is given by $4x^2 + y^2 = 1$. 1 mark

- c. Sketch the path followed by the particle on the set of axes below. Indicate the particle's starting point and label any axis intercepts with their coordinates. 2 marks



Question 8 (3 marks)

Find the value of $\lambda \in \mathbb{R}$, such that the vectors $\underline{a} = 2\underline{i} - \underline{j} + 3\underline{k}$, $\underline{b} = 3\underline{i} + 3\underline{j} - 2\underline{k}$ and $\underline{c} = -2\underline{i} + \lambda\underline{j} + \underline{k}$ are linearly dependent.

Question 10 (5 marks)

- a.** Evaluate $\int_0^1 x e^{2x} dx$. Give your answer in the form $\frac{1}{a}e^2 + \frac{1}{a}$ where $a \in \mathbb{Z}$. 2 marks

- b.** Evaluate $\int_0^1 \frac{5x-3}{(x+1)(x^2+3)} dx$. Give your answer in the form $\frac{\sqrt{b}\pi}{c} - \log_e(b)$,
where $b, c \in \mathbb{Z}$ 3 marks

Mensuration

area of a circle segment	$\frac{r^2}{2}(\theta - \sin(\theta))$	volume of a sphere	$\frac{4}{3}\pi r^3$
volume of a cylinder	$\pi r^2 h$	area of a triangle	$\frac{1}{2}bc \sin(A)$
volume of a cone	$\frac{1}{3}\pi r^2 h$	sine rule	$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$
volume of a pyramid	$\frac{1}{3}Ah$	cosine rule	$c^2 = a^2 + b^2 - 2ab \cos(C)$

Algebra, number and structure (complex numbers)

$z = x + iy = r(\cos(\theta) + i \sin(\theta)) = r \text{cis}(\theta)$	$ z = \sqrt{x^2 + y^2} = r$	
$-\pi < \text{Arg}(z) \leq \pi$	$z_1 z_2 = r_1 r_2 \text{cis}(\theta_1 + \theta_2)$	
$\frac{z_1}{z_2} = \frac{r_1}{r_2} \text{cis}(\theta_1 - \theta_2)$	de Moivre's theorem	$z^n = r^n \text{cis}(n\theta)$

Data analysis, probability and statistics

for independent random variables $X_1, X_2 \dots X_n$	$E(aX_1 + b) = aE(X_1) + b$ $E(a_1 X_1 + a_2 X_2 + \dots + a_n X_n)$ $= a_1 E(X_1) + a_2 E(X_2) + \dots + a_n E(X_n)$	
	$\text{Var}(aX_1 + b) = a^2 \text{Var}(X_1)$ $\text{Var}(a_1 X_1 + a_2 X_2 + \dots + a_n X_n)$ $= a_1^2 \text{Var}(X_1) + a_2^2 \text{Var}(X_2) + \dots + a_n^2 \text{Var}(X_n)$	
for independent identically distributed variables $X_1, X_2 \dots X_n$	$E(X_1 + X_2 + \dots + X_n) = n\mu$	
	$\text{Var}(X_1 + X_2 + \dots + X_n) = n\sigma^2$	
approximate confidence interval for μ	$\left(\bar{x} - z \frac{s}{\sqrt{n}}, \bar{x} + z \frac{s}{\sqrt{n}} \right)$	
distribution of sample mean $\bar{X}$	mean	$E(\bar{X}) = \mu$
	variance	$\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1}x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a}e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e x + c$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$	$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$
$\frac{d}{dx}(\cot(ax)) = -a \text{cosec}^2(ax)$	$\int \text{cosec}^2(ax) dx = -\frac{1}{a} \cot(ax) + c$
$\frac{d}{dx}(\sec(ax)) = a \sec(ax) \tan(ax)$	$\int \sec(ax) \tan(ax) dx = \frac{1}{a} \sec(ax) + c$
$\frac{d}{dx}(\text{cosec}(ax)) = -a \text{cosec}(ax) \cot(ax)$	$\int \text{cosec}(ax) \cot(ax) dx = -\frac{1}{a} \text{cosec}(ax) + c$
$\frac{d}{dx}(\sin^{-1}(ax)) = \frac{a}{\sqrt{1-(ax)^2}}$	$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\cos^{-1}(ax)) = \frac{-a}{\sqrt{1-(ax)^2}}$	$\int \frac{-1}{\sqrt{a^2 - x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\tan^{-1}(ax)) = \frac{a}{1+(ax)^2}$	$\int \frac{a}{a^2 + x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$
	$\int (ax + b)^n dx = \frac{1}{a(n+1)}(ax + b)^{n+1} + c, n \neq -1$
	$\int \frac{1}{ax + b} dx = \frac{1}{a} \log_e ax + b + c$

Calculus - continued

product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$	
quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$	
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$	
integration by parts	$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$	
Euler's method	If $\frac{dy}{dx} = f(x, y), x_0 = a$ and $y_0 = b$ then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + h \times f(x_n, y_n)$	
arc length parametric	$\int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$	
surface area Cartesian about x-axis	$\int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$	
surface area Cartesian about y-axis	$\int_{y_1}^{y_2} 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$	
surface area parametric about x-axis	$\int_{t_1}^{t_2} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$	
surface area parametric about y-axis	$\int_{t_1}^{t_2} 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$	

Kinematics

acceleration	$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$	
constant acceleration formulas	$v = u + at$	$s = ut + \frac{1}{2}at^2$
	$v^2 = u^2 + 2as$	$s = \frac{1}{2}(u + v)t$

Vectors in two and three dimensions

$\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j} + z(t)\vec{k}$	$ \vec{r}(t) = \sqrt{x(t)^2 + y(t)^2 + z(t)^2}$
	$\dot{\vec{r}}(t) = \frac{d\vec{r}}{dt} = \frac{dx}{dt}\vec{i} + \frac{dy}{dt}\vec{j} + \frac{dz}{dt}\vec{k}$
for $\vec{r}_1 = x_1\vec{i} + y_1\vec{j} + z_1\vec{k}$ and $\vec{r}_2 = x_2\vec{i} + y_2\vec{j} + z_2\vec{k}$	vector scalar product $\vec{r}_1 \cdot \vec{r}_2 = \vec{r}_1 \vec{r}_2 \cos(\theta) = x_1x_2 + y_1y_2 + z_1z_2$
	vector cross product $\vec{r}_1 \times \vec{r}_2 = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} = (y_1z_2 - y_2z_1)\vec{i} + (x_2z_1 - x_1z_2)\vec{j} + (x_1y_2 - x_2y_1)\vec{k}$
vector equation of a line	$\vec{r}(t) = \vec{r}_1 + t\vec{r}_2 = (x_1 + x_2t)\vec{i} + (y_1 + y_2t)\vec{j} + (z_1 + z_2t)\vec{k}$
parametric equation of a line	$x(t) = x_1 + x_2t \quad y(t) = y_1 + y_2t \quad z(t) = z_1 + z_2t$
vector equation of a plane	$\vec{r}(s, t) = \vec{r}_0 + s\vec{r}_1 + t\vec{r}_2$ $= (x_0 + x_1s + x_2t)\vec{i} + (y_0 + y_1s + y_2t)\vec{j} + (z_0 + z_1s + z_2t)\vec{k}$
parametric equation of a plane	$x(s, t) = x_0 + x_1s + x_2t, \quad y(s, t) = y_0 + y_1s + y_2t, \quad z(s, t) = z_0 + z_1s + z_2t$
Cartesian equation of a plane	$ax + by + cz = d$

Circular functions

$\cos^2(x) + \sin^2(x) = 1$	
$1 + \tan^2(x) = \sec^2(x)$	$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$
$\sin(x + y) = \sin(x)\cos(y) + \cos(x)\sin(y)$	$\sin(x - y) = \sin(x)\cos(y) - \cos(x)\sin(y)$
$\cos(x + y) = \cos(x)\cos(y) - \sin(x)\sin(y)$	$\cos(x - y) = \cos(x)\cos(y) + \sin(x)\sin(y)$
$\tan(x + y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$	$\tan(x - y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$
$\sin(2x) = 2\sin(x)\cos(x)$	
$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$	$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$
$\sin^2(ax) = \frac{1}{2}(1 - \cos(2ax))$	$\cos^2(ax) = \frac{1}{2}(1 + \cos(2ax))$

Mathematics Formula Sheets reproduced by permission; © VCAA. The VCAA does not endorse or make any warranties regarding this study resource. Past VCAA VCE® exams and related content can be accessed directly at www.vcaa.vic.edu.au