

2023 VCE Specialist Mathematics Year 12 Trial Examination 1



Kilbaha Education

Quality educational content

**Kilbaha Education
PO Box 2227
Kew Vic 3101
Australia**

**Tel: (03) 9018 5376
kilbaha@gmail.com
<https://kilbaha.com.au>**

All publications from Kilbaha Education are digital and are supplied to the purchasing school in both WORD and PDF formats with a school site licence to reproduce for students in both print and electronic formats.



Kilbaha Education

Quality educational content

Kilbaha Education (Est. 1978) (ABN 47 065 111 373)
PO Box 2227
Kew Vic 3101
Australia

Tel: +613 9018 5376
Email: kilbaha@gmail.com
Web: <https://kilbaha.com.au>

IMPORTANT COPYRIGHT NOTICE FOR KILBAHA PUBLICATIONS

- (1) The material is copyright. Subject to statutory exception and to the provisions of the relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Kilbaha Pty Ltd.
- (2) The contents of these works are copyrighted. Unauthorised copying of any part of these works is illegal and detrimental to the interests of the author(s).
- (3) For authorised copying within Australia please check that your institution has a licence from <https://www.copyright.com.au> This permits the copying of small parts of the material, in limited quantities, within the conditions set out in the licence.
- (4) All pages of Kilbaha files must be counted in Copyright Agency Limited (CAL) surveys.
- (5) Kilbaha files must **not** be uploaded to the Internet.
- (6) Kilbaha files may be placed on a password protected school Intranet.

Kilbaha educational content has no official status and is not endorsed by any State or Federal Government Education Authority.

While every care has been taken, no guarantee is given that the content is free from error. Please contact us if you believe you have found an error.

CAUTION NEEDED!

All Web Links when created linked to appropriate Web Sites. Teachers and parents must always check links before using them with students to ensure that students are protected from unsuitable Web Content. Kilbaha Education is not responsible for links that have been changed in its publications or links that have been redirected.

**Victorian Certificate of Education
2023**

STUDENT NUMBER

Figures
Words

Letter

--

SPECIALIST MATHEMATICS

Trial Written Examination 1

Reading time: 15 minutes

Total writing time: 1 hour

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
10	10	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or white out liquid/tape.

Materials supplied

- Question and answer book of 22 pages with a detachable sheet of miscellaneous formulas at the end of this booklet.

Instructions

- Detach the formula sheet from the end of this book during reading time.
- Write your **student number** in the space provided above on this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Answer **all** questions in the spaces provided.
 Unless otherwise specified an **exact** answer is required to a question.
 In questions where more than one mark is available, appropriate working **must** be shown.
 Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.
 Take the **acceleration due to gravity** to have magnitude $g \text{ m/s}^2$, where $g = 9.8$.

Solve the equation $z^4 + 16 = 0$, $z \in \mathbb{C}$, giving your answers in rectangular $a + bi$ form.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Solve the differential equation $\frac{dy}{dx} = \frac{x(9+4y^2)}{6}$ given $y(0) = \frac{3}{2}$.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question 3 (3 marks)

Let $f : D \rightarrow R$, $f(x) = \frac{x}{x^2 - 4}$, where D is the maximal domain of f .


- i. Find $f'(x)$ and hence show that the graph of f has no turning points.

1 mark

- ii. Given that $f''(x) = \frac{2x(x^2 + 12)}{(x^2 - 4)^3}$ find the coordinates of any points of inflexion and

sketch the graph of $f(x) = \frac{x}{x^2 - 4}$ on the axes below, labelling the equations of all asymptotes.

2 marks



Question 4 (3 marks)

A particle moves in a straight line. At a time t seconds, the particle has a displacement of x metres and a velocity of $v \text{ ms}^{-1}$ and an acceleration $a \text{ ms}^{-2}$. Initially the particle is at the origin and has a velocity of 2 ms^{-1} . If the acceleration $a = -\frac{4}{3}e^{-\frac{2x}{3}}$ and the velocity of the particle is always positive.

- a. Show that $v = 2e^{-\frac{x}{3}}$

1 mark

- b. After three seconds, the particle has a displacement of $\log_e(s)$, find the value of s .

2 marks

Question 5 (3 marks)

The weights of a piece of corn are normally distributed with a mean of 100 gm with a variance of 12 gm². Z has the standard normal distribution and given that $\Pr(Z < 1.5) = 0.933$.

- a. Determine the probability that the total weight of three independent pieces of corn will be between 291 and 300 gms. Give your answer correct to three decimal places.


2 marks

- b. An **approximate** 95% confidence interval for n pieces of corn was found to be 99 to 101 gm. Given that $\Pr(-2 < Z < 2) = 0.95$, where Z is a standard normal random variable. Determine the value of n .

1 mark

- b.** It is known that the number of kangaroos on the plantation was increasing most rapidly around the year 2011. Sketch the graph of K versus t on the axes below, clearly labelling the point of inflexion, the equations of any asymptotes and coordinates of any axial intercepts.

1 mark



Prove by induction that $5^{2n} + 3n - 1$ is divisible by 9 for $n \in N$.

[illegible]

Question 9 (6 marks)

A plane has the vector equation $\vec{r}(s, t) = (3 + 4t + 3s)\vec{i} + (-2 + t + 2s)\vec{j} + (1 + 2t - s)\vec{k}$

- a. Show that the equation of the plane can be written in the form $x - 2y - z = 6$.

2 marks

- b. One line has the vector equation $\vec{r}(t) = 2\vec{i} - \vec{j} + z_0\vec{k} + t(\vec{i} - 2\vec{j} + c\vec{k})$, find the values of z_0 and c for which the line and the plane do not intersect.

2 marks

- 2 marks

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

c. Hence evaluate I_3 .

1 mark

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

**Kilbaha Education
PO Box 2227
Kew Vic 3101
Australia**

Tel: (03) 9018 5376
kilbaha@gmail.com
<https://kilbaha.com.au>

SPECIALIST MATHEMATICS

Written examination 1

FORMULA SHEET

Directions to students

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

Specialist Mathematics formulas

Mensuration

area of a circle segment	$\frac{r^2}{2}(\theta - \sin(\theta))$	volume of a sphere	$\frac{4}{3}\pi r^3$
volume of a cylinder	$\pi r^2 h$	area of a triangle	$\frac{1}{2}bc \sin(A)$
volume of a cone	$\frac{1}{3}\pi r^2 h$	sine rule	$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$
volume of a pyramid	$\frac{1}{3}Ah$	cosine rule	$c^2 = a^2 + b^2 - 2ab \cos(C)$

Algebra, number and structure (complex numbers)

$z = x + yi = r(\cos(\theta) + i \sin(\theta)) = r \operatorname{cis}(\theta)$	$ z = \sqrt{x^2 + y^2} = r$	
$-\pi < \operatorname{Arg}(z) \leq \pi$	$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$	
$\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$	de Moivre's theorem	$z^n = r^n \operatorname{cis}(n\theta)$

Circular (trigonometric) functions

$\cos^2(x) + \sin^2(x) = 1$	
$1 + \tan^2(x) = \sec^2(x)$	$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$
$\sin(x + y) = \sin(x)\cos(y) + \cos(x)\sin(y)$	$\sin(x - y) = \sin(x)\cos(y) - \cos(x)\sin(y)$
$\cos(x + y) = \cos(x)\cos(y) - \sin(x)\sin(y)$	$\cos(x - y) = \cos(x)\cos(y) + \sin(x)\sin(y)$
$\tan(x + y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$	$\tan(x - y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$
$\sin(2x) = 2\sin(x)\cos(x)$	
$\cos(2x) = \cos^2(x) - \sin^2(x)$ $= 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$	$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$
$\sin^2(ax) = \frac{1}{2}(1 - \cos(2ax))$	$\cos^2(ax) = \frac{1}{2}(1 + \cos(2ax))$

Data analysis, probability and statistics

for independent random variables $X_1, X_2, \dots, X_n$	$E(aX_1 + b) = aE(X_1) + b$ $E(a_1X_1 + a_2X_2 + \dots + a_nX_n)$ $= a_1E(X_1) + a_2E(X_2) + \dots + a_nE(X_n)$	
	$\text{Var}(aX_1 + b) = a^2\text{Var}(X_1)$ $\text{Var}(a_1X_1 + a_2X_2 + \dots + a_nX_n)$ $= a_1^2\text{Var}(X_1) + a_2^2\text{Var}(X_2) + \dots + a_n^2\text{Var}(X_n)$	
for independent identically distributed variables $X_1, X_2, \dots, X_n$	$E(X_1 + X_2 + \dots + X_n) = n\mu$	
	$\text{Var}(X_1 + X_2 + \dots + X_n) = n\sigma^2$	
approximate confidence interval for μ	$\left(\bar{x} - z \frac{s}{\sqrt{n}}, \bar{x} + z \frac{s}{\sqrt{n}} \right)$	
distribution of sample mean $\bar{X}$	mean	$E(\bar{X}) = \mu$
	variance	$\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$

Vectors in two and three dimensions

$\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j} + z(t)\vec{k}$	$ \vec{r}(t) = \sqrt{x(t)^2 + y(t)^2 + z(t)^2}$
	$\dot{\vec{r}}(t) = \frac{d\vec{r}}{dt} = \frac{dx}{dt}\vec{i} + \frac{dy}{dt}\vec{j} + \frac{dz}{dt}\vec{k}$
for $\vec{r}_1 = x_1\vec{i} + y_1\vec{j} + z_1\vec{k}$ and $\vec{r}_2 = x_2\vec{i} + y_2\vec{j} + z_2\vec{k}$	vector scalar product $\vec{r}_1 \cdot \vec{r}_2 = \vec{r}_1 \vec{r}_2 \cos(\theta) = x_1x_2 + y_1y_2 + z_1z_2$ vector cross product $\vec{r}_1 \times \vec{r}_2 = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} = (y_1z_2 - y_2z_1)\vec{i} + (x_2z_1 - x_1z_2)\vec{j} + (x_1y_2 - x_2y_1)\vec{k}$
vector equation of a line	$\vec{r}(t) = \vec{r}_1 + t\vec{r}_2 = (x_1 + x_2t)\vec{i} + (y_1 + y_2t)\vec{j} + (z_1 + z_2t)\vec{k}$
parametric equation of line	$x(t) = x_1 + x_2t \quad y(t) = y_1 + y_2t \quad z(t) = z_1 + z_2t$
vector equation of a plane	$\vec{r}(s, t) = \vec{r}_0 + s\vec{r}_1 + t\vec{r}_2$ $= (x_0 + x_1s + x_2t)\vec{i} + (y_0 + y_1s + y_2t)\vec{j} + (z_0 + z_1s + z_2t)\vec{k}$
parametric equation of a plane	$x(s, t) = x_0 + x_1s + x_2t \quad y(s, t) = y_0 + y_1s + y_2t \quad z(s, t) = z_0 + z_1s + z_2t$
Cartesian equation of a plane	$ax + by + cz = d$

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e(x) + c$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$	$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$
$\frac{d}{dx}(\cot(ax)) = -a \operatorname{cosec}^2(ax)$	$\int \operatorname{cosec}^2(ax) dx = -\frac{1}{a} \cot(ax) + c$
$\frac{d}{dx}(\sec(ax)) = a \sec(ax) \tan(ax)$	$\int \sec(ax) \tan(ax) dx = \frac{1}{a} \sec(ax) + c$
$\frac{d}{dx}(\operatorname{cosec}(ax)) = -a \operatorname{cosec}(ax) \cot(ax)$	$\int \operatorname{cosec}(ax) \cot(ax) dx = -\frac{1}{a} \operatorname{cosec}(ax) + c$
$\frac{d}{dx}(\sin^{-1}(ax)) = \frac{1}{\sqrt{1-(ax)^2}}$	$\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\cos^{-1}(ax)) = \frac{-1}{\sqrt{1-(ax)^2}}$	$\int \frac{-1}{\sqrt{a^2-x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\tan^{-1}(ax)) = \frac{a}{1+(ax)^2}$	$\int \frac{a}{a^2+x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$
	$\int (ax+b)^n dx = \frac{1}{a(n+1)} (ax+b)^{n+1} + c, n \neq -1$
	$\int (ax+b)^{-1} dx = \frac{1}{a} \log_e(ax+b) + c$

Calculus- continued

product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$
quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
integration by parts	$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$
Euler's method	If $\frac{dy}{dx} = f(x)$, $x_0 = a$ and $y_0 = b$, then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + hf(x_n, y_n)$
arc length parametric	$\int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$
surface area Cartesian about the x-axis	$\int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$
surface area Cartesian about the y-axis	$\int_{y_1}^{y_2} 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$
surface area parametric about the x-axis	$\int_{t_1}^{t_2} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$
surface area parametric about the y-axis	$\int_{t_1}^{t_2} 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$

Kinematics

acceleration	$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$	
constant acceleration formulas	$v = u + at$	$s = ut + \frac{1}{2}at^2$
	$v^2 = u^2 + 2as$	$s = \frac{1}{2}(u + v)t$

END OF FORMULA SHEET