

Victorian Certificate of Education 2024

STUDENT NUMBER

Figures
Words

Letter

--

SPECIALIST MATHEMATICS

Trial Written Examination 1

Reading time: 15 minutes

Total writing time: 1 hour

QUESTION AND ANSWER BOOK

Structure of book

<i>Number of questions</i>	<i>Number of questions to be answered</i>	<i>Number of marks</i>
9	9	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers.
- Students are NOT permitted to bring into the examination room: any technology (calculators or software), notes of any kind, blank sheets of paper and/or white out liquid/tape.

Materials supplied

- Question and answer book of 21 pages with a detachable sheet of miscellaneous formulas at the end of this booklet.

Instructions

- Detach the formula sheet from the end of this book during reading time.
- Write your **student number** in the space provided above on this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.

Instructions

Answer **all** questions in the spaces provided.

Unless otherwise specified an **exact** answer is required to a question.

In questions where more than one mark is available, appropriate working **must** be shown.

Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Take the **acceleration due to gravity** to have magnitude $g \text{ m/s}^2$, where $g = 9.8$.

Question 1 (5 marks)

Given the differential equation $\frac{dy}{dx} = 2x\sqrt{16-9y^2}$, $y(1) = \frac{2}{3}$.

- a. Solve the differential equation, expressing your answer in the form $y = f(x)$.

3 marks

- b. Find the value of $\frac{d^2y}{dx^2}$ when $x=1$ and $y = \frac{2}{3}$.

2 marks

Question 2 (4 marks)

Let $f : D \rightarrow R$, $f(x) = \frac{x^2 - 5x + 4}{x}$, where D is the maximal domain of f .

- i. Determine the equations of all the asymptotes on the graph of $y = f(x)$.

1 mark

- ii. Determine the coordinates of all the turning points on the graph of $y = f(x)$.


1 mark

- iii. Show that the graph of $y = f(x)$ has no points of inflexion.

1 mark

- iv. Sketch the graph of $y = f(x) = \frac{x^2 - 5x + 4}{x}$ on the axes below, labelling turning points, axial intercepts and the equations of all asymptotes.

1 mark



b. Show that $v = \frac{7 \left(1 - e^{-\frac{14t}{5}} \right)}{1 + e^{-\frac{14t}{5}}}$.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Question 4 (5 marks)

Given the three points $P(1, 2, 3)$, $Q(-1, 2, -1)$, $R(-1, -2, 5)$.

- a. Show that the equation of the plane which passes through the points P , Q and R is given by $-4x + 3y + 2z = 8$.

2 marks

- b. The plane $-4x + 3y + 2z = 8$ and the line $2 - x = \frac{y - 4}{2} = \frac{z - 2}{c}$ intersect at an acute angle of $\cos^{-1}\left(\frac{5\sqrt{29}}{29}\right)$, determine the value of c .

3 marks

- c. Let $u = z + \frac{1}{z}$ show that $Q(z) = 0$ can be expressed in the form $u^2 - u - 1 = 0$.

1 mark

- d. Hence find the value $\cos\left(\frac{3\pi}{5}\right)$

1 mark

Question 6 (3 marks)

The weights of a carrots from a local farm are normally distributed with a mean of 70 grams with a standard deviation of 5 grams.

Z has the standard normal distribution and given that $\Pr(-1.2 < Z < 1.2) = 0.770$.

- a.** Determine the probability that the total weight of four independent carrots from the local farm will have a weight greater than 292 grams.

Give your answer correct to three decimal places.

2 marks

- b.** Find the probability that the mean weight of four carrots is between 67 and 73 grams.
Give your answer correct to three decimal places.

1 mark

[illegible]

2 marks

[illegible]

Find $\int t^3 \sin(t^2) dt$

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

Kilbaha Education (Est. 1978) (ABN 47 065 111 373) PO Box 3229 Cotham Vic 3101 Australia	PayID: 47065111373 Email: kilbaha@gmail.com Tel: (03) 9018 5376 Web: https://kilbaha.com.au
---	--

SPECIALIST MATHEMATICS

Written examination 1

FORMULA SHEET

Directions to students

Detach this formula sheet during reading time.

This formula sheet is provided for your reference.

Specialist Mathematics formulas

Mensuration

area of a circle segment	$\frac{r^2}{2}(\theta - \sin(\theta))$	volume of a sphere	$\frac{4}{3}\pi r^3$
volume of a cylinder	$\pi r^2 h$	area of a triangle	$\frac{1}{2}bc \sin(A)$
volume of a cone	$\frac{1}{3}\pi r^2 h$	sine rule	$\frac{a}{\sin(A)} = \frac{b}{\sin(B)} = \frac{c}{\sin(C)}$
volume of a pyramid	$\frac{1}{3}Ah$	cosine rule	$c^2 = a^2 + b^2 - 2ab \cos(C)$

Algebra, number and structure (complex numbers)

$z = x + yi = r(\cos(\theta) + i \sin(\theta)) = r \operatorname{cis}(\theta)$	$ z = \sqrt{x^2 + y^2} = r$	
$-\pi < \operatorname{Arg}(z) \leq \pi$	$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$	
$\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$	de Moivre's theorem	$z^n = r^n \operatorname{cis}(n\theta)$

Circular (trigonometric) functions

$\cos^2(x) + \sin^2(x) = 1$	
$1 + \tan^2(x) = \sec^2(x)$	$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$
$\sin(x + y) = \sin(x)\cos(y) + \cos(x)\sin(y)$	$\sin(x - y) = \sin(x)\cos(y) - \cos(x)\sin(y)$
$\cos(x + y) = \cos(x)\cos(y) - \sin(x)\sin(y)$	$\cos(x - y) = \cos(x)\cos(y) + \sin(x)\sin(y)$
$\tan(x + y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$	$\tan(x - y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$
$\sin(2x) = 2\sin(x)\cos(x)$	
$\cos(2x) = \cos^2(x) - \sin^2(x)$ $= 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$	$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$
$\sin^2(ax) = \frac{1}{2}(1 - \cos(2ax))$	$\cos^2(ax) = \frac{1}{2}(1 + \cos(2ax))$

Data analysis, probability and statistics

for independent random variables $X_1, X_2, \dots, X_n$	$E(aX_1 + b) = aE(X_1) + b$ $E(a_1X_1 + a_2X_2 + \dots + a_nX_n)$ $= a_1E(X_1) + a_2E(X_2) + \dots + a_nE(X_n)$	
	$\text{Var}(aX_1 + b) = a^2\text{Var}(X_1)$ $\text{Var}(a_1X_1 + a_2X_2 + \dots + a_nX_n)$ $= a_1^2\text{Var}(X_1) + a_2^2\text{Var}(X_2) + \dots + a_n^2\text{Var}(X_n)$	
for independent identically distributed variables $X_1, X_2, \dots, X_n$	$E(X_1 + X_2 + \dots + X_n) = n\mu$	
	$\text{Var}(X_1 + X_2 + \dots + X_n) = n\sigma^2$	
approximate confidence interval for μ	$\left(\bar{x} - z \frac{s}{\sqrt{n}}, \bar{x} + z \frac{s}{\sqrt{n}} \right)$	
distribution of sample mean $\bar{X}$	mean	$E(\bar{X}) = \mu$
	variance	$\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$

Vectors in two and three dimensions

$\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j} + z(t)\vec{k}$	$ \vec{r}(t) = \sqrt{x(t)^2 + y(t)^2 + z(t)^2}$
	$\dot{\vec{r}}(t) = \frac{d\vec{r}}{dt} = \frac{dx}{dt}\vec{i} + \frac{dy}{dt}\vec{j} + \frac{dz}{dt}\vec{k}$
for $\vec{r}_1 = x_1\vec{i} + y_1\vec{j} + z_1\vec{k}$ and $\vec{r}_2 = x_2\vec{i} + y_2\vec{j} + z_2\vec{k}$	vector scalar product $\vec{r}_1 \cdot \vec{r}_2 = \vec{r}_1 \vec{r}_2 \cos(\theta) = x_1x_2 + y_1y_2 + z_1z_2$
	vector cross product $\vec{r}_1 \times \vec{r}_2 = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} = (y_1z_2 - y_2z_1)\vec{i} + (x_2z_1 - x_1z_2)\vec{j} + (x_1y_2 - x_2y_1)\vec{k}$
vector equation of a line	$\vec{r}(t) = \vec{r}_1 + t\vec{r}_2 = (x_1 + x_2t)\vec{i} + (y_1 + y_2t)\vec{j} + (z_1 + z_2t)\vec{k}$
parametric equation of line	$x(t) = x_1 + x_2t \quad y(t) = y_1 + y_2t \quad z(t) = z_1 + z_2t$
vector equation of a plane	$\vec{r}(s, t) = \vec{r}_0 + s\vec{r}_1 + t\vec{r}_2$ $= (x_0 + x_1s + x_2t)\vec{i} + (y_0 + y_1s + y_2t)\vec{j} + (z_0 + z_1s + z_2t)\vec{k}$
parametric equation of a plane	$x(s, t) = x_0 + x_1s + x_2t \quad y(s, t) = y_0 + y_1s + y_2t \quad z(s, t) = z_0 + z_1s + z_2t$
Cartesian equation of a plane	$ax + by + cz = d$

Calculus

$\frac{d}{dx}(x^n) = nx^{n-1}$	$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$
$\frac{d}{dx}(e^{ax}) = ae^{ax}$	$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$
$\frac{d}{dx}(\log_e(x)) = \frac{1}{x}$	$\int \frac{1}{x} dx = \log_e(x) + c$
$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$	$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$
$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$	$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$
$\frac{d}{dx}(\tan(ax)) = a \sec^2(ax)$	$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$
$\frac{d}{dx}(\cot(ax)) = -a \operatorname{cosec}^2(ax)$	$\int \operatorname{cosec}^2(ax) dx = -\frac{1}{a} \cot(ax) + c$
$\frac{d}{dx}(\sec(ax)) = a \sec(ax) \tan(ax)$	$\int \sec(ax) \tan(ax) dx = \frac{1}{a} \sec(ax) + c$
$\frac{d}{dx}(\operatorname{cosec}(ax)) = -a \operatorname{cosec}(ax) \cot(ax)$	$\int \operatorname{cosec}(ax) \cot(ax) dx = -\frac{1}{a} \operatorname{cosec}(ax) + c$
$\frac{d}{dx}(\sin^{-1}(ax)) = \frac{1}{\sqrt{1-(ax)^2}}$	$\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\cos^{-1}(ax)) = \frac{-1}{\sqrt{1-(ax)^2}}$	$\int \frac{-1}{\sqrt{a^2-x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$
$\frac{d}{dx}(\tan^{-1}(ax)) = \frac{a}{1+(ax)^2}$	$\int \frac{a}{a^2+x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$
	$\int (ax+b)^n dx = \frac{1}{a(n+1)} (ax+b)^{n+1} + c, n \neq -1$
	$\int (ax+b)^{-1} dx = \frac{1}{a} \log_e(ax+b) + c$

Calculus- continued

product rule	$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$
quotient rule	$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$
chain rule	$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$
integration by parts	$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$
Euler's method	If $\frac{dy}{dx} = f(x)$, $x_0 = a$ and $y_0 = b$, then $x_{n+1} = x_n + h$ and $y_{n+1} = y_n + hf(x_n, y_n)$
arc length parametric	$\int_{t_1}^{t_2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$
surface area Cartesian about the x-axis	$\int_{x_1}^{x_2} 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$
surface area Cartesian about the y-axis	$\int_{y_1}^{y_2} 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$
surface area parametric about the x-axis	$\int_{t_1}^{t_2} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$
surface area parametric about the y-axis	$\int_{t_1}^{t_2} 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$

Kinematics

acceleration	$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$	
constant acceleration formulas	$v = u + at$	$s = ut + \frac{1}{2}at^2$
	$v^2 = u^2 + 2as$	$s = \frac{1}{2}(u + v)t$

END OF FORMULA SHEET