

CHEMISTRY

Written examination 2

DATA BOOK

Directions to students

This data book is provided as a reference.

Make sure that you remove this data book from the question and answer book during reading time.

Any writings, jottings, notes or drawings made on this data book will **not** be considered in the marking.

At the end of the examination, ensure that you do **not** leave the data book in the question and answer book.

You may keep this data book.

Table of contents

	page
1. Periodic table of the elements	3
2. The electrochemical series	4
3. Physical constants	5
4. SI prefixes, their symbols and values	5
5. ^1H NMR data	5–6
6. ^{13}C NMR data	7
7. Infrared absorption data	7
8. 2-amino acids (α -amino acids)	8–9
9. Formulas of some fatty acids	10
10. Structural formulas of some important biomolecules	10
11. Acid-base indicators	11
12. Acidity constants, K_{a} , of some weak acids	11
13. Values of molar enthalpy of combustions of some common fuels at 298 K and 101.3 kPa	11

1. Periodic table of the elements

1 H 1.0 Hydrogen		atomic number										79 Au 197.0 Gold		symbol of element										2 He 4.0 Helium											
3 Li 6.9 Lithium		4 Be 9.0 Beryllium		relative atomic mass																		10 Ne 20.1 Neon													
11 Na 23.0 Sodium		12 Mg 24.3 Magnesium		name of element																		17 Cl 35.5 Chlorine													
19 K 39.1 Potassium		20 Ca 40.1 Calcium		21 Sc 44.9 Scandium		22 Ti 47.9 Titanium		23 V 50.9 Vanadium		24 Cr 52.0 Chromium		25 Mn 54.9 Manganese		26 Fe 55.9 Iron		27 Co 58.9 Cobalt		28 Ni 58.7 Nickel		29 Cu 63.6 Copper		30 Zn 65.4 Zinc		31 Ga 69.7 Gallium		32 Ge 72.6 Germanium		33 As 74.9 Arsenic		34 Se 79.0 Selenium		35 Br 79.9 Bromine		36 Kr 83.8 Krypton	
37 Rb 85.5 Rubidium		38 Sr 87.6 Strontium		39 Y 88.9 Yttrium		40 Zr 91.2 Zirconium		41 Nb 92.9 Niobium		42 Mo 95.9 Molybdenum		43 Tc 98.1 Technetium		44 Ru 101.1 Ruthenium		45 Rh 102.9 Rhodium		46 Pd 106.4 Palladium		47 Ag 107.9 Silver		48 Cd 112.4 Cadmium		49 In 114.8 Indium		50 Sn 118.7 Tin		51 Sb 121.8 Antimony		52 Te 127.6 Tellurium		53 I 126.9 Iodine		54 Xe 131.3 Xenon	
55 Cs 132.9 Caesium		56 Ba 137.3 Barium		57 La 138.9 Lanthanum		72 Hf 178.5 Hafnium		73 Ta 180.9 Tantalum		74 W 183.8 Tungsten		75 Re 186.2 Rhenium		76 Os 190.2 Osmium		77 Ir 192.2 Iridium		78 Pt 195.1 Platinum		79 Au 197.0 Gold		80 Hg 200.6 Mercury		81 Tl 204.4 Thallium		82 Pb 207.2 Lead		83 Bi 209.0 Bismuth		84 Po (209) Polonium		85 At (210) Astatine		86 Rn (222) Radon	
87 Fr (223) Francium		88 Ra (226) Radium		89 Ac (227) Actinium		104 Rf (261) Rutherfordium		105 Db (262) Dubnium		106 Sg (266) Seaborgium		107 Bh (264) Bohrium		108 Hs (277) Hassium		109 Mt (268) Meitnerium		110 Ds (271) Darmstadtium		111 Rg (272) Roentgenium		112 Uub (285) Ununbium		114 Uuq (289) Ununquadium		116 Uuh (294) Ununhexium		118 Uuo (294) Ununoctium							

2. The electrochemical series

	E° in volt
$\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-(\text{aq})$	+2.87
$\text{H}_2\text{O}_2(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$	+1.77
$\text{Au}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Au}(\text{s})$	+1.68
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$	+1.23
$\text{Br}_2(\text{l}) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-(\text{aq})$	+1.09
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2(\text{aq})$	+0.68
$\text{I}_2(\text{s}) + 2\text{e}^- \rightleftharpoons 2\text{I}^-(\text{aq})$	+0.54
$\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightleftharpoons 4\text{OH}^-(\text{aq})$	+0.40
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}(\text{aq})$	+0.15
$\text{S}(\text{s}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+0.14
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0.00
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ni}(\text{s})$	-0.23
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Fe}(\text{s})$	-0.44
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Mn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mn}(\text{s})$	-1.03
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightleftharpoons \text{Al}(\text{s})$	-1.67
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mg}(\text{s})$	-2.34
$\text{Na}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ca}(\text{s})$	-2.87
$\text{K}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{K}(\text{s})$	-2.93
$\text{Li}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Li}(\text{s})$	-3.02

3. Physical constants

Avogadro's constant (N_A) = $6.02 \times 10^{23} \text{ mol}^{-1}$

Charge on one electron = $-1.60 \times 10^{-19} \text{ C}$

Faraday constant (F) = $96\,500 \text{ C mol}^{-1}$

Gas constant (R) = $8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

Ionic product for water (K_w) = $1.00 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$ at 298 K
(Self ionisation constant)

Molar volume (V_m) of an ideal gas at 273 K, 101.3 kPa (STP) = 22.4 L mol^{-1}

Molar volume (V_m) of an ideal gas at 298 K, 101.3 kPa (SLC) = 24.5 L mol^{-1}

Specific heat capacity (c) of water = $4.18 \text{ J g}^{-1} \text{ K}^{-1}$

Density (d) of water at 25°C = 1.00 g mL^{-1}

1 atm = 101.3 kPa = 760 mm Hg

0°C = 273 K

4. SI prefixes, their symbols and values

SI prefix	Symbol	Value
giga	G	10^9
mega	M	10^6
kilo	k	10^3
deci	d	10^{-1}
centi	c	10^{-2}
milli	m	10^{-3}
micro	μ	10^{-6}
nano	n	10^{-9}
pico	p	10^{-12}




5. ^1H NMR data

Typical proton shift values relative to TMS = 0

These can differ slightly in different solvents. Where more than one proton environment is shown in the formula, the shift refers to the ones in bold letters.

Type of proton	Chemical shift (ppm)
R-CH ₃	0.9
R-CH ₂ -R	1.3
RCH = CH-CH ₃	1.7
R ₃ -CH	2.0
$\text{CH}_3-\text{C} \begin{array}{l} \text{O} \\ \parallel \\ \text{OR} \end{array}$ or $\text{CH}_3-\text{C} \begin{array}{l} \text{O} \\ \parallel \\ \text{NHR} \end{array}$	2.0

TURN OVER

Type of proton	Chemical shift (ppm)
$\begin{array}{c} \text{R} \quad \text{CH}_3 \\ \diagdown \quad / \\ \text{C} \\ \\ \text{O} \end{array}$	2.1
R-CH ₂ -X (X = F, Cl, Br or I)	3–4
R-CH ₂ -OH	3.6
$\begin{array}{c} \text{O} \\ // \\ \text{R}-\text{C} \\ \backslash \\ \text{NHCH}_2\text{R} \end{array}$	3.2
R-O-CH ₃ or R-O-CH ₂ R	3.3
	4.1
$\begin{array}{c} \text{O} \\ // \\ \text{R}-\text{C} \\ \backslash \\ \text{OCH}_2\text{R} \end{array}$	4.1
R-O-H	1–6 (varies considerably under different conditions)
R-NH ₂	1–5
RHC=CH ₂	4.6–6.0
	7.0
	7.3
$\begin{array}{c} \text{O} \\ // \\ \text{R}-\text{C} \\ \backslash \\ \text{NHCH}_2\text{R} \end{array}$	8.1
$\begin{array}{c} \text{O} \\ // \\ \text{R}-\text{C} \\ \backslash \\ \text{H} \end{array}$	9–10
$\begin{array}{c} \text{O} \\ // \\ \text{R}-\text{C} \\ \backslash \\ \text{O}-\text{H} \end{array}$	11.5

6. ^{13}C NMR data

Type of carbon	Chemical shift (ppm)
R-CH ₃	8–25
R-CH ₂ -R	20–45
R ₃ -CH	40–60
R ₄ -C	36–45
R-CH ₂ -X	15–80
RC-NH ₂	35–70
R-CH ₂ -OH	50–90
RC≡CR	75–95
RC=CR	110–150
RCOOH	160–185

7. Infrared absorption data




Characteristic range for infrared absorption

Bond	Wave number (cm ⁻¹)
C-Cl	700–800
C-C	750–1100
C-O	1000–1300
C=C	1610–1680
C=O	1670–1750
O-H (acids)	2500–3300
C-H	2850–3300
O-H (alcohols)	3200–3550
N-H (primary amines)	3350–3500

TURN OVER

8. 2-amino acids (α -amino acids)


Name	Symbol	Structure
alanine	Ala	$\begin{array}{c} \text{CH}_3 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
arginine	Arg	$\begin{array}{c} \text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
asparagine	Asn	$\begin{array}{c} \text{O} \\ \\ \text{CH}_2-\text{C}-\text{NH}_2 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
aspartic acid	Asp	$\begin{array}{c} \text{CH}_2-\text{COOH} \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
cysteine	Cys	$\begin{array}{c} \text{CH}_2-\text{SH} \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
glutamine	Gln	$\begin{array}{c} \text{O} \\ \\ \text{CH}_2-\text{CH}_2-\text{C}-\text{NH}_2 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
glutamic acid	Glu	$\begin{array}{c} \text{CH}_2-\text{CH}_2-\text{COOH} \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
glycine	Gly	$\text{H}_2\text{N}-\text{CH}_2-\text{COOH}$
histidine	His	$\begin{array}{c} \text{Imidazole ring} \\ \\ \text{CH}_2- \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$
isoleucine	Ile	$\begin{array}{c} \text{CH}_3-\text{CH}-\text{CH}_2-\text{CH}_3 \\ \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$

Name	Symbol	Structure
leucine	Leu	$ \begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_3 \\ \\ \text{CH}_2 \\ \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array} $
lysine	Lys	$ \begin{array}{c} \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{NH}_2 \\ \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array} $
methionine	Met	$ \begin{array}{c} \text{CH}_2 - \text{CH}_2 - \text{S} - \text{CH}_3 \\ \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array} $
phenylalanine	Phe	$ \begin{array}{c} \text{CH}_2 - \text{C}_6\text{H}_5 \\ \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array} $
proline	Pro	
serine	Ser	$ \begin{array}{c} \text{CH}_2 - \text{OH} \\ \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array} $
threonine	Thr	$ \begin{array}{c} \text{CH}_3 - \text{CH} - \text{OH} \\ \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array} $
tryptophan	Trp	
tyrosine	Tyr	
valine	Val	$ \begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_3 \\ \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array} $


9. Formulas of some fatty acids

Name	Formula
Lauric	$C_{11}H_{23}COOH$
Myristic	$C_{13}H_{27}COOH$
Palmitic	$C_{15}H_{31}COOH$
Palmitoleic	$C_{15}H_{29}COOH$
Stearic	$C_{17}H_{35}COOH$
Oleic	$C_{17}H_{33}COOH$
Linoleic	$C_{17}H_{31}COOH$
Linolenic	$C_{17}H_{29}COOH$
Arachidic	$C_{19}H_{39}COOH$
Arachidonic	$C_{19}H_{31}COOH$


10. Structural formulas of some important biomolecules




sucrose




glycerol




deoxyribose




adenine




guanine



cytosine



thymine



phosphate

11. Acid-base indicators

Name	pH range	Colour change		K_a
		Acid	Base	
Thymol blue	1.2–2.8	red	yellow	2×10^{-2}
Methyl orange	3.1–4.4	red	yellow	2×10^{-4}
Bromophenol blue	3.0–4.6	yellow	blue	6×10^{-5}
Methyl red	4.2–6.3	red	yellow	8×10^{-6}
Bromothymol blue	6.0–7.6	yellow	blue	1×10^{-7}
Phenol red	6.8–8.4	yellow	red	1×10^{-8}
Phenolphthalein	8.3–10.0	colourless	red	5×10^{-10}

12. Acidity constants, K_a , of some weak acids

Name	Formula	K_a
Ammonium ion	NH_4^+	5.6×10^{-10}
Benzoic	$\text{C}_6\text{H}_5\text{COOH}$	6.4×10^{-5}
Boric	H_3BO_3	5.8×10^{-10}
Ethanoic	CH_3COOH	1.7×10^{-5}
Hydrocyanic	HCN	6.3×10^{-10}
Hydrofluoric	HF	7.6×10^{-4}
Hypobromous	HOBr	2.4×10^{-9}
Hypochlorous	HOCl	2.9×10^{-8}
Lactic	$\text{HC}_3\text{H}_5\text{O}_3$	1.4×10^{-4}
Methanoic	HCOOH	1.8×10^{-4}
Nitrous	HNO_2	7.2×10^{-4}
Propanoic	$\text{C}_2\text{H}_5\text{COOH}$	1.3×10^{-5}

13. Values of molar enthalpy of combustions of some common fuels at 298 K and 101.3 kPa

Substance	Formula	State	$\Delta H_c (\text{kJ mol}^{-1})$
hydrogen	H_2	g	–286
carbon(graphite)	C	s	–394
methane	CH_4	g	–889
ethane	C_2H_6	g	–1557
propane	C_3H_8	g	–2217
butane	C_4H_{10}	g	–2874
pentane	C_5H_{12}	l	–3509
hexane	C_6H_{14}	l	–4158
octane	C_8H_{18}	l	–5464
ethene	C_2H_4	g	–1409
methanol	CH_3OH	l	–725
ethanol	$\text{C}_2\text{H}_5\text{OH}$	l	–1364
1-propanol	$\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$	l	–2016
2-propanol	$\text{CH}_3\text{CHOHCH}_3$	l	–2003
glucose	$\text{C}_6\text{H}_{12}\text{O}_6$	s	–2816