

CHEMISTRY

Units 3 & 4 – Written examination



(TSSM's 2014 trial exam updated for the current study design)

SOLUTIONS

SECTION A: Multiple-choice questions (1 mark each)

Question 1

Answer: D

Explanation:

The half equation for the reaction is $\text{Al}^{3+} + 3\text{e}^- \rightarrow \text{Al}$. Therefore the number of electrons is three times the number of aluminium atoms. $3 \times 0.12 = 0.36$

Question 2

Answer: B

Explanation:

In all redox cells oxidation occurs at the anode. In the case of electrolysis the anode is positive.

Question 3

Answer: C

Explanation:

The diagram shows magnesium being oxidized to magnesium ions and sulfur being reduced. From the electrochemical series this could occur as a spontaneous galvanic cell.

Question 4

Answer: C

Explanation:

Aluminium will not deposit from an aqueous solution hence C is correct.

Ag^+ will have the same number of mole as the electrons.

Cu^{2+} will have half the number of mole of the electrons.

Question 5

Answer: C

Explanation:

In this cell, Cl_2 will react with Mn(s) . Cl_2 forming Cl^- is a reduction reaction, hence will occur at the cathode. Mn(s) reaction is an oxidation reaction and it will occur at the anode.

Question 6

Answer: B

Explanation:

$2\text{SO}_2 + 2\text{e}^- \rightarrow \text{S}_2\text{O}_4^{2-}$ is a balanced reduction reaction. Reduction occurs at the cathode.

Question 7

Answer: D

Explanation:

In SO_2 , the sulfur is +4. In $\text{S}_2\text{O}_4^{2-}$, the charge on sulfur is +3

Question 8

Answer: A

Explanation:

Bioethanol can be produced from renewable plant matter at a sustainable rate. All other options are non-renewable.

Question 9

Answer: C

Explanation:


$$3 \times 24.8\text{L/mol} = 74.4\text{L}$$

Question 10*Answer: A**Explanation:*

At both temperatures, there are some reactants with enough energy to overcome the activation energy. The graph at 100 °C however, is higher than the graph of 50 °C at the activation energy. This means there are more particles at 100 °C with sufficient energy to react.

Question 11*Answer: D**Explanation:*

To obtain the reaction $\text{H}_2\text{O(g)} \rightarrow \text{H}_2\text{O(s)}$ the first equation supplied needs to be reversed and then the two equations added together;



ΔH will therefore be $-6 + -44 = -50 \text{ kJ mol}^{-1}$

Question 12*Answer: A**Explanation:*

Compared to the first equation, the second has been reversed then doubled. This leads to K being the reciprocal of the first value and then squared.

$$K = \frac{1}{24.2^2} = 0.00171$$

Question 13

Answer: D

Explanation:

$$1.23 \times 6.12 \times 1 = 7.52 \text{kJ}$$

$$n(\text{NH}_4\text{NO}_3) = 1/(14+4+14+48) \\ = 0.0125 \text{mol}$$

$$\Delta H = 7.52/0.0125 \\ = 601.6 \text{kJ/mol}$$

Question 14

Answer: B

Explanation:

Add a catalyst will increase the rate of reaction.

Question 15

Answer: D

Explanation:


The question is answered by looking up each hydrogen atoms shift in the data book. The alkanes have a low shift and carboxylic acids have very high shifts.

Question 16

Answer: A

Explanation:

The graph shows two reactants and one product. The product concentration grows at the same rate one of the reactants is used up, so they must share the same coefficient in the reaction ($2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$). The other reactant drops at half the rate of the first, so its coefficient must be half of the other reactant ($\text{NO} + \frac{1}{2}\text{O}_2 \rightarrow \text{NO}_2$)



Question 17

Answer: B

Explanation:

Chiral molecules contain at least one carbon with four non identical substituents. Therefore only II and IV contain chiral centres.

Question 18

Answer: B

Explanation:

They have the same percentage composition, molecular masses (137.9g/mol) and molecular formula (C_3H_7OBr) except for chirality.

Question 19

Answer: D

Explanation:

Glucose has a molecular formula of $C_6H_{12}O_6$ as does fructose. Ethanoic acid is $C_2H_4O_2$. The empirical formula of these molecules is CH_2O .

Question 20

Answer: D

Explanation:


The alkanol section of this molecule is ethanol and the acid section is methanoic acid, hence the ester is ethyl methanoate.

Question 21

Answer: C

Explanation:

Glycine has an amine group and a carboxyl group. The carboxyl group can react with a base and the amine group can react with an acid.



Question 22

Answer: A

Explanation:

Use of the Data Book shows the three amino acids as leucine, glycine and serine

Question 23

Answer: C

Explanation:

The peak corresponding to the molecular ion peak since it is the strongest and the heaviest ion.

Question 24

Answer: C

Explanation:

Infrared radiation has longer wavelength than visible light. Wavenumber is number of waves in an unit distance, since IR has long wavelength, the wavenumber becomes lower.

Question 25

Answer: C

Explanation:

Presence of a broad peak -OH at 3300cm^{-1} , absence of $\text{C}=\text{O}$ at 1700 cm^{-1} . Therefore molecule is ethanol.

Question 26

Answer: B

Explanation:

Sulfur undergoes reduction which is equation B.

Question 27

Answer: D

Explanation:


A monounsaturated molecule has one double bond. Its formula is evident as $C_nH_{2n-2}O_2$

Question 28

Answer: B

Explanation:

Glycerol has 4 different environments, marked with an *



Question 29

Answer: D

Explanation:

$$n(CH_4) = \frac{1.5}{40.16} = 0.9375 \text{ mol}$$

$$\begin{aligned} \Delta H &= 83 / 0.09375 \\ &= - 885.3 \text{ KJ/mol} \end{aligned}$$

Question 30

Answer: B

Explanation:

$$n(C) = \frac{2}{12} = 0.166$$


$$n(H) = 10/4 \ n(C) = 10 \times 0.166/4 = 0.417$$

$$mass = 0.417 \text{ g}$$

SECTION B : Short-answer questions

Question 1 (10 marks)

- | | | | | | |
|-----------|---------------------------------|--------------------|---------|--------|----------------------|
| a. | i. Example: | glucose | sucrose | starch | 1 mark |
| | ii. Molecular formula: | $C_6H_{12}O_6$ | * | : | $C_{12}H_{22}O_{11}$ |
| | | | | | * |
| | | | | | |
| | iii. Structural isomer : | fructose/galactose | | | 2 marks |
| | | | | | 1 mark |
| b. | | | | | 1 mark |



- c. i. pepsin is close to its optimum reaction rate* while trypsin does not function at this pH* 2 marks
- ii. The tertiary and secondary structures of the enzyme are destroyed. The enzyme loses its particular 3-D shape. Its primary structure however, is still intact. 1 mark
- iii. The mass of the individual amino acids will be greater than the mass of the protein.* When hydrolysis occurs, water is added back onto the molecules. 19999 water molecules would be needed to allow hydrolysis * 2 marks


Question 2 (10 marks)

a


$$\text{O}_2(\text{g}) \quad 3 \underline{\hspace{2cm}}$$
H₂O (l) 4
$$\text{CO}_2(\text{g}) \quad \underline{\quad 2 \quad}$$

CH₃OH (l) 2

- b.** Full equation: $2\text{CH}_3\text{OH}(\text{g}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 4\text{H}_2\text{O}(\text{l})$ $E^\circ = +0.80 \text{ V}$ 1 mark



1 mark



1 mark


$$E^\circ = (1.23 \times 3 - 0.80) / 2$$

$$= 1.445 \text{ V}$$

1 mark


Question 3 (10 marks)

a.



i. See graph: H_2 drops at twice the rate of CO from balanced equation ** 2 marks

ii. Methanol starts at zero and rises at the same rate CO drops – dashed line below




1 mark

b. i. $K = \frac{[\text{CH}_3\text{OH}]}{[\text{CO}][\text{H}_2]^2}$ * 2 marks

$$= \frac{0.3}{0.3 \times 0.4^2} = 6.25 \text{ M}^{-2} *$$

ii. $c = 0.3 \Rightarrow n = c \times V = 0.3 \times 2 = 0.6 \text{ mol}$ 1 mark

- c. i. each concentration is halved. Keep in mind that the concentration drop is not the same for each chemical – the graph should show the values halving. 2 marks



- ii. system will move to the left, creating more particles*. The concentration of carbon monoxide and hydrogen will increase* 2 marks

Question 4 (6 marks)

- a. i. methyl propanoate 1 mark

- ii. $\text{C}_2\text{H}_4\text{O}$ 1 mark

- b. i. $\text{C}_4\text{H}_8\text{O}_2(\text{l}) + \text{H}_2\text{O}(\text{l}) \rightarrow \text{CH}_4\text{O}(\text{aq}) + \text{C}_3\text{H}_6\text{O}_2(\text{aq})$ 1 mark

- ii. $2\text{CH}_3\text{OH}(\text{l}) + 3\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + 4\text{H}_2\text{O}(\text{l})$ 1 mark

- iii. ΔH for methanol is $-725 \text{ kJ mol}^{-1} *$
 ΔH per g = $725/32 = 22.7 \text{ kJ g}^{-1} *$

2 marks

Question 5 (7 marks)**a.** Cell contents: KCl(aq) *

3 marks

Anode half equation: $2\text{H}_2\text{O}(\text{l}) \rightarrow \text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- *$ Cathode half equation: $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq}) *$ **b.** Metal will be deposited in the CuCl₂ solution. *

$$Q = It = 2.55 \times 20 \times 60 = 3060 \text{ C} *$$

$$n(\text{e}) = \frac{3060}{96500} = 0.0317 \text{ mol} *$$

$$n(\text{Cu}) = \frac{1}{2} n(\text{e}) = 0.0159 \text{ mol}$$

$$\text{mass} = 0.0159 \times 63.5 = 1.00 \text{ g} *$$

4 marks

Question 6 (9 marks)**a. i.** $\text{Pb}(\text{s}) + \text{PbO}_2(\text{s}) + 2\text{H}_2\text{SO}_4(\text{aq}) \rightarrow 2\text{PbSO}_4(\text{s}) + 2\text{H}_2\text{O}(\text{l})$ 1 mark**ii.** As this cell discharges the sulfuric acid reacts. Therefore the pH will rise as the cell discharges. 1 mark**iii.** A secondary cell can be recharged but a primary cell is disposed of once flat.

1 mark

iv. Lead is very heavy hence the vehicle is heavy*. Lead is neither abundant nor completely safe to handle.*

2 marks

b. $\text{PbSO}_4(\text{s}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow \text{PbO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) + 2\text{e}^-$ 1 mark**c. i.** anode: $\text{Na} \rightarrow \text{Na}^+ + \text{e}^-$
cathode: $3\text{S} + 2\text{e}^- \rightarrow \text{S}_3^{2-}$

2 marks

ii. Sodium reacts very vigorously with water, especially at 300 °C.

1 mark

Question 7 (12 marks)


a. i. Mass of chlorine = $3.6 - 1.65 - 0.322 = 1.628$ g 1 mark

ii. $\frac{1.65}{12} : \frac{0.322}{1} : \frac{1.628}{35.5} = 0.138 : 0.322 : 0.0459 \times = 3 : 7 : 1 = \text{C}_3\text{H}_7\text{Cl}$ 2 marks

b. i. Chlorine has two isotopes, ^{35}Cl and ^{37}Cl . There is a peak present for each isotope. 1 mark

ii. 63 is 15 less than 78. This is probably the removal of a methyl group, CH_3 . 1 mark


iii. Molecular formula of the compound matches empirical formula $\text{C}_3\text{H}_7\text{Cl}$ 1 mark




Isomer 1: 1-chloropropane**

Isomer 2: 2-chloropropane**

d. molecule is 2-chloropropane* as it has two different hydrogen environments. One environment is shown on the spectra as a doublet (one H neighbour) and the other as a septet (six H neighbours).*. 2 marks



Question 8 (9 marks)

- i.** Propene is a hydrocarbon. It has no significant dipoles to lead to any polarity. It is non-polar, hence low in solubility in water. 1 mark
 - ii.** Again, the lack of dipoles means that the forces between molecules are dispersion forces only. They are weak and the boiling point is low. 1 mark
- b.** See diagram 1 mark
- c.**
- i.** See diagram 1 mark
 - ii.** A 'bromine test' is a test for whether a molecule is saturated or not*. Bromine is brown in colour. If a double or triple bond is present, it will react and the brown colour disappears.* 2 marks
- d.**
- i.** See diagram. (could also be 2-propanol)
 - ii.** Several possible answers. Molecule B will react with dichromate to form a carboxylic acid. Molecule B will have an absorption band around 3200 cm⁻¹ where the -O-H absorbs. (1 mark method, 1 mark why method works) 2 marks

Question 9 (7 marks)

a. HMe and Me⁻ have different colours, therefore the position of equilibrium can be monitored by the colour. 1 mark

b. [H₃O⁺] is increased*. This favours the back reaction and produces the red colour*. 2 marks

c. The OH⁻ reacts with H₃O⁺. This lowers the [H₃O⁺]*. This favours the forward reaction and produces the orange colour.* 2 marks

d. $K_a = \frac{[\text{H}_3\text{O}^+][\text{Me}^-]}{[\text{HMe}]} = 2 \times 10^{-4}$ * 2 marks

Since [HMe] = [Me⁻] at transition, $K_a = 2 \times 10^{-4}$

pH = -log(2 × 10⁻⁴) = 3.7 *

Question 10 (7 marks)

a. i. $n(\text{ethanol}) = \frac{1.5}{46} = 0.0326 \text{ mol}$ * 2 marks


$E = 1364 \times 0.0326 = 44.5 \text{ kJ}$ *

ii. $CF = \frac{\text{energy}}{\Delta T} = \frac{44.5}{8.9} = 5.00 \text{ kJ } ^\circ\text{C}^{-1}$ 1 mark

b. $E = CF \times \Delta T = 5 \times 0.87 = 4.35 \text{ kJ}$ *

$n(\text{Na}) = \frac{0.145}{23} = 0.00630 \text{ mol}$ *


Energy/mole = $\frac{4.35}{0.00630} = 690 \text{ kJ mol}^{-1}$



$\Delta H = -1380 \text{ kJ mol}^{-1}$ *


4 marks

Question 11 (6 marks)



a. In order: hydroxyl, amine and carboxyl 3 marks

b. i. 1 mark



ii. What other molecule is formed when paracetamol is formed? Water 1 mark

iii. Benzene: molecular formula is C_6H_6 1 mark